Mitigation of COVID-19 in Workplaces: When Push comes to Shove in Offshore Oil Operation.

Abstract

Introduction: As COVID-19 pandemic is a novel disease with no prior antidote or proven preventive measures;

global response in occupational settings significantly defaulted to remote working otherwise referred to as

'Working from Home - WFH' or 'Remote Working'. While remote working favoured some industries and

businesses, it was impractical for industries with processes requiring physical contact to operate such as in the

offshore oil and gas industry. This meant workers in the offshore industry must dare the risk of exposure to

access offshore work environment thereby increasing their exposure to the dreaded and incurable virus. The aim

of this study was to explore and present effective response strategies to COVID-19 pandemic in offshore work

environment when exposure becomes inevitable. The method adopted was an observational descriptive study

(ODS) using the ad libitum sampling technique. A Walk-Through Survey (WTS) was conducted to observe and

discuss the measures adopted by a representative offshore oil company in the Gulf of Guinea. The result showed

significant potentials for increased transmission in the offshore oil industry. Multiple contact points and surfaces

in offshore facilities, shared amenities and tools, and constrained proximity of working position constitute

significant risk factors in offshore oil and gas facilities. Mitigation measures could be categories as general

measures; pre-embarkation strategies; transit strategies; aboard strategies; and remedial strategies. Combined

measures have been found a workable and effective model to mitigate the spread and transmissibility; the

eminence of distancing, serial testing, personal and community hygiene, and vaccination was established as a

more effective model.

Keywords: Work-From-Home, Severe Acute Respiratory Syndrome-Coronavirus-2, vaccination, transmission,

COVID-19

Introduction

Coronavirus disease 2019 (COVID-19 or C-19) outbreak was initially reported to have occurred in Wuhan,

China on the 31st of December 20191,2 as a novel type of respiratory tract infection. This was corroborated by

the World Health Organisation (WHO) as a coronavirus-related pneumonia on the 9th of January 2020 and later

confirmed as a pandemic on the 11th of March 2020.34 It is known to be caused by the deadly Severe Acute

Respiratory Syndrome-Coronavirus-2 (SARS-CoV-2) which is transmitted as respiratory droplets through

1

sneezing, coughing, inhalation of infected airborne droplet, and contact with infected surfaces or objects. Worldwide, C-19 pandemic negatively impacted operations of industries, commerce, and trade with the offshore oil industry amongst the worst hit. While the initial response strategy by governments was enforced lockdowns and varying degrees of restrictions of movements; industries and organisations defaulted to working remotely in a strategy dubbed 'Work from Home (WFH)' or 'Remote Working'.

Key learning points

What is already known about this subject:

- Offshore oil workers are constrained to continue offshore commute and work despite uptick in the incidence rate of COVID-19 (C-19).
- Surging number of C-19 cases impacted the industry in several fronts.
- Different measures have been adopted by different oil and gas players with no universally acceptable model.

What this study adds:

- We now know the critical health risk factors in the offshore oil and gas industry.
- A bouquet of proven measures and exemplar model has been synthesised

What impact this may have on practice or policy:

- Risk management will factor the critical risk factors into C-19 response strategy in the offshore industry.
- Bespoke models should be generated from the collage of proven response measures.

The WFH response strategy meant staff would work remotely from home via online platforms and telecommunication systems. While the remote approach worked seamlessly for certain industries as a strategy for meetings, engagement, trainings, communications, reviews, and other contactless work; industries with activities and field work requiring manual operations were handicapped as they required specialised skill pool for operations and maintenance at site. Some of the remotely operated offshore facilities encountered telecommunication challenges at the onset of COVID-19 due to inadequate remote working technologies, insufficient internet bandwidths. One of the global industries that could not completely operate remotely was the offshore petroleum industry because of being largely and remotely field base, and heavy dependence on industrial processes requiring human operation of technology hardware's.

Comment [AS1]: Please write down the table number and title.

The offshore petroleum industry was hence classified among essential industries granted waivers by governments. This meant workers would have to commute and work during the pandemic as the industry was found a major economic artery for several economies globally, thus portend significant risk exposure necessitating a robust risk management strategy.

While the health and safety of workers took severe hit; production and warfare of economies plummeted with several cases of mortality and morbidity reported⁷. In Mexico, Pemex (Petroleos Mexicanos), the state-owned oil and gas exploration and production company reported the death of 314 direct staff and 7 contractors from start of pandemic to September 2020.8 Pemex is reported to have over 240 offshore platforms in the Gulf of Mexico with an average of 200 -300 personnel working and living in clustered and shared spaces. 9 Measures deployed by Pemex include enforced social distancing, encouraged self-reporting, de-manning of platforms, sanitization of work and living spaces, pre-embarkation temperature checks, rapid testing, early evacuation of suspected ill workers. In Brazil, several COVID-19 outbreaks have been severally reported in offshore platforms. The nexus of oil and gas operations in the United Kingdom, the North Sea, witnessed upsurge in aeromedical evacuations from several offshore platforms due to workers testing positive to COVID-19.10 The oil workers federation of Brazil (FUP) reported 8,500 oil and gas workers in Brazil have tested positive to COVID-19 by December 31, 2020. Petrobras (Brazil's national oil company) had performed 450,000 COVID-19 tests; 4,030 workers have tested positive with 3 deaths by January 11, 2021. Other measures employed by Petrobras beside testing including a buffer waiting period of 14 days (quasi quarantine) before embarkation on any offshore platform, and compulsory training. In Norway, cases of COVID-19 infections have been reported on offshore platforms. 10,111 Overall, the cumulative psychological effects of layoffs, social restrictions and lockdowns, inevitable work-related exposures, incidences of increasing COVID-19 cases and deaths, sliding economic forecast for the upstream petroleum industry, presented a perfect storm for mental health challenges for the offshore cohorts of workers. Beside the health and safety implications of the pandemic, global business outlook was heavily impacted. The industry witnessed a slump in crude demand, forced decline in crude oil production, crude oil stockpiles, debt rise, loss in share value, decrease demand for end products, decrease in asset value, decreased or near zero investment, lay-offs, increased divestment, difficulties in logistics, halted business travels, deferred for among others (see Figure 1 and Figure 2).

Given that the offshore petroleum industry is a significant national cash cow and continual working during the pandemic is inevitable; a robust risk mitigation strategy will be required to mitigate the impacts on workers'

Comment [AS2]: Adjust the intended image description's positioning in relation to the desired sentence explanation.

health and productivity. The need for mitigation is further heightened by the reported uptick in morbidity and mortality occurring in upstream oil industries across the world.




Figure 1: Collapse of Oil Price in 2020¹⁰




Figure 2 Oil Majors 2020 capex cuts¹¹

The novelty of the COVID-19 pandemics further necessitates sharing generalisable mitigation practices that could be adapted by other players in industry. The aim and objective of this study is to identify potential risk factors in the offshore oil and gas industry and present workable measures that could mitigate exposures and protect the health of at-risk workers.

Methods

This is an observational descriptive study (ODS) conducted using the ad libitum sampling technique. A Walk-Through Survey (WTS) was undertaken to conduct the observational study between January 2020 and April 2021. The WTS involved direct observation of COVID-19 mitigation processes and review of procedures adopted by an offshore petroleum industry in the Gulf of Guinea. WTS is a research process used to obtain primary information about a workplace through direct observation and discussion. ^{12,13,14}

The study setting was an offshore oil and gas servicing facility in the Gulf of Guinea with over 1500 workforce. The Observational Descriptive Study (ODS) technique was adopted in this study because of its usefulness and adequacy in accurate description of observed processes and procedure. It is known to be accurate, factual, verified source of primary data. Technique constitutes 70% -80% of research in scientific journals. It is the commonly used research technique used in scientific research to observe and was achieved in this study through filed visits, note taking, discussion, meaning making and reflection. Permission was granted by the process owners for educational purposes; however, the organisation requested to be de-identified.

Results

The walk-through survey (WTS) revealed a study setting replete with COVID risk factors and mitigation measures that has been grouped into general, pre-embarkation, transit, onboard and remedial strategies. Commute to offshore location involves staff reporting at the local airstrip from their respective homes to board flight to offshore locations. At the airstrip workers will undergo check-in protocol, safety briefing and board helicopters. There are usually 7-9 passengers onboard the 1-hour flights to the offshore platform. On arrival and disembarkation, workers undergo briefing and induction about the facility and disperse to their respective rooms. Other shared and communal amenities include the toilets, large restaurant, library, meeting rooms, banking hall and workshops. Work cycle is 28 days off and on rotational cycle. Besides the several machinery units are the living areas which accommodate 4 persons in each room of two double bunk bed. Other communal and shared areas offshore include the mess, meeting rooms, open-plan offices, gymnasium, and workshops. Categories of workers include the operations, maintenance, and support staff. While operation staff undertake activities needed to produce oil and gas, maintenance staff undertakes corrective and preventive maintenances while the support staff cater for welfare and material management.

The WTS revealed significant health risk factors associated with working in an offshore oil and gas facility.

These include use of shared amenities including mess, open plan offices, accommodation (rooms), meeting spaces or concourse, recreational spaces, transport medium, computer stations, communication facilities (phone

Comment [AS3]: Explain what was observed and, if an interview was conducted, detail the questions asked of the respondent. Additionally, convey the respondent's identity or the respondent's criteria.

& handheld radios), laundry services, hand tools and equipment; clustered layout of machinery, office settings and accommodation; multiple contact points and surfaces including operational controls (buttons, levers, wheel, jockeys); compulsory handling of rails of stairways, walkways, gangways and door handles; interconnected ventilation ducts and multinational workforce. These risk factors could potentially trap and foster temporary storage of COVID-19 from an infected worker and potentially facilitate its transmission to unaffected workers.

General strategies include measure's workers observed when not at work in their personal spaces. These include regular handwashing with soap and free flowing water, use of sanitisers and mask, social distancing of at least 1.5 meters, compliance with lockdowns and restriction by jurisdictions, avoidance of potentially super spreader events, hospital call when respiratory and symptoms are developed.

Pre-embarkation strategies involved measures undertaken by the company and staff before embarking on flight to offshore locations. These measures include completion of online medical questionnaire form eight days prior to travelling, detailing their present state of health with regards to presence or absence of respiratory tract infection, sore throat, cough, fever, a recent travel history and close relation of any febrile or ill person suspected of having symptoms. A normal case would however visit a pre-travel quarantine centre in a selected, prepared, and secured hotels not available to the public. While completion of a normal questionnaire will automatically activate chip in staff identity card to access quarantine centre; the presence of symptoms will however deny access to the premises but will trigger the C-19 remedial response strategy by referring the person to specialised hospital equipped to manage respiratory cases including C-19. At the quarantine centre, a PCR C-19 test will be conducted before being accommodated in the centre for 7 -days. A negative test result is followed by a confirmatory test on the fourth day of quarantine. Staff with two negative tests are transported in a bus to the airstrip for onward trip to offshore location; the bus and helicopters are periodically decontaminated.

Transit strategies involved measures committed to mitigating exposures during commute in company buses from quarantine centres to the airstrip and onboard flight from the airstrip to offshore location. Strategies include a pre-boarding briefing on covid safe measures, temperature scan, hand sanitization, mask, social distance and helicopter. On alighting, the bus and helicopter undergo light cleaning with deep cleaning done weekly.

Aboard strategies involved measures undertaken to mitigate exposures while on-board the offshore facility.

These include, arrival temperature check with the telethermographic system to assess temperature on

disembarkation, safety induction which includes safe COVID-19 measures, departure temperature check, mandatory use of N95 mask, placement of disinfectant wipes and sanitizer stations, use of sanitizers after touching 'high-touch-surfaces', arrangement of workstation desks to be 1.5 feet apart from each other in the offices, weekly cleaning, replacement of air filters in ventilation systems, floor markers of 6-feet apart to maintain physical distancing in common areas, two hourly cleaning of high touch surfaces (doorknobs, lift call buttons etc) and conveniences and twice daily disinfection of employee workstation.

Remedial strategies refer to measures deployed when anyone test positive to PCR test or presents suspicious symptoms. The person is isolated in a secured room, medic and attendants wear disposable hazmat suit with air supplied respirator. Medical screening is commenced, an antigen and PCR C-19 test is performed, and treatment is started after a positive confirmation in consultation with supervising physician. As soon as reasonably practicable, the individual is evacuated. Once onshore, the patient is transported to an appropriate medical facility. All environmental cleaning and disinfection procedures are followed consistently and correctly. Increased diligence in sanitary measures and infection control is implemented once a possible case of a communicable disease is identified. Identification of close contacts who may be infected or other individuals at risk for infection commence. The appropriate public health authority is notified and with advice on further actions.

DISCUSSION

Our study showed that COVID-19 poses significant threat to the health of offshore oil workers as evidenced by surging cases of positive cases in different scenarios, multiple health risk factors and the need for a collage of mitigation measures. The strength of this study lies in the validity of using a tested process and study setting. The non-availability of incidence rate before and after the study potentially weakens measurement of impact of the mitigation measures. The quest for crude oil in offshore locations pitch the workforce against the dreaded virus as offshore facilities are remotely located requiring exposure to several C-19 risk factors. While robust measures were mobilised to mitigate the exposures in this facility; the study by Wilbanks, Abulhassan & Kilpatrick revealed that additional measures should be included as shown in Table 1. Although there are similar measures in both models, the authors differ in opinion on 'no air travel or hotel and opening of all doors to avoid contact'. Authors of this study posit that with suitable and adequate measures put in place air travel and closed doors could still be achieved. Measures such as frequent cleaning of door handles and other touch surfaces could

mitigate transmission by contact. Contrary to the two models, the BOHS strategy (see Table 2) opined the importance of Covid Risk Assessment (CRA) which assesses similar exposure groups in occupational settings, exposure scenarios or events, control banding and control options. The response approach should target the pathogen at the source, pathway, and receptor point. While the model in this study were seemingly robust, it would have been more potent if there was strategic engagement at the source, pathway and receptor or receiver end.

Given the itinerant and multicultural demography of the offshore workforce, activities in the industry is buoyed by significant intranational and international business travels making the sector vulnerable to transmission of the virus. The industry being a major employer of labour, engages staff, contractors and sub-contractors running in thousands especially multinationals upstream organisations. There exists significant potential for transmission within offshore facilities given the bunched-up layout with linear ventilation system. It possesses several shared features that portend the risk of C-19 transmission such as control rooms, training rooms, public conveniences, meeting rooms, handles of entrances and exits, group safety and team meetings, walkways, and stairways less than 1.5m width, shared ventilation systems (Heating Ventilation Air Conditioning – HVAC system). Potential activities include crew change activities, meetings, trainings, eating, handling, and touching of surfaces, ventilation within open plan offices, field accommodations and use of handled tools, controls, and surfaces. Risk factors above supports and corroborates the identification of the Oil and Gas industry as a major frontline sector requiring response in the United State OF America³. More importantly, the following category of workers constitutes a high-risk group amongst offshore workers, namely the cleaners, laundry workers, logistic personnel, utility workers, stewards, security, medical, team leaders and emergency response team workers.

Screening	Worker	Barriers	Touchable surfaces	Cleaning/	Communication	Reporting and quarantine
	interaction			disinfecting		
All access control	 Spreading shifts 	•Physical	• Open	•Trained and	Awareness training	• Procedures
point	•Visitor access	transparent	doors to reduce	adequate	for all personnel and	Self-quarantine and
Screening	controls	barriers in	contact	cleaning personnel	visitors	return-to-work guidelines
Temperature	 Online meetings 	open plan offices	 Disposable or 	 Automatic hand 	prior to entry	published
checks	•Work process	•Face mask	Single use	sanitizer stations	 Daily and periodic 	 Engagement with union
 Identity cards access 	modification.	required	packaging	•Easy access	employee	On COVID-19 procedures
linked to	 Telehealth 	unless stationary	 Installation of 	disinfectant wipes	communication	 Mandatory compensatory
screenings	 Embargo on air, 	and	hands-free or foot	stations	Additional	leave if
Infrared	sea travel and hotel	social distancing	door	•Electrostatic	newsletters, electronic	confirmed case or
scanning,	Accommodation	can	 Electronic form 	cleaning	board messaging	exposed to confirmed
cameras, kiosks		be achieved	screening	•Ultraviolet	brochures, postings,	case

 Table 1: Response Strategies

Comment [AS4]: It should be placed above the table section.

Give an explanation for the table you've been given.

Self-screening	•Social distancing	•Hands-free testing	sterilizer at	handouts	Periodic COVID-19 drills
• Daily	Markers on the		high-traffic areas	 Precautionary signage 	Contact tracing
questionnaires	floor		 Post-cleaning 	• Inter and Intra	procedures
Developed	Face shields		swab	departmental	 Configuring badge
organisations app	•Partitions		(ATP) testing	pandemic response	system for use in contact
• PCR testing			•Portable or mobile	marshals	tracing and QR-coding
Electronic door			disinfection		
screening			misting devices		

Source: Adapted [4]

Control Ordinar							
Control Options							
Source	Pathway	Receptor					
Infection control measures and	Cleaned and maintained Local	Powered Air Purifying Respirator					
patient isolation, restricted access,	Exhaust Ventilation (LEV),	(PAPR), otherwise minimum					
regular surface disinfection with	General ventilation, regular	Filtering Face Piece (FFP3) and					
acceptable chemicals. Visor or	surface disinfection	visor, gown, gloves and/or					
face covering on medical staff and		personal hygiene - hand					
patient		washing/hand sanitizing.					
Isolation of patient, restricted	Barrier or enclosure. General	Minimum FFP3 and visor, gown,					
access, regular surface	ventilation, regular surface	gloves and/or hygiene - hand					
disinfection. Visor or face	disinfection	washing/hand sanitizing.					
covering on both staff and patient							
Distancing, hand washing,	Barriers, regular surface	FFP2 should be considered for					
sanitisation.	disinfection of touch points and	prolonged contact, otherwise fluid					
	surfaces, one-way systems as far	resistant masks, visor, gloves and					
	as reasonably practicable, general	or hygiene - hand access.					
	ventilation, avoid handling cash.						
Distancing, frequent hand washing	Barriers, regular surface	Visor or safety spectacles & fluid					
and sanitisation.	disinfection of touch points and	resistant mask and hygiene - hand					
	surfaces.	washing and sanitizing					
Distancing, frequent hand washing	Regular surface disinfection of	Visor or safety spectacles and					
and sanitisation	touch points and surfaces.	fluid resistant mask and hand					
		washing and sanitization.					
Social distancing	Surface disinfection of touch	Hand washing and sanitization.					
	points and surfaces.						

Source: Adapted from British Occupational Hygiene Society¹⁶

While above strategies may have contributed to slowing down the transmission of the virus as evidenced by low incidence rate in this organisation, the advent of the Delta strain might after the period under consideration may have increased the number of infected workers given the significant potentials, clustered layout in offshore environment and logistic medium. Inclusion of vaccination into the bouquet of strategies will drastically reduce transmission, mitigate exposure, improve resilience, and enhance productivity. It is however important to align the strategies with an adapted hierarchy of controls¹⁷ (see figure 3).




Figure 3: Hierarchy of Controls¹⁷

Conclusion

In conclusion, we found that there are significant potentials for transmission and spread of C-19 in the offshore oil industry occasioned by the travel pattern of the workforce, commute method, design of offshore oil facility, presence of several shared facility and abundant touch points. Combination of the response strategies will be a potent approach to mitigating exposure and transmission of COVID-19 in an offshore facility and operations. The strength of this study lies in the validity of the presented measures which have been tested and proven. Limitation lies in the short duration of the study and inability to validate the measures for 1 year to authenticate in the long term.

Comment [AS6]: Is there any evidence for this? If you compare it to the findings of your research, is it conclusive? If not, why not?

Comment [AS7]: There is no evidence to back this up. Please give proof with valid explanations and data.

References

- 1. Osakwe K, Ola K, Omotosho P. Contactless Academia The Case for Automated Essay Scoring (AES) System in COVID 19 Pandemic. *CJAST* 2021; 40:17-29.
- 2. Osakwe Adakporia K, Menkiti O, Ukaegbu K. Three-month Epidemiological Trend of Covid-19 in Nigeria Rapid Review. *CJAST* 2020: 39:9-18.
- Krebs CC. Memorandum of Identification of Essential Critical Infrastructure Workers during COVID-19
 Response. US Department of Homeland Security 2020.
- 4. Wilbanks DW, Abulhassan Y, Kilpatrick RS. COVID-19 Early Planning, Response & Lessons in OSH. *Professional Safety* 2021; 66:32-37.
- 5. Kumar A, Singh R, Kaur J, Pandey S, Sharma V, Thakur L, Sati S, Mani S, Asthana S, Sharma TK, Chaudhuri S, Bhattacharyya S, Kumar N. Wuhan to World: The COVID-19 Pandemic. *Frontiers in cellular and infection microbiology*, 2021;11:596201
- University of Rochester. Guideline for De-identification of Data for Publishing. Research Subject Review Board 2018.

https://www.rochester.edu/ohsp/documents/ohsp/pdf/policiesAndGuidance/901b GDL Deidentification Data
Publishing.pdf (accessed 5th February 2022).

- 7. Norouzi N. (2021). Post-COVID-19 and globalization of oil and natural gas trade: Challenges, opportunities, lessons, regulations, and strategies. *International journal of energy research*, 10.1002/er.6762. Advance online publication. https://doi.org/10.1002/er.6762
- 8. Bloomberg https://www.bloomberg.com/news/features/2020-09-10/oil-producer-pemex-has-the-most-covid-19-deaths-of-any-company-in-the-world 2020 (accessed 5th February 2022)
- 9. Bnamericas. COVID-19 cases up in Brazil's oil and gas sector 2021.

https://www.bnamericas.com/en/news/covid-19-cases-up-in-brazils-oil-and-gas-sector (accessed 5th February 2022).

- 10. Offshore technology. How coronavirus has spread through North Sea platforms 2020. https://www.offshore-technology.com/features/coronavirus-north-sea-infections-evacuations/ (accessed 5th February 2022).
- 11. Yu K. Multiple COVID-19 shocks mean no respite for oil and gas sector in near future 2020.https://www.maplecroft.com/insights/analysis/multiple-covid-19-shocks-mean-no-respite-for-oil-and-gas-sector-in-near-future/ (accessed 5th February 2022).

- 12. Hiller J. Exxon lops 30% off 2020 spending, deeper and later than rivals.2020 Reuters https://www.reuters.com/article/us-global-oil-exxon-mobil/exxon-lops-30-off-2020-spending-deeper-and-later-than-rivals-idUKKBN21P1II (accessed 5th February 2022).
- 13. Moen EB (2014). The Importance of Walk-Through Surveys at Workplaces. Occup Med Health Aff 2014; 2:171.
- 14. Quizon RR, Hernandez PMR, Lacsamana GC, Remaneses JI. Identification of Occupational Hazards by Walk-Through Surveys Among Selected Public Hospitals in the Philippines. Phil J Health Res Dev. 2015;19 (3):41-47.
- 15. Manterola C, Otzen T. Checklist for reporting results using observational descriptive studies as research designs. The MInCir initiative. *Int. J. Morphol* 2017; 35(1):72-76.
- 16. British Occupational Hygiene Society -BOHS. Covid 19 Control Measures Risk Matrix. 2020; Version 1. https://mk0bohsx5kak7rlajjs.kinstacdn.com/app/uploads/2020/10/BOHS-Covid-19-Control-Measures-Risk-Matrix-Version-1.0 23.07-1.pdf (accessed 5th February 2022).
- 17. Dehghani F, Omidi F, Yousefinejad S, Taheri E. The hierarchy of preventive measures to protect workers against the COVID-19 pandemic: A review. Work (Reading, Mass.), 2020; 67(4), 771–777.