

# Agronomic Characterization of Bambara groundnuts accessions (*Vigna subterranea* (L.)) on Sudano-guinean zone, Adamawa-Cameroon

## ABSTRACT

**Aims:** ~~this~~ Study aims to identify agronomic traits responsible ~~of~~ ~~for~~ the variation between 20 accessions ~~of Bambara groundnut~~ in order to retain adapted genotype on pedoclimatic conditions of the Sudano-sahelean area of Cameroon.

**Place and Duration of Study:** This study was carried in Dang on Sudano-~~guinean~~ ~~Guinean~~ zone of Cameroon in order to identify agronomic traits responsible for variation among 20 accessions of Bambara groundnuts.

**Methodology:** The experimental design in field is a block completely randomized with 3 repetitions. Eleven quantitative characters selected among the describer of Bambara groundnut have been evaluated by the multivariate analysis.

**Results:** The analysis of variance shows a significant difference for ten of the eleven studied characters among eleven characters studied which show an important variability between these characters. Seven characters are correlated on two prime of principles components which explaining 44.32 % of total variability. Dendrogram rising of hierarchical classification permit to differentiate 5 groups of morphological diversity. The fifth group gathers the earliest varieties with a time of 121 days maturity. Highly positive correlation is ( $r=0.72$ ). Characters analyzed may constitute some base of criterion to differentiate the varieties from others regions and must be used on the program of improvement varieties of Bambara groundnut in Cameroon.

**Conclusion:** In all of ~~the~~ 11 characters evaluated in this study, only one of them that not permit to differentiate the accessions. Important variability between ~~theses-these~~ characters ~~was-were~~ observed. ~~Ri~~ ~~The rising~~ rising of hierarchical classification showed one of important agronomic diversity and regroup the accessions in 5 groups.

**Keywords:** Bambara groundnut, Analysis of principles components, rising of hierarchical classification, correlation.

## 1. INTRODUCTION

For their development, several African countries massively investigated ~~in~~ industrial crops to the detriment of food crops of which some of them are abandoned and threatened as

**Comment [QA1]:** Please add months and year with complete details

disappearance (Aké, 1988). The majority of annual culture has low yields due to climate change, but there are adapted on their middle (Altieri, 1986).

**Comment [QA2]:** Please used most latest citations at least from 2010 and onwards

Bambara groundnut (*Voandzeia subterrannea* (L.)) is the third food leguminous most important on terms of production and consummation in Africa, after peanut (*Arachis hypogaeae* (L.)) and Cowpea (*Vigna unguiculata* (L.)) (Linnemann, 1992; Howell, 1994). Their seeds are rich in calories, mineral elements, vitamin and proteins (Amarteifio *et al.*, 2006; Sévérin and Yao, 2011; Ndiang *et al.*, 2012). ~~She-It~~ have therapeutic property ~~that for and hence is popular among the~~ rural populations (Nacoulma, 1996; Sévérin and Yao, 2011). Bambara groundnut ~~have positive impact on~~ ~~contributes~~ on soil ~~fertilization-fertility by due of~~ their ability to fix ~~atmospheric~~ nitrogen and ~~amelioratee enhance~~ yield of plants around 350 to 800 kg/ha in region where soil is poor and rainfall is relatively low (Bonny and Djè, 2011; Touré *et al.*, 2013). This stability of production could be due to their adaptation traits on tropical climate and genetic diversity of seeds sowed (Azam-Ali *et al.*, 2001). Some morphotypes resist well on insect ~~attackattacks~~, diseases and on critical conditions of dryness (Mungate, 1997; Touré *et al.*, 2013). According to their importance, Bambara groundnut is considered like minor culture and not used ~~on-in~~ varietal selection programs (Zerihun, 2009). However, these cultures have several potentialities for genetic amelioration according to their diversity founded on many accessions collected ~~around-from the~~ country (Basu *et al.*, 2004; Ntundu *et al.*, 2004; Djè *et al.*, 2005; Bonny et Djè, 2011).

**Formatted:** Font: Italic

**Formatted:** Font: Italic

**Formatted:** Font: Italic

**Formatted:** Font: Italic

**Formatted:** Font: Italic

**Formatted:** Font: Italic

In Sudano-sahelean area, the production of seeds is relatively low due to ~~the decrease of poor~~ soil ~~ferilityfertility~~ (Minader, 2012). However, ~~in-from a-an~~ agronomic context, acknowledge of genetical mechanisms could help to ~~know-develop~~ strategies of selection in order to retain high genotype adapted on pedo-climatic conditions of Sudano-sahelean zone. Many accessions of Bambara groundnut collected on different area of Cameroon have different characters (Pasquet et Fotso, 1991; Ndiang *et al.*, 2012; Sobda *et al.*, 2013). Research or new knowledge on variety on high yield of Bambara groundnut is a principal objective of selection. General objective of this research is to identify agronomic traits responsible of the variation between 20 accessions in order to retain adapted genotype on pedoclimatic conditions of the Sudano-~~sahelean-Sahelian~~ area of Cameroon.

## I- MATERIALS AND METHODS

### I.1. DESCRIPTION OF STUDY

Dominant trait of this area (Sudano-Sudano-sahelean climate type) of Cameroon is a massif of Adamawa with ~~a~~ mountainous arch which separate northern and southern part of country. Adamawa plateau is dominated with a plain of Bénoué in the north and in west with Nigeria. This region extends to south-west by the high volcanic mountain with the altitude of 2460 m (Rippstein, 1985).

### I.2. EXPERIMENTAL SITE

The study was conducted in Dang situated at 700 m of Campus University of Ngaoundere at Bini-Dang, village situated at 15 kilometers of the town of Ngaoundere (1113 m of altitude with latitude of 7.28° North and longitude of 13.34° East). Climatic conditions of this site are

characteristic of Sudano-guinnean zone. Geographic localization of this experimental site is 13°34'238" East and 7°24'671" North with altitude of 1155.8 m.

### I.3 VEGETAL MATERIALS

Biological material is constituted of 20 morphotypes of Bambara groundnuts (*Vigna subterranea*) obtained from laboratory team of Biodiversity of University of Ngaoundere (Figure 1).

Formatted: Font: Italic



**Figure 1:** Accessions of Bambara groundnut.

#### I.4 EXPERIMENTAL DESIGN

The experimental design for this study is a block completely randomized with 3 replications. Ridge distances are respectively 1m, with 0.5m of length and width. The total surface of ridge is 0.5 m<sup>2</sup>. A total of 60 ridges are materialized ~~on a surface of~~with total experimental plot size of 61.95 m<sup>2</sup> (4.2 m x 14.75 m). Distance among ridge is of 60 cm and 25 cm. Seeds are sowing in the soil at 3 cm of depth with a space of 20 cm between poquets. For each 20 morphotypes, twelve seeds were sowed ~~by~~on each ridge. Each morphotype is represented by 36 plants. Sowing process was done during 10 July 2015 at Bini-Dang. Data collection is made according to the list of describer of Bambara groundnut (IPGRI/IITA/BAMNET, 2000).

#### I.5. EVALUATION OF AGRONOMIC PARAMÈTERS

Eleven parameters were chosen within the descriptors of Bambara groundnut to characterize and evaluate different morphotypes (Table 1). All the measurements ~~is~~was done on 5 plants of each ~~morphotypes~~morphotype.

**Table 1:** Parameters used to characterize the morphotypes.

N°	Parameters
1	Germination rate (%)
2	Date of first flowering (DAS)
3	Date of 50 % flowering (DAS)
4	Height of plants (cm)
5	Number of leaves per plant
6	Number of stems per plant
7	Number of pods per plant
8	Number of seeds per pod
9	Weight of 100 seeds (g)
10	Yields of seeds per plant (g)
11	Cycle period (DAS)

#### I.6. STATISTICAL ANALYSIS

**Formatted:** Superscript

**Comment [QA3]:** This statement is bit confusing for reader as their are two distances give 60 and 25 cm. What these two distances represents please explain.

**Comment [QA4]:** ???

**Comment [QA5]:** This work is very old

**Comment [QA6]:** When was data collected?? Please write the age of experiment.

**Comment [QA7]:** Please add brief descriptions how these paraeters were measured

For the different characters studied, we had proceeded with a comparison of means within morphotypes by ANOVA test with software STATGRAPHICS PLUS Version 5.0. For a highest significant difference between morphotypes for one character given, ANOVA is completed by the test of small significant difference (ppds). Which permit to identify the different morphotypes that ~~significant~~ significantly differed ~~to from~~ others (Dagnelie, 1998). Different performances of morphotypes for each quantitative character were evaluated on determining the minimum, maximum, average, standard-error and variation index. Evaluation of morphological diversity structuration was done by a Principal Component Analysis (ACP), a rising of hierarchical Classification (CAH) in order to appreciate the degree of similarity between analyzed unities and understand the phylogenetic relationship existing between them with the software XLSTAT Version 2007.8.04. Correlation coefficient of Bravais Pearson was used to test and appreciate the degree of association between the different characters studied.

## II. RESULTS AND DISCUSSION

### II.1 VARIABILITY ANALYSIS OF AGRONOMIC PARAMETERS

~~Exam's Results in of the~~ table 2a and 2b present the variability of different morphotypes for the eleven agronomic parameters. ANOVA ~~reveal~~ revealed a significant difference ( $p < 0.05$ ) between variety used  $\rightarrow$  germination rate, date of first flowering, date of 50 % of flowering, height of plants, number of stems per plant, number of pods per plant, number of seeds per pods, weight of 100 seeds, yields of seeds per plant and maturation time. Contrary, ~~not a no~~ significant ~~difference~~ difference for the for number of leaves per plant were obtained.

**Table 2a:** Genetic variability of 20 morphotypes for the 11 agronomic parameters.

Variety	TG (%)	DFI (JAS)	DF50%I (JAS)	HaP (cm)	NFe	NTg
CM/EN/AE/O2	63,88±33,67 <sup>defgh</sup>	50,33±0,57 <sup>cdefg</sup>	65,33±2,51 <sup>bcd</sup>	22±1,8 <sup>ab</sup>	95±8,52	6,33±0,76 <sup>bcd</sup>
CM/EN/DW/03	27,77±4,8 <sup>ab</sup>	46±2,00 <sup>abcd</sup>	62,33±1,52 <sup>bcd</sup>	22,68±3,48 <sup>ab</sup>	141,33±67,66	7,33±1,44 <sup>ef</sup>
CM/AD/MC/04	19,44±4,81 <sup>a</sup>	52±6,08 <sup>fg</sup>	62,33±2,08 <sup>bcd</sup>	22,56±1,25 <sup>ab</sup>	138,5±57,82	6,16±0,76 <sup>abcde</sup>
CM/AD/MC/05	33,33±14,43 <sup>abc</sup>	45,66±3,78 <sup>abc</sup>	62,33±1,52 <sup>bcd</sup>	24,25±4,02 <sup>ab</sup>	197,5±21,54	6,33±0,76 <sup>bcd</sup>
CM/EN/DW/07	69,44±39,38 <sup>efgh</sup>	45,33±2,51 <sup>ab</sup>	62,33±1,15 <sup>bcd</sup>	23,1±0,4 <sup>ab</sup>	115,5±10,81	4,83±0,28 <sup>ab</sup>
CM/EN/DW/09	47,22±4,8 <sup>abcdef</sup>	53±1,00 <sup>g</sup>	65,33±2,08 <sup>bcd</sup>	21,78±2,33 <sup>ab</sup>	118±43,53	4,66±0,76 <sup>a</sup>
CM/EN/AE/10	75±8,33 <sup>fgh</sup>	43±0,00 <sup>a</sup>	61±0,00 <sup>bc</sup>	24,58±3,64 <sup>abc</sup>	151,5±13,74	6±1,32 <sup>abcde</sup>
CM/EN/AE/13	66,66±22,04 <sup>efgh</sup>	48,33±3,51 <sup>bcd</sup>	64,66±3,51 <sup>bcd</sup>	21,58±2,96 <sup>ab</sup>	162,5±23,82	5,6±0,28 <sup>abcd</sup>
CM/EN/DW/14	75±8,33 <sup>fgh</sup>	49,33±2,30 <sup>bcd</sup>	66,66±5,77 <sup>de</sup>	23,33±2,02 <sup>ab</sup>	158,5±83,37	6±0,86 <sup>abcde</sup>
CM/AD/AE/15	77,77±12,73 <sup>gh</sup>	50,66±1,15 <sup>defg</sup>	54,33±8,08 <sup>a</sup>	30,13±2,87 <sup>d</sup>	191±45,92	6,66±1,75 <sup>cdef</sup>
CM/AD/MC/16	66,66±36,32 <sup>efgh</sup>	52,66±1,52 <sup>g</sup>	62,33±1,52 <sup>bcd</sup>	32,91±6,87 <sup>d</sup>	138,5±36,13	5,16±1,04 <sup>abc</sup>
CM/AD/MC/17	72,22±9,61 <sup>fgh</sup>	51,66±1,52 <sup>efg</sup>	62,66±1,52 <sup>bcd</sup>	33,66±1,75 <sup>d</sup>	143,5±42,64	5,66±1,15 <sup>abcd</sup>
CM/AD/MC/18	75±8,33 <sup>fgh</sup>	48±0,00 <sup>bcd</sup>	60±0,00 <sup>b</sup>	29,16±1,66 <sup>cd</sup>	171,5±64,73	6,66±0,76 <sup>cdef</sup>
CM/EN/DW/19	41,66±16,66 <sup>abcde</sup>	51,66±3,51 <sup>efg</sup>	68,66±1,52 <sup>e</sup>	20±2,94 <sup>a</sup>	150±68,73	5,33±0,76 <sup>abc</sup>
CM/EN/DW/20	55,55±4,8 <sup>bcd</sup>	47,33±4,04 <sup>abcde</sup>	64,33±5,13 <sup>bcd</sup>	23,08±2,09 <sup>ab</sup>	186,5±36,81	7±0,86 <sup>def</sup>
CM/EN/DW/21	80,55±12,72 <sup>gh</sup>	50,33±2,08 <sup>cdefg</sup>	66,33±2,51 <sup>cde</sup>	22,01±1,76 <sup>ab</sup>	161±50,58	6±1,00 <sup>abcde</sup>
CM/EN/DW/23	58,33±8,33 <sup>cdefg</sup>	50,66±5,13 <sup>defg</sup>	62,66±4,61 <sup>bcd</sup>	21,58±2,4 <sup>ab</sup>	132±37,58	8±0,86 <sup>f</sup>
CM/EN/DW/27	36,11±4,81 <sup>abcd</sup>	53±2,264 <sup>g</sup>	62,66±2,51 <sup>bcd</sup>	22,93±2,72 <sup>ab</sup>	202±42,46	7±0,86 <sup>def</sup>
CM/EN/DW/28	61,11±9,62 <sup>cdefgh</sup>	45,66±3,05 <sup>abc</sup>	62,66±3,51 <sup>bcd</sup>	25,05±2,22 <sup>bc</sup>	140±29,63	6,66±1,6 <sup>cdef</sup>
CM/EN/DW/29	88,89±4,81 <sup>h</sup>	48,66±3,51 <sup>bcd</sup>	62,66±2,51 <sup>bcd</sup>	23,48±1,34 <sup>ab</sup>	143±19,34	5,66±0,76 <sup>abcd</sup>
<b>Moyenne</b>	<b>59,58±23,66</b>	<b>49,16±3,76</b>	<b>63,08±3,79</b>	<b>24,49±4,45</b>	<b>151,86±41,18</b>	<b>6,15±1,16</b>
<b>PPDS (5%)</b>	<b>30,55</b>	<b>5</b>	<b>6</b>	<b>5,03</b>		<b>1,66</b>

**NB:** On the same lines, means following by the same letter are not significantly different at the level of probability considered ( $P \leq 0.05$ ).

**TG**=germination rate; **DFI**=date of first flowering; **DF50%I**=date of 50% flowering; **HaP**=height of plant; **NFe**=number of leaves; **NTg**=number of stems.

**Table 2b:** Genetic variability of 20 accessions for the 11 agronomic parameters.

Accessions	NGs	NGr	P100gr (g)	RGr (g)	TEm (JAS)
CM/EN/AE/O2	15,83±5,00 <sup>abc</sup>	1,19±0,17 <sup>a</sup>	78,46±2,07 <sup>gh</sup>	15,7±4,85 <sup>abc</sup>	123,33±2,88 <sup>cde</sup>
CM/EN/DW/03	25,16±4,85 <sup>ef</sup>	1,22±0,38 <sup>a</sup>	68,53±5,18 <sup>def</sup>	22,33±3,01 <sup>bcde</sup>	125±0,00 <sup>def</sup>
CM/AD/MC/04	14,66±8,60 <sup>a</sup>	1,11±0,19 <sup>a</sup>	105,7±0,00 <sup>i</sup>	18,06±9,12 <sup>abcd</sup>	125±0,00 <sup>def</sup>
CM/AD/MC/05	21±4 <sup>abcde</sup>	1,08±0,14 <sup>a</sup>	76,23±7,67 <sup>gh</sup>	20,46±4,81 <sup>bcde</sup>	123,33±2,88 <sup>cde</sup>
CM/EN/DW/07	14,33±1,52 <sup>a</sup>	1,80±0,17 <sup>bcde</sup>	93,46±7,82 <sup>i</sup>	20,13±3,03 <sup>bcde</sup>	126,66±2,88 <sup>ef</sup>
CM/EN/DW/09	19±7,21 <sup>abcde</sup>	1,44±0,38 <sup>abc</sup>	55±12,12 <sup>ab</sup>	14,46±3,99 <sup>ab</sup>	125±0,00 <sup>def</sup>
CM/EN/AE/10	17±6,08 <sup>abcd</sup>	1,00±0,00 <sup>a</sup>	60,6±6,00 <sup>abcd</sup>	10,76±4,29 <sup>a</sup>	118,33±5,77 <sup>ab</sup>
CM/EN/AE/13	23,66±1,15 <sup>cdef</sup>	1,5±0,17 <sup>abc</sup>	55,36±3,70 <sup>ab</sup>	19,6±1,25 <sup>bcde</sup>	128,33±2,88 <sup>f</sup>
CM/EN/DW/14	19±3,12 <sup>abcde</sup>	1,5±0,17 <sup>abc</sup>	51,46±4,04 <sup>a</sup>	18,1±2,26 <sup>abcd</sup>	123,33±2,88 <sup>cde</sup>
CM/AD/AE/15	15,5±4,92 <sup>ab</sup>	1,89±0,19 <sup>cde</sup>	54,13±3,53 <sup>ab</sup>	18,93±7,44 <sup>bcde</sup>	121,66±2,88 <sup>bcd</sup>
CM/AD/MC/16	14,83±2,02 <sup>a</sup>	2,33±0,33 <sup>e</sup>	60±4,47 <sup>abcd</sup>	21,16±1,94 <sup>bcde</sup>	125±0,00 <sup>def</sup>
CM/AD/MC/17	16,66±5,75 <sup>abcd</sup>	2,22±0,69 <sup>de</sup>	52,76±5,10 <sup>a</sup>	16,13±9,13 <sup>abc</sup>	123,33±2,88 <sup>cde</sup>
CM/AD/MC/18	16,33±2,92 <sup>abcd</sup>	1,77±0,38 <sup>bcd</sup>	62,66±2,05 <sup>bcde</sup>	16,4±6,2 <sup>abc</sup>	120±0,00 <sup>bc</sup>
CM/EN/DW/19	24,16±6,00 <sup>def</sup>	1,77±0,38 <sup>bcd</sup>	54,76±2,21 <sup>ab</sup>	22,76±5,44 <sup>cde</sup>	126,66±2,88 <sup>ef</sup>
CM/EN/DW/20	15,66±1,52 <sup>ab</sup>	1,5±0,17 <sup>abc</sup>	67,3±7,06 <sup>cdef</sup>	15,4±0,95 <sup>abc</sup>	120±5,00 <sup>bc</sup>
CM/EN/DW/21	13,83±2,92 <sup>a</sup>	1,30±0,33 <sup>ab</sup>	66,5±9,10 <sup>cde</sup>	14,53±3,8 <sup>ab</sup>	120±5,00 <sup>bc</sup>
CM/EN/DW/23	14±4,27 <sup>a</sup>	1,41±0,52 <sup>abc</sup>	83,6±2,00 <sup>h</sup>	16,43±4,62 <sup>abc</sup>	125±0,00 <sup>def</sup>
CM/EN/DW/27	30,83±5,00 <sup>f</sup>	1,5±0,17 <sup>abc</sup>	61,06±2,55 <sup>abcd</sup>	26,76±3,06 <sup>e</sup>	125±0,00 <sup>def</sup>
CM/EN/DW/28	23,33±5,00 <sup>bcdef</sup>	1,4±0,41 <sup>abc</sup>	58,73±9,20 <sup>abc</sup>	23,26±1,95 <sup>cde</sup>	115±0,00 <sup>a</sup>
CM/EN/DW/29	19,83±5,34 <sup>abcde</sup>	1,41±0,22 <sup>abc</sup>	70,86±4,75 <sup>efg</sup>	25,36±4,21 <sup>de</sup>	120±5,00 <sup>bc</sup>
<b>Moyenne</b>	<b>18,73±6,01</b>	<b>1,52±0,43</b>	<b>66.86±14,98</b>	<b>18,84±5,61</b>	<b>123±4,03</b>
<b>PPDS (5%)</b>	<b>8</b>	<b>0,55</b>	<b>9,8</b>	<b>8,23</b>	<b>5</b>

**NB:** On the same lines, means following by the same letter are not significantly different at the level of probability considered ( $P \leq 0.05$ ).

**NGs**=Number of pods per plant; **NGr**=Number of seeds per plant; **P100gr**=Weight of 100 seeds; **RGr**=Yields of seeds per plant; **TEm**=Maturation time.



## II.1 VARIABILITY OF QUANTITATIVE PARAMETERS OF AGRONOMIC DIVERSITY

Results of **Table 3** indicate important standard between the minimum and maximum. Variation index are between 2.66 % (time of maturation) from 32.38 % (germination rate). High values (CV>20 %) was observed for 5 of 11 characters analyzed. All of results traduce the existence of important diversity within the species. This important phenotypic variability observed could result from the expression of high genotypic heterogeneity but also of the influence of some environmental factors. Theses different accessions have a germination rate that varying from 19.44 to 88.89 %

These results corroborate to works of results of Touré *et al.* (2013) in Côte d'Ivoire, were reported that germination rate on Bambara groundnut varying from 26.8 to 91.25 %. Contrary, theses results did not corroborate to works of Djé *et al.* (2005) were founded that germination rate varying from 64 to 88 % on fourth morphotypes. Variability response expressed within ours conditions may traduce heterogeneity of germinative quality of seeds that arising principally from state purity of seeds and their physiology.

Contrary to this observation, Massawe *et al.* (2002) showed a favor effect of height of seeds on appearance of plants. Theses different accessions of Bambara groundnut have a flowering at 50 % that varying from 54 to 89 days. Theses results are not similar to works results obtained by Brink *et al.* (2006) and Amadou *et al.* (2015), who reported respectively a period of 32 to 42 days and 33 to 40 days respectively. Theses different results could be explain by the difference in number of variety studied in this testour study and geographical provenance are unknown. Dimakatso (2006) reported that flowering on Bambara groundnut is undetermined. The date of flowering could be recorded like good trait agronomic in the program of amelioration of Bambara groundnut in Cameroon. Accessions of Bambara groundnut tested have a height around 20 and 33.66 cm with 5 to 8 stems.

Theses observations corroborate to the works obtained by Ndiang *et al.* (2012) in Cameroon, showed that the height of plants varies from 20 to 38 cm with 5 to 8 stems. Contrary, these values are different of those obtained by Sobda *et al.* (2013) in Cameroon that reported the number of stems per plant varying between 16 and 24. Number of pods per plants in this study varying respectively between 14 and 31. Theses results is-are different to the works-observations found-made by Ouédraogo *et al.* (2008) in Burkina Faso on 310 accessions and prove that the number of pods on Bambara groundnut varying from 4 to 50. The difference in observationsed within ours-our conditions-experiment could be explain by the difference between morphotypes, state of fertility of experimental plots and local climatic conditions. Weight of 100 seeds varying from 51.46 to 105.7g. Theses s results corroborate to with the works obtained-byof Bonny et Djé (2011) in Côte d'Ivoire, who reported that weight of 100 seeds on Bambara groundnut varyvarieding between 35.83 and 111.3g. Results are in contradiction of works of Ndiang *et al.* (2012) in Cameroun and Touré *et al.* (2012) in Côte d'Ivoire that founded respectively-total weight of 100 seeds were in range of 52 to 82g and 40.71 to 76.54g respectively.

Theses differences obtained could be explained by the cultivation of accessions on a humid border on different climatic and ecologic conditions of the production area (on a high humidity of soil due to rain). Yield of seeds per plant in this study fluctuate from 10.76 to 26.76g. ThesesThese results corroborate of-with resultworks obtained by Shegro *et al.*

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic



(2013) in South Africa, who reported that the yield of seeds per plant on Bambara groundnut ~~varying-varied~~ between 10.20g and 57g. Amplitude time for maturation varies from 115 to 128 days. Results correspond to works founded by Goli *et al.* (1997) in Burkina Faso that reported the duration of lifecycle of Bambara groundnut ~~varying varied~~ between 100 and 160 DAS. However, results differed ~~of-from~~ those obtained by par Ndiang *et al.* (2014) in Cameroon. These ~~authors-scientists~~ reported that the duration of lifecycle on Bambara groundnut is 90 days. ~~InfaetIn fact~~, numerous previous works showed that the length of day (Harris & Azam-ali, 1993; Linnemann *et al.*, 1995; Brink, 1999), temperature (Linnemann & Craufurd, 1994; Brink, 1998) and humidity (Collinson *et al.*, 1996) cause variable effects on vegetative and physiological development of Bambara groundnut.

Formatted: Font: Italic

Formatted: Font: Italic

**Table 3:** Performances of 11 quantitative characters measured for 20 accessions.

Variables	Observations	Minima	Maxima	Means	Standard error	CV (%)
TG (%)	20	19.44	88.89	59.58	19.30	32.38
DF1 (JAS)	20	43.00	53.00	49.16	2.92	5.94
DF50%1 (JAS)	20	54.33	68.66	63.08	2.93	4.64
Hap (cm)	20	20.00	33.66	24.49	3.85	15.72
Nfe	20	95.00	202.00	151.87	27.92	18.34
NTg	20	4.66	8.00	6.15	0.84	13.66
NGs	20	13.83	30.83	18.73	4.63	24.72
NGr	20	1.00	2.33	1.52	0.35	23.03
P100gr (g)	20	51.46	105.70	66.86	14.43	21.58
RGr (g)	20	10.76	26.76	18.84	4.02	21.33
TEm (JAS)	20	115.00	128.33	122.99	3.27	2.66

TG=germination rate; DF1=date of first flowering; DF50%1=date of 50% of flowering; HaP=height of plants; Nfe=number of leaves per plant; NTg=number of stems per plant; NGs=number of pods per plant; NGr=number of seeds per pods; P100gr=weight of 100 seeds; RGr= yield of seeds per plant; TEm=maturation time.

## II.2. STRUCTURATION OF AGRO-MORPHOLOGIC DIVERSITY BY PRINCIPAL COMPONENT ANALYSIS (ACP)

**Table 4** ~~show-shows~~ proper values and contribution of characters on the axis of analysis of principal component (ACP). ACP ~~indicate-indicates~~ that the fourth primary component ~~explain-explains respectively~~ 23.53 %; 19.79%; 19.13 % and 12.15 % of the variability, with around 75.6 % of the total variability. The two primary components which recorded 44.31% of the variance are retained for analyzing the agronomic variability of accessions. **Axis 1** that explains 24.53 % of the variability is highest correlated with the height of plants, germination rate, and yield of seeds per plant. **Axis 2** with 19.79 % of variance defines the phenology and yield parameters. The date of first flowering, the date of 50 % flowering, maturation time, number of pods per plant, number of seeds per ~~pods-pod~~ and yield of seeds per plant are positively correlated in this axis. ~~Theses-These~~ characters that are positively correlated on two axes are factors most discriminant suggesting their consideration in the selection for the important agronomic traits.

**Table 4:** Proper ~~v~~Values and contribution of characters on the axis of component of principal analysis.

	Axe 1	Axe 2	Axe 3	Axe 4
Valeur propre	2,698	2,176	2,104	1,337
Variabilité (%)	24,528	19,786	19,129	12,150
% cumulé	24,528	44,314	63,443	75,593

#### caractères définissant les axes et leurs valeurs

##### propres

TG	0,740	-0,169	-0,114	-0,405
DFI	0,029	0,630	-0,326	0,283
DF50%l	-0,505	0,203	-0,477	-0,541
Hap	0,854	0,212	0,151	0,279
Nfe	0,070	0,154	0,778	0,022
NTg	-0,194	-0,345	0,622	0,314
NGs	-0,520	0,516	0,542	-0,295
NGr	0,667	0,632	-0,133	0,154
P100gr	-0,408	-0,458	-0,300	0,621
RGr	-0,306	0,594	0,406	0,108
Tem	-0,385	0,542	-0,421	0,339

TG=taux de germination ; DFI=date de 1<sup>ère</sup> floraison ; DF50%l=date de 50% de floraison ; HaP=hauteur des plantes ; Nfe=nombre des feuilles par plante ; NTg=nombre des tiges par plante ; NGs=nombre de gousses par plante ; NGr=nombre des graines par gousse ; P100gr=poids de 100 graines ; RGr=rendement en graine par plante ; TEm=temps de maturation.


### II.3. ANALYSE DE LA DIVERSITÉ DES ACCESSIONS PAR LA CLASSIFICATION ASCENDANTE HIÉRARCHIQUE (CAH)

Comment [QA8]: English please

Dendrogram rising to hierarchical classification (**Figure 2**) reveal a repartition of 20 accessions in 5 groups. The **groups 1** is constituted of 3 accessions (CM/EN/AE/02, CM/EN/DW/07, CM/EN/DW/23). The **groups 2** that regroup 3 accessions (CM/EN/DW/03, CM/EN/DW/09, CM/EN/DW/19), characterized by a cycle of maturation of 126 days. Variety (CM/AD/MC/04) constitute the **group 3** with, with a weight of 100 seeds most high (105.7g). This variety could be selected for a food security. The **group 4** includes 3 accessions (CM/AD//04, CM/EN/DW/20, CM/EN/DW/27). This group is characterized by a yield of seeds per plant most high (20.7g). The **group 5** is composed of 10 accessions (CM/EN/AE/10, CM/EN/AE/13, CM/EN/DW/14, CM/AD/AE/15, CM/AD/MC/16, CM/AD/MC/17, CM/AD/MC/18, CM/EN/DW/21, CM/EN/DW/28, CM/EN/DW/29).

It is characterized by a germination rate most high (73.88 %), a flowering between 48 and 62 days, with a height of 26.56 cm. ~~these~~this group is constituted of morphotype with earlier cycle. Precocity~~The precocity~~ of cycle of these accessions made this culture interesting for rural farmers during the ~~weld~~ period, because they achieve their maturity before others variety and constitute one important source of income for producers and reseller.

Morphological and phenological difference observed between different groups of phenotypic diversity suggested that accessions are maintained under **evolutive** processes which are different to their agro-ecosystem respectively. Agro-ecosystems are susceptible to exert variable selective pressure on genotypes (Doku & Karikari, 1971; Sadiki & Jarvis, 2005).



**Figure 2:** Dendrogram rising to hierarchical classification of accessions of Bambara groundnut.

Average performance of characters of different groups formed rising to hierarchical classification are represented on **Table 5**.

**Table 5:** Means performances of characters of different groups obtained by CAH.

Groupes	TG	DFI	DF50%I	Hap	Nfe	NTg	NGs	NGr	P100gr	RGr	TEm
1	63.87	48.77	63.44	22.22	114.16	6.38	14.72	1.46	85.17	17.42	124.99
2	38.88	50.22	65.44	21.48	136.44	5.77	22.77	1.47	59.43	19.85	125.5
3	19.44	52.00	62.33	22.56	138.50	6.16	14.66	1.11	105.70	18.06	125.00
4	41.66	48.66	63.10	23.42	195.33	6.77	22.4	1.36	68.19	20.87	122.77
5	73.88	48.82	62.32	26.58	156.10	6.01	17.99	1.63	59.30	18.42	121.49

**TG**=germination rate; **DFI**=date of first flowering; **DF50%I**=date of 50% of flowering; **Hap**=height of plants; **Nfe**=number of leaves per plant; **NTg**=number of stems per plant;

**NGs**=number of pods per plant; **NGr**=number of seeds per pods; **P100gr**=weight of 100 seeds; **RGr**= yield of seeds per plant; **TEm**=maturation time.

#### II.4 CORRELATIONS BETWEEN PARAMETERS STUDIED

**Table 6** shows the correlation concerning 11 characters that represent the variability. High positive correlation  $r=0.72$  is observed between the height of plant and number of seeds per plant. A correlation  $r=0.67$  bind the number of pods per plant and the yields of seeds per plant. The high negative correlation ( $r=-0.58$ ) is considerate between the height of plants and the date of 50 % flowering. The positive correlation observed between characters could facilitate the genetic amelioration of all others characters correlated positively. Contrary, negative correlation observed between the characters studied could limited the selection, possibility to obtain cultivar with earlier cycle and a high yield producing heavy seeds that producers and consumers needed. Correlations constitute an important tool for amelioration of characters on integrating on a program of selection.

**Table 6:** Correlation Matrix of Pearson of 11 agronomic parameters.

Variables	TG	DFI	DF50%	Hap	Nfe	NTg	NGs	NGr	P100g	RGr	TEm
<b>TG</b>	<b>1</b>										
<b>DFI</b>	-0,149	<b>1</b>									
<b>DF50%</b>	-0,162	0,189	<b>1</b>								
<b>Hap</b>	0,361	0,126	<b>-0,576*</b>	<b>1</b>							
<b>Nfe</b>	-0,085	-0,069	-0,311	0,157	<b>1</b>						
<b>NTg</b>	-0,221	-0,155	-0,303	-0,095	0,355	<b>1</b>					
<b>NGs</b>	-0,426	-0,013	0,197	-0,293	0,338	0,130	<b>1</b>				
<b>NGr</b>	0,344	0,406	-0,173	<b>0,712**</b>	0,011	-0,328	-0,151	<b>1</b>			
<b>P100g</b>	-0,361	-0,115	-0,067	-0,318	-0,321	0,142	-0,370	-0,391	<b>1</b>		
<b>RGr</b>	-0,261	0,126	-0,013	-0,058	0,211	0,051	<b>0,674*</b>	0,187	-0,017	<b>1</b>	
<b>TEm</b>	-0,406	0,425	0,263	-0,218	-0,166	-0,266	0,156	0,191	0,185	0,188	<b>1</b>

**NB:**\*significatif (5%); \*\*très significatif (1%); **TG**=germination rate; **DFI**=date of first flowering; **DF50%**=date of 50% of flowering; **Hap**=height of plants; **Nfe**=number of leaves per plant; **NTg**=number of stems per plant; **NGs**=number of pods per plant; **NGr**=number of seeds per pods; **P100g**=weight of 100 seeds; **RGr**= yield of seeds per plant; **TEm**=maturation time.

## CONCLUSION

~~Objective~~ The objective of ~~these~~ this study were to identify agronomic traits responsible ~~of~~ for the variation between 20 accessions of Bambara groundnut. In all of 11 characters evaluated in this study, only one of them that not ~~permit~~ permitted to differentiate the accessions. Important variability between ~~theses~~ these characters was observed. ~~Rising~~ The rising of hierarchical classification showed one of important agronomic diversity and regroup the accessions in 5 groups. Agro-morphological diversity observed in this study permit to explain that ~~theses~~ these accessions constitute a good reservoir for the genetical characteristic of Bambara groundnut. Correlation observed between morphologic and agronomic parameters made the characters concerned as important index to use in the program of amelioration. The characters analyzed could constitute of base criteria to differentiate variety to others regions and could be used in the program of varietal amelioration of Bambara groundnut in Cameroon.

## REFERENCE:

- [1] Aké Assi L. (1988). La diversité génétique des espèces végétales sous-exploitées d'Afrique. In P. Perrino (Ed.) Crop genetic resources of Africa, vol. 1. Nairobi, Kenya : IBPGR/IITA/UNEP, p. 53-88.
- [2] Altieri M. A. (1986). L'agro-écologie: bases scientifiques d'une agriculture alternative. Paris: Debard, 237 p.
- [3] Amadou H. I., Doumma A., Katsilero A., Zoubeirou A. M., Nourou S. S. M. (2015). Agro-morphological variability in fourteen bambara groundnut (*Vigna subterranea* (L.) Verdc.) morphotypes cultivated in Niger. *Scholars Academic Journal of Biosciences*, 3(9):774-781.
- [4] Amarteifio J. O., Tibe O., Njogu R. M. (2006). The mineral composition of Bambara groundnut (*Vigna subterranea* (L.) Verdc.) Grown in Southern Africa. *African Journal of Biotechnology* 5(23): 2408-2411.
- [5] Azam-ali S. N., Sesay A., Karikari S. K., Massawe F. J., Aguilar-manjarrez J., Bannayan M., Hampson K. J. (2001). Assessing the potential of an underutilized crop- A case study using Bambara groundnut. *Experimental Agriculture*, 37: 433-472.
- [6] Basu S., Roberts J., Mithen R., Azam-Ali S., and Pasquet R. (2004). The genetics of Bambara groundnut and the construction of a genetic linkage map. In New directions for a diverse planet: Proceedings of the 4th International Crop Science Congress Brisbane, Australia. 7p.
- [7] Bonny B. S et Djè Yao. (2011). Variabilité morphologique et agronomique des variétés traditionnelles de voandzou (*Vigna subterranea* (L.) Verdc. (Fabaceae) de Cote d'Ivoire. *Journal of Applied Bioscience*, 41: 2820-2835.
- [8] Boutrais J. (1978). Deux études sur l'élevage en zone tropicale humide (Cameroun). ORSTOM, Paris, France, 194p.
- [9] Brink M. (1998). Matching crops and environments: quantifying photothermal influences on reproductive development in Bambara groundnut (*Vigna subterranea* (L.) Verdc). PhD Thesis, Wageningen Agricultural University, pp.161.
- [10] Brink M. (1999). Development, growth and dry matter partitioning in Bambara groundnut (*Vigna subterranea*) as influenced by photoperiod and shading. *Journal of Agricultural Science*, 133: 159-166.
- [11] Brink M., Ramolemana G. M., Sibuga K. P. (2006). *Vigna subterranea* (L.) Verdc. In: Brink M., Belay G. (Eds). Plant Resources of Tropical African. Cereals and pulses. PROTA Foundation, Wageningen, Netherlands. pp. 213-218.
- [12] Brink M. and Belay G. (2006). Plant Resources of Tropical Africa 1. Cereals and pulses. PROTA Foundation, Wageningen, Netherlands/Backhuys Publishers, Leiden, Netherlands/CTA, Wageningen, Netherlands. Pp. 213-218.
- [13] Brough S. H., Azam-Ali S. N. (1992). The effect of soil moisture on the proximate composition of Bambara groundnut (*Vigna subterranea* (L.) Verdc.). *Journal of Science Food and Agriculture*, 60: 197-203.
- [14] Brough S. H., Azam-Ali S. N. & Taylor A. J. (1993). The potential of Bambara groundnut (*Vigna subterranea*) in vegetable milk production and basic protein functionality systems. *Food Chemistry* 47, 277-283.

Comment [QA9]: Missing or incorrect citation

- [15] Collinson S. T., Azam-Ali S. N., Chavula K. M., Hodson D. A. (1996). Growth, development and yield of bambara groundnut (*Vigna subterranea*) in response to soil moisture. *Journal of Agricultural Science*, 126: 307-318.
- [16] Collinson S. T., Sibuga K. P., Tarimo A. J. P., Azam-Ali S. N. (2000). Influence of sowing date on the growth and yield of Bambara groundnut landraces in Tanzania. *Exp. Agric.* 36, p. 1-13.
- [17] Dagnelie P. (1998). Statistique théorique et appliquée (Tome 1). Bruxelles : De Boeck & Larcier.
- [18] Dimakatso R. M. (2006). Evaluation of bambara groundnut (*Vigna subterranea*) for yield stability and yield related characteristics. Agricultural Research Council - Grain Crop Institute (ARC-GCI).
- [19] Djè Y., Bonny B. S., Zoro Bi I. A. (2005). Observations préliminaires de la variabilité entre quelques morphotypes de voandzou (*Vigna subterranea* (L.) Verdc., Fabaceae) de Côte d'Ivoire. *Biotechnologie, Agronomie, Société et Environnement* 9 (4): 249-258pp.
- [20] Doku E. V., Karikari S. (1971). The roles of plants in pollination and pod production of Bambara groundnut. *Econ. Bot.* 25, p. 357-362.
- [21] Edjé O. T., Sesay A. (2004). Effects of seed source on performance and yield of Bambara groundnut (*Vigna subterranea* (L.) Verdc.) landraces. In: Proceeding of the International Bambara Groundnut Symposium" European Union Framework Programme 5 Botswana College of Agriculture, Botswana 8 -12 September 2003. PP. 141-152.
- [22] Goli A. E., Begemann F., Ng N. Q. (1997). Characterization and evaluation of IITA's Bambara groundnut collection. In Heller J., Begemann E., Mushonga J. (Eds). Promotion of the conservation and use of underutilized and neglected crops. 9. Proceedings of the workshop on conservation and improvement of bambara groundnut (*Vigna subterranea* (L.) Verdc.), 14-16 November 1995, IPGRI, Harare, Zimbabwe, p. 101-118.
- [23] Goli A. E. (1997). Bibliographical review of Bambara groundnut. Cotonou: IPGRI.
- [24] Harris D., Azam-Ali S. N. (1993). Implications of day length sensitivity in Bambara groundnut (*Vigna subterranea*) for production in Botswana. *Agricultural Science*, 120: 75-78.
- [25] Howell J. A., Eshbaugh W. H., Guttman S. & Rabakonandrianina E. (1994). Common names given to bambara groundnut (*Vigna subterranea*: Fabaceae) in central Madagascar. *Economic Botany*, 48, 217-221.
- [26] IPGRI, IITA, BAMNET. (2000). Descripteur du pois bambara (*Vigna subterranea*). Institut international des ressources phytogénétiques, Rome, Italie; Institut International d'Agriculture Tropicale, Ibadan, Nigeria; Réseau International de pois bambara, Allemagne. 48.
- [27] Karikari S. K. (1971). Economic importance of bambara groundnut. *World Crops* 23(4):195-196.
- [28] Kumaga F. K., Adiku S. G. K., and Ofori K. (2003). Effect of Post-flowering Water Stress on Dry Matter and Yield of Three Tropical Grain Legumes. *International journal of agriculture & biology*, 1560-8530/2003/05-4-405-407.
- [29] Linnemann A. R., Craufurd P. Q. (1994). Effects of temperature and photoperiod on phenological development in three genotypes of Bambara groundnut (*Vigna subterranea*). *Annals of Botany*, 74: 675-681.
- [30] Linnemann A. R., Westphal M., Wessel E. (1995). Photoperiod regulation of development and growth in bambara groundnut (*Vigna subterranea*). *Field Crops Research*, 40: 39-47.
- [31] Linnemann A. R. (1993). Phenological development in bambara groundnut (*Vigna subterranea*) at constant exposure to photoperiods of 10 to 16 h. *Ann. Bot.* 71, p. 5-452.
- [32] Linnemann A. R. (1994). Phenological development in bambara groundnut (*Vigna subterranea*) at alternate exposure to 12 and 14 h photoperiods. *Journal of Agricultural Sciences*. 123, p. 40.
- [33] Makanda I., Tongona P., Madamba R., Icishahayo D and Derera J. (2009). Path Coefficient Analysis of Bambara Groundnut Pod Yield Components at Four Planting Dates. *Research Journal of Agriculture and Biological Sciences*, 5(3): 287.
- [34] Massawe F. J., Dickinson M., Roberts J. A., Azam-Ali S. N. (2002). Genetic diversity in Bambara groundnut (*Vigna subterranea* (L.) Verdc) landraces revealed by AFLP markers. *Genome*. 45: 1175-1180.
- [35] Massawe F. J., Mwale S. S., Azam-Ali S. N., Roberts J. A. (2005). Breeding in Bambara groundnut (*Vigna subterranea* (L.) Verdc.): strategic considerations. *African Journal of Biotechnology*, 4(6): 463-471.
- [36] MINADER. (2012). AGRI-STAT N°17: Annuaire des statistiques du secteur agricole Campagnes 2009 et 2010 123 p.
- [37] Mungate D. (1997). Bambara groundnut (*Vigna subterranea* (L.) Verdc.). In: Heller J., Begemann F., Mushonga J., eds. Promoting the conservation and use of underutilized and neglected crops. 9. Proceedings of the

Comment [QA10]: Missing citation

Comment [QA11]: Missing or incorrect citation in text

Comment [QA12]: missing

Comment [QA13]: Missing citation

Comment [QA14]: Missing or incorrect citation

Comment [QA15]: Missing citation



- workshop on Conservation and Improvement of Bambara Groundnut (*Vigna subterranea* (L.) Verdc.), 14-16 November 1995, Harare, Zimbabwe. 3-4.
- [38] Nacoulma-Ouédraogo O. G. (1996). Plantes médicinales et pratiques médicales traditionnelles au Burkina-Faso: cas du plateau central, Thèse de Doctorat ès Sciences naturelles, Université de Ouagadougou, Ouagadougou, Burkina-Faso.
- [39] Ndiang Z., Bell J. M., Missoup A. D., Fokam P. E., Amougou Akoa. (2012). Etude de la variabilité morphologique de quelques variétés de voandzou au Cameroun. *Journal of Applied Biosciences*, 60: 4394-4409.
- [40] Ndiang Z. Bell., J. M. Fokam, P. E. Ouattara, B. Simo, C. Dibong, D. S. (2014) Agro-morphological variability in twelve Bambara groundnut (*Vigna subterranea* (L.) Verdc) accessions in Cameroon. *Sciences, Technologies et Development*, 2014; 16:38-45.
- [41] Ntundu W. H., Bach I. C., Christiansen J. L., Andersen S. B. (2004). Analysis of genetic diversity in Bambara groundnut (*Vigna subterranea* (L.) Verdc.) Landraces using (AFLP) markers. *African Journal of Biotechnology*, 3 (4): 220-225.
- [42] Onwubiko N. I. C., Odum O. B., Utazi C. O And Poly-Mbah P. C. (2011). Studies on the adaptation of Bambara Groundnut (*Vigna Subterranea* (L.) Verdc) in Owerri Southeastern Nigeria. *New York Science Journal*, 4(2):60-67.
- [43] Ouedraogo M., Ouedraogo J. T., Tignere J. B., Balma D., Dabire C. B. and Konate G. (2008). Characterization and evaluation of accessions of Bambara groundnut (*Vigna subterranea* (L.) Verdc) from Burkina Faso, *Science and Nature*, 5: 191-197.
- [44] Pasquet R. S. and Fotso M. (1991). Les légumineuses alimentaires du Cameroun, premiers résultats. In Boutrais, J., ed, Du politique à l'économique, études historiques dans le bassin du lac Tchad, ORSTOM, Paris: 317-360.
- [45] Rippstein G. (1985). Etude de la végétation de l'Adamaoua, évolution, conservation, régénération et amélioration d'un écosystème pâturé au Cameroun. Maison Alfort, IEMVT 14, France, 367p.
- [46] Sadiki M., Jarvis D. (2005). Conservation in situ de la diversité génétique des cultures par sa gestion à la ferme dans les agroécosystèmes marocains. Les Actes du BRG, 5: 445-464.
- [47] Sobda Gonné, Wassouo F. A. Koubala Bargui B. (2013). Assessment of twenty bambara groundnut (*Vigna subterranea* (L.) Verdc) landraces using quantitative morphological traits. *International Journal of Plant Research*, 3(3): 39-45.
- [48] Touré Y., Koné M., Silué S., Kouadio Y. J. (2013). Prospection, collecte et caractérisation agromorphologique des morphotypes de voandzou (*Vigna subterranea* (L.) Verdc. (Fabaceae) de la zone savanicole en Côte d'Ivoire. *European Scientific Journal*, 9(24): 308-325.
- [49] Touré Y., Koné M., Kouakou T. H. & D. Koné. (2012). Agromorphological and Phenological Variability of 10 Bambara Groundnut (*Vigna subterranea* (L.) Verdc. (Fabaceae)) Landraces cultivated in the Ivory Coast, *Tropicultura*, 30(4): 216-221
- [50] Zerihun Tadele. (2009). Role of Orphan Crops in Enhancing and Diversifying Food Production in Africa. *African Technology Development Forum Journal*, Volume 6, Issue 3 and 4, 83 p.

**Comment [QA16]:** Missing citation from article text