

Study Protocol

Some Toxic Effects of Formalin on Ist (First) Year M.B.B.S. (Don't use this abbreviation) Students: A Cohort Study

Abstract: (Why not use 'Project Summary' instead of Abstract)

Background: Formalin can lead to allergy, toxicity and carcinomas. In anatomy department students gets exposed to formalin either by inhaling it, or by (contact with the) skin as fluids containing formaldehyde may be absorbed. Formalin interact with molecules on cell membrane producing changes in the nuclei, changes in proteins, vacuolation of cytoplasm, and changes in DNA.

Objective: To estimate toxic effects of formalin on Ist year medical students.

Methodology: The participants for this prospective cohort study will be Ist year M.B.B.S. (write in full) students at Jawaharlal Nehru Medical College, Sawangi (meghe), Wardha. Self-administered questionnaire will be given and complete blood count and thyroid function tests will be done prior to study. After completion of cadaveric dissection, repeat blood sample will be collected to estimate the complete blood count and thyroid function tests.

Expected Results: Exposure to formalin may lead to abnormal values of complete blood count(CBC) and thyroid function tests (TFT).

Conclusion: The study can focus the toxic effects of formalin on Ist year M.B.B.S. students ,which may necessitate to reconsider the formalin volume and ventilation issue in the dissecting halls. There should be regular assessment of working practices to control effects of formalin at the department of Anatomy.

Keywords: Formalin, complete blood count (CBC), thyroid function tests(TFT).

Introduction:

For centuries the fundamental teaching tool in anatomy has been human cadaveric dissection. Even if the technology is revolutionized gross anatomy teaching over certain years, research clearly indicated the uniqueness and value of dissection of cadaver in anatomical sciences. Cadaveric dissection can inculcate human values in medical students, as cadaver acts as first silent mentor to them. This will definitely contribute to make the future generation of doctors more empathetic (1).

August Wilheld Von Hofmann discovered Formaldehyde in 1867. At room temperature, it is inflammable and possess a highly irritating odour.(2) Formalin contains 40% by volume or 37% by weight of formaldehyde gas in water. Embalming of cadaver is done usually by using

formalin. There is widespread use of formalin in tissue fixation and embalming of cadaver but the major issue is safety.(3)

Formalin can lead to allergy, toxicity and carcinomas.(4,5) . In anatomy department students get(s) exposed to formalin either by inhaling it, or by skin as fluids containing formaldehyde may be absorbed. Exposure to formalin leads to irritation and also obstructive disorders of air passages, ocular irritations, clouding of cornea, various leukemias, nasopharyngeal carcinomas. (6)It also leads to various obstetrics and gynaecological problems as spontaneous abortions, congenital malformations, and irregular menstrual cycles. (7) Formalin may also lead to dermatitis.(8)

As exposed person gets habitual within a ??? some hours of exposure to formalin, toxic effects of formalin worsens. Accordingly, those person(s) remaining in environment of gradually increased level of formaldehyde concentrations, faces (???) increased exposure levels and may develop(s) chronic toxic effects.(9)

Various standard(s) are provided by concerned authorities to protect exposed workers to formaldehyde and include all occupational exposures to formalin like dissection hall staff. The exposure limit that can be permitted for an 8-hour shift in the working place is 0.75 ppm of air (reference needed). The second limit permits exposure of 2 ppm as the maximum exposure that can be allowed for 15-minutes (reference needed).

In medical colleges, medical students and faculty members working in Anatomy department are prone to develop the toxic effects of formalin. Studies have shown that because of formalin used for embalming of cadavers in the anatomy dissection halls, students get(s) exposed to formalin and effects can be dangerous if the ventilation of dissection hall is not properly maintained. Formalin can interact with molecules on cell membrane producing changes in the nuclei, vacuoles in cytoplasm, changes in proteins and DNA(10) . It was observed that some female students develop(s) menstrual cycle disorders after some months of attending cadaveric dissection(reference needed). Endocrinological effects of formalin are not much explored, especially in human beings that leads us to think about the possibility of toxic effects of formalin on thyroid gland(reference needed).

Aim: To estimate the effects of formalin on Ist year M.B.B.S. students attending cadaveric dissection at Anatomy department.

Objectives:

1. To estimate the immediate effects of formalin.
2. To estimate the effect of formalin on complete blood count.
3. To estimate the effect of formalin on thyroid gland.

Material and methods

Study design: Prospective Cohort Study

At Jawaharlal Nehru Medical College, Sawangi (Meghe), Wardha one dissection hall measuring 500 sq m. is situated at the ground floor of the college main building. There are 16 windows in the side walls for natural ventilation with 20 cadavers. Artificial ventilation comprised of 38 ceiling fans, 9 exhaust fans attached to the roof and 2 suction devices in dissection hall. During practical's, cut sections and dissected parts of the cadaver are sometimes kept in formalin solution either in trays or in basins for demonstration and teaching purposes.

The study participants for this prospective cohort study will be 30 (15 male and 15 female) first year medical students attending cadaveric dissection for 2 hours daily. Study will start with first day of dissection and will end with completion of dissection of cadavers i.e. after around one year. Participants will be selected randomly by coin flip method. Self-administered questionnaire will be given and complete blood count and thyroid function tests will be done prior to study. After completion of cadaveric dissection, blood sample again will be collected to observe the chronic toxic effect of formalin on complete blood count and thyroid function tests. Comparison group will comprise of 30 (15 male and 15 female) final year medical students.

Informed consent will be taken from the study participants.

Inclusion criteria

1st year medical students exposed to formalin during cadaveric dissection.

Exclusion criteria

- Diagnosed cases of asthma, hypothyroidism and polycystic ovarian disease.
- Those who have undergone surgeries on thyroid.
- Those on steroids.
- Those having chronic diseases like chronic renal failure, multiple sclerosis or malignancy.

Institutional ethical clearance will be received prior to the study.

Research tools:

Study participants will complete a questionnaire reflecting the formalin effects. Subject experts will validate the questionnaire responses, about symptoms of exposure to formalin like unpleasant smell, unusual thirsty feeling, itching and redness in eyes, headache, unusual tiredness, sore throat, itching of the hands. Symptoms will be graded by a scale of 1–4; grade (1): absent, grade (2): minimally recognizable (3): strong and grade (4): unable to tolerate.

Blood samples will be collected from all the study participants for Complete Blood Count (CBC) and thyroid function tests, which will be done at the Clinical Pathology department, Acharya Vinoba Bhave Hospital.

Procedure

5ml of blood will be collected from each participant and transported to Department of pathology for doing CBC and TFT by ELISA method. The name, age, sex will be recorded in the Proforma. Each participant will be given a specific code number. After entering the data in the excel sheet, names of student will be deleted.

Statistical analysis

The data will be typed onto computer files. The values of arithmetic mean (X), standard deviation (S), frequency will be calculated and Student's t test and Chi-square test will be applied to analyse data.

Table 1: Immediate exposure symptoms of formalin.

Symptoms of immediate exposure to formalin	Grade 0	Grade I	Grade II	Grade III
Unpleasant smell				
Unusual thirsty feeling				
Itching of eyes				
Dry, sore throat				
Redness of eyes				
Itching of the hands				
Headache				
Unusual tiredness and dizziness				

Table 2: Distribution of exposed and unexposed female students according to Complete Blood Count findings.

Complete Blood Count parameters	Formalin exposed students	Unexposed students	Test of significance
Hb			
RBC			
Ht			
MCV			
MCH			
MCHC			
Platelets			
WBC			

Table 3: Distribution of values of thyroid function tests in formalin exposed and unexposed students.

Parameters	Formalin exposed students	Unexposed students	Test of significance
T3			
T4			
TSH			

Discussion:

Medical Council of India has prescribed the standards of dissection hall in relation to its dimensions at all medical colleges. But issues like ventilation, exhaust appliances to reduce the exposure of formalin is not considered. In India, for ventilation air flowing through windows and doors is only considered. When exhaust fans and suction devices are provided, the capacity and ventilation rate of them should be considered.

In Japan, a medical school with dissection hall had 25 cadavers. Initially ventilation was provided by exhaust which has three outlets, at the rate of 247 m³/min. To reduce the exposure of formalin to staff and student(s), it was increased to 280 m³/min by adding local ventilation devices to every table in dissection hall. (11)

In Thailand medical school, the gross anatomy dissection hall measured 420 m³ had 18 cadavers. It had 2 doors, 4 windows and effective ventilation appliances like 10 air conditioners, 3 exhausts and 8 air cleaners to deodorize the hall. They were kept switched on before and after class at least for 30 minutes to minimize the exposure. (12)

In Poland, study reported that formaldehyde concentrations in the anatomy practical lab with cadavers as 0.47–0.57 mg/m³. Still medical students complained of various reactions like lacrimation was reported in about 85.9%, red eyes, dry and itchy eyes, runny nose, sneezing, and headache was observed in more than 50% students, cough and dry throat or throat irritation was the complaint found in 44% and 42% of students respectively (13).

Wineski and English (14) observed exposures of formaldehyde in an anatomy lab in a range of 0.07 to 2.95 ppm. It was observed that there was significant difference in formaldehyde concentration level in the cadaver storage room after opening of the tanks, among two institutes, which was actually about marginal before opening tanks. It was explained by difference in the total number of cadavers stored /unit floor area.

At Indian Medical College formalin levels was found as 0.11 to 1.07 mg/m³ in the storage room for cadavers whereas in the gross anatomy lab it was 0.06–1.12 mg/m³. That was above the upper limit of proposed guidelines laid down by concerned authorities, definitely harmful for health status of students and staff. (15)

A study in Japan found that level of formalin evaporated from cadavers was elevated at the beginning of dissection of cadaver; but, it crosses the guidelines for indoor chemical concentrations with respect to specified workplaces by 0.25 ppm each day of measurement except the last days of dissection.

Research done at the Department of Medicine and Physiology at the David Geffen School of Medicine at UCLA observed that environmental agents like formalin interfere with thyroid function at more than one sites. They can reduce circulating thyroid hormone levels impairing their actions; these agents may act as partial thyroid hormone agonists. They may trigger autoimmune thyroid diseases also. (16)

In a cross-sectional study conducted by Asha latha PR and Anilakumari V P, it was found that 15.1% formalin exposed person had hypothyroidism as compared to only 6.4% non-exposed persons. Formalin exposed group was comprised of staff and residents of Anatomy, Pathology and Forensic Medicine departments, while non exposed group included staff and residents of Physiology, Pharmacology, Community Medicine, Microbiology department. Each group was comprised of 120 participants. They observed values of T3 and T4 were slightly elevated, but within normal limits in exposed group as compared to non-exposed group. The value of TSH was slightly lower in exposed group.(17)

Elshaer N.S.M. and Mahmoud M.A.E. conducted the cross-sectional study to observe the acute effects of formalin on medical students and chronic effects on faculty members of Anatomy department. Medical students of Ist, IInd and IIIrd year were given a questionnaire to report the acute effects of formalin. They found that symptoms like unpleasant smell, dry or sore nose, unusual thirst, itching in the eyes, running or congested nose, redness in the eyes, excessive lacrimation, disturbance in sight and headache were present in more than 50% students. Whereas symptoms like syncope, dry or sore throat, nausea, gastrointestinal tract disturbances, unusual tiredness or dizziness, itching of the hands, skin eruptions on the face/neck, respiratory distress and disturbed nocturnal sleep were present in less than 50% students.

16 faculty and staff members of Anatomy department were the study participants as formalin exposed group for observing the chronic effects of formalin. More than 50% of the participants had more than 10 years duration of experience in the department. Among 16 participants, 7 were males and 9 were females. Comparison group was comprised of 19 staff members from Community medicine department as non-exposed group. They reported symptoms of skin disorders, such as burning, cracking, scaling, erythema, drying, eczema and allergic contact dermatitis in more than 50% and edema in less than 50% of exposed participants. These symptoms were totally absent in non-exposed individual.

Also formalin-exposed female participants reported anemia and menstrual disorders as 44.4% and 33.3% respectively as compared with 6.25% and 0.0% of the unexposed female participants respectively. Besides number of formalin-exposed females having history of spontaneous abortion, baby with congenital anomalies was more as compared to the unexposed females.

Regarding ocular disorders some of formalin exposed members had eye discomfort and irritation, corneal clouding and permanent alteration of vision. All formalin exposed participants reported upper and lower airway irritation. Bronchial asthma and its exacerbation were not encountered among the unexposed members. Nausea and gastro-intestinal tract hemorrhage was also found in some study members. One staff at the Anatomy department had cancer and received treatment.

Mean value of RBCs count, mean Ht%, mean value of platelets count was significantly lower while mean corpuscular volume, mean corpuscular hemoglobin and mean corpuscular hemoglobin concentration level was significantly higher among formalin exposed group as compared to non-exposed group. White blood cell count was also abnormal in exposed group (18). Different studies conducted among medical students on various health issues are available(19-25).

Interpretation

This study will help to point out the toxic effects of formalin on various systems, complete blood count and thyroid gland.

References:

1. Ghosh SK. Human cadaveric dissection: a historical account from ancient Greece to the modern era. *Anat Cell Biol* 2015;48:153-69.
2. Raja DS. Potential health hazards for students exposed to formaldehyde in the gross anatomy laboratory. *J Environ Health*. 2012;74:36-40.
3. International Agency for Research on Cancer IARC. Monographs on the evaluation of carcinogenic risks to humans 88, formaldehyde, 2-butoxyethanol and I-tert-butoxypropanolol-2-ol. In: IARC. 2006;36-325.
4. Binawara BK, Rajnee, Choudhary S, Mathur KC, Sharma H, Goyal K. Acute effect of formalin on pulmonary function tests in medical students. *Pak J Physiol*. 2010;6:8.
5. Hauptmann M, Stewart PA, Lubin JH, et al.. Mortality from lymphohaematopoietic malignancies and brain cancer among embalmers exposed to formaldehyde. *J Nat Can Inst*. 2009;1001:1696-1708.
6. Hauptmann M, Lubin JH, Stewart PA, Hayes RB, Blair A. Mortality from solid cancers among workers in formaldehyde industries. *Am J Epidemiol*. 2004;159:1117-1130.
7. Khaliq F, Tripathi P. Acute effects of formalin on pulmonary functions in gross anatomy laboratory. *Ind J Physiol Pharmacol*. 2009;53:93-96.
8. Keil CE, Akbar-Khanzadeh F, Konency KA. Characterizing formaldehyde emission rates in a gross anatomy laboratory. *Appl Occup Environ Hyg*. 2001;16:967-972.
9. Emue BE, Ayanniyi AA, Nwegbu MM, Ibekwe TS. Acute effects of formalin treated cadaver on Nigerian medical students. *Am J Trop Med Pub Health*. 2011;1:89-96.
10. Fischer MH. The toxic effects of formaldehyde and formalin. *The Journal of experimental medicine*. 1905 Feb 1;6(4-6):487.
11. Yamato H, Nakashima T, Kikuta A, Kunugita N, Arashidani K, Nagafuchi Y, Tanaka I. A novel local ventilation system to reduce the levels of formaldehyde exposure during a gross anatomy dissection course and its evaluation using real-time monitoring. *J Occup Health* 2005;47:450-3.
12. Saowakon N, Ngernsoungnern P, Watcharavitoon P, Ngernsoungnern A, Kosanlavit R. Formaldehyde exposure in gross anatomy laboratory of Suranaree University of Technology: a comparison of area and personal sampling. *Environ Sci Pollut Res Int* 2015;22:19002-12.
13. Uba G, Pachorek D, Bernstein J, Garabrant DH, Balmes JR, Wright WE, Amar RB. Prospective study of respiratory effects of formaldehyde among healthy and asthmatic medical students. *Am J Ind Med* 1989;15:91-101.
14. Wineski LE, English AW. Phenoxxyethanol as a nontoxic preservative in the dissection laboratory. *Acta Anat (Basel)* 1989;136: 155-8.
15. Gahukar S, Ramteke U, Majumdar D, Malviya R, Patil D, Trivedi J, Rao C, Kale AH. Prevalence of formaldehyde in indoor air of gross anatomy laboratory and cadaver storage room of a medical college. *J Environ Occup Sci* 2014;3:181-5.
16. Brent A *et al.* UCLA. Environmental Exposures and Autoimmune Thyroid Diseases. *Thyroid* 2010; Jul, 20(7): 755-761.

17. Asha latha PR and Anilakumari VP. Can occupational exposure to formaldehyde cause thyroid dysfunction? –a cross sectional comparative study. International journal of current advanced research 2019;8(1):17724-27.
18. Elshaer N.S.M. , Mahmoud M.A.E. Toxic effects of formalin-treated cadaver on medical students, staff members, and workers in the Alexandria Faculty of Medicine. Alexandria Journal of Medicine.2017;53:337-43.
19. Gajbhiye DR, Wadnerwar NN. Assessment of Awareness Regarding Heavy Metal Toxicity in School Children. JOURNAL OF PHARMACEUTICAL RESEARCH INTERNATIONAL. 2021;33(37B).
20. Hora, Srajan, Tanvi Jaiswal, and Seema Sathe. “Awareness among Post-Graduate Students about Ergonomic Practice in Vidarbha Region.” JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS 9, no. 5 (February 3, 2020): 271–74. <https://doi.org/10.14260/jemds/2020/61>.
21. Rathi, A., Ransing, R.S., Mishra, K.K., Narula, N., 2018. Quality of sleep among medical students: Relationship with personality traits. Journal of Clinical and Diagnostic Research 12, VC01–VC04. <https://doi.org/10.7860/JCDR/2018/24685.12025>
22. Keche, Harsha Atul, Preeti Prabhakar Thute, Ujwal Lehandas Gajbe, Atul Shankarrao Keche, and Darshna Gulabrao Fulmali. “Cadaveric Oath - Perceptions of First Year Medical Students.” JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS 9, no. 37 (September 14, 2020): 2722–25. <https://doi.org/10.14260/jemds/2020/591>.
23. Muley, Parikshit Ashok, Karan Ramyank Thakkar, Praveen Kanaram Gehlot, Ashok Jaykumar Vankudre, Pranjali Parikshit Muley, and Pradip Bhanudas Barde. “Association between Body Mass Index and Cognitive Functions in Medical Students.” JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH 14, no. 11 (November 2020): CC8–14. <https://doi.org/10.7860/JCDR/2020/45081.14256>.
24. Nimbalkar, Mayur Anil, and Swarupa Chakole. “Risk Factors of Non-Communicable Diseases (NCD) among Students of Junior College in Wardha District.” JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS 9, no. 43 (October 26, 2020): 3214–17. <https://doi.org/10.14260/jemds/2020/705>.
25. Sawarkar, Gaurav, Punam Sawarkar, and Vaishali Kuchewar. “Ayurveda Students’ Perception toward Online Learning during the COVID-19 Pandemic.” JOURNAL OF EDUCATION AND HEALTH PROMOTION 9, no. 1 (December 2020). https://doi.org/10.4103/jehp.jehp_558_20.