

# A COMPARATIVE EVALUATION OF THE MEASUREMENT OF INTERLEUKIN-1 $\beta$ AS SALIVARY BIOMARKER IN PERIODONTITIS PATIENTS WITH AND WITHOUT DIABETES MELLITUS

## ABSTRACT

**Introduction:** Periodontitis is characterized by inflammatory destruction of connective tissue, loss of periodontal attachment, and resorption of the alveolar bone. Because of its activity as an inflammatory mediator as well as a modulator of the extracellular matrix and bone, the pro-inflammatory cytokine interleukin (IL)-1 $\beta$ . The study included 30 patients, 15 males and 15 females, ranging in age from 30 to 60 years old. Group A: Clinically healthy, Group B: Periodontitis with diabetes mellitus, and Group C: Periodontitis without diabetes mellitus were the three groups of patients. Expectoration into sterile bulbs was used to collect whole unstimulated saliva from all patients. Following that, collected samples are sent to the laboratory, where the amount of IL-1 $\beta$  is determined using the ELISA sandwich technique and the data are statistically analysed using One-Way ANOVA. has been identified as a periodontal disease marker.

**Aim:** The study aims to evaluate Interleukin-1 $\beta$  as a salivary biomarker in periodontitis patients with and without diabetes mellitus.

**Materials and methods:** The study included 30 patients, 15 males and 15 females, ranging in age from 30 to 60 years old. Group A: Clinically healthy, Group B: Periodontitis with diabetes mellitus, and Group C: Periodontitis without diabetes mellitus were the three groups of patients under this investigation. Expectoration into sterile bulbs was used to collect whole unstimulated saliva from all patients. Following that, collected samples are sent to the laboratory, where the amount of IL-1 $\beta$ , as a periodontal disease marker was determined using the ELISA sandwich technique. and The data were statistically analysed using One-Way ANOVA. The Newman-Keuls multiple comparison test was used to test the significance at the levels of  $p < 0.05$ .

**Results:** IL-1 $\beta$  level was found to be significantly higher ( $p < 0.05$ ) in periodontitis with diabetes mellitus ( $75 \pm 7$  pg/L) when compared with patients with periodontitis only ( $45.06 \pm 7$  pg/L) and also when compared with healthy controls ( $36.37 \pm 5.6$  pg/L).

**Conclusion:** From the study, it was concluded that IL-1 $\beta$  level was significantly increased with periodontitis patients, and patients with periodontitis and diabetes mellitus, when compared to patients with clinically healthy group. In summary, diabetes mellitus and periodontal diseases are closely associated with each other. Salivary IL-1 $\beta$  level can be used as a biomarker in early diagnosis of chronic periodontitis in patients with diabetes mellitus.

**Keywords:** Interleukin-1 $\beta$ , Diabetes mellitus, Innovative technology, Periodontitis, Salivary biomarker.

## 1. INTRODUCTION

Periodontitis is characterized by the inflammatory destruction of the connective tissue, loss of periodontal attachment, and resorption of the alveolar bone. Though bacterial pathogens were needed to initiate the disease progress, it's evident that their presence alone isn't enough to cause the rate of tissue destruction in periodontitis [1]. Pathogenesis of the periodontal disease suggests that tissue destruction can result

from the change of host defense process by bacterial and host products, which stimulate the host inflammatory processes [2]. During the onset and course of periodontal disease, pro-inflammatory cytokines are a significant modulator of inflammation [3,4]. The pro-inflammatory cytokine interleukin (IL)-1 $\beta$  is involved in immunological modulation as well as a variety of inflammatory responses. Because of its activity as an inflammatory mediator as well as a modulator of the extracellular matrix and bone, the pro-inflammatory cytokine interleukin (IL)-1 $\beta$  has been identified as a periodontal disease marker[5,6]. Although both isoforms of IL-1 $\beta$  have similar biologic activity, IL-1 $\beta$  is more effective in stimulating bone resorption and is the type found more frequently in periodontitis [7].

Diabetes mellitus (DM) may be a group of metabolic disorders characterized by a high blood glucose level over a protracted period of your time-. Type I diabetes mellitus is an autoimmune disease whereby antigen presenting T cells selectively destroy insulin producing pancreatic  $\beta$ -cells. The activated T cells first invade the islets, leading to insulinitis. This is followed by destruction of islets, mediated by a complex interaction between activated lymphocytes, cytokines and macrophages [8]. Apoptosis is a fundamental process involved in destruction of  $\beta$ -cells [9]. More recently,  $\beta$ -cell apoptosis shows presentation of islet antigens for disease onset. B cell derived antigens are presented by antigen presenting cells such as dendritic cells in local pancreatic-draining lymph nodes.

Both periodontitis and diabetes are widespread and complex chronic illnesses with a well-established bidirectional link [10,11]. Diabetes is a key risk factor for periodontitis, with epidemiological studies indicating that it increases the risk threefold when compared to non-diabetic persons, especially if glycemic control is inadequate [12]. It's becoming obvious that there are interactions between the two diseases that have significant clinical consequences for dental practitioners' disease management. Periodontal treatment incorporating non-surgical periodontal therapy has been linked to improved glycemic control in diabetes patients, as well as a decrease in HbA1C. Infections from untreated periodontal disease raise blood sugar levels, making diabetes difficult to manage. For these reasons, in diabetes patients' periodontitis must be managed carefully. Our team has extensive knowledge and research experience that has translated into high quality publications.[13–25],[26–30] [31,32] The aim of the present study was to evaluate the measurement of IL-1 $\beta$  as a salivary biomarker in periodontitis patients with and without diabetes mellitus.

## 2. MATERIALS AND METHODS

Between December 2020 and February 2021, patients aged 30 to 60 years old who visited the department of periodontics at Saveetha dental college and hospitals in Chennai, (country???) were assessed. The test groups included thirty patients (15 males, 15 females) with pre-existing moderate to severe periodontitis.

### 2.1. Inclusion criteria

- Not more than two teeth missing in each quadrant;

- greater than or equal to 30% of periodontal sites with periodontal depth greater than or equal to 4 mm;
- greater than or equal to 20% of periodontal sites with interproximal clinical attachment loss >2 mm;
- greater than or equal to 30% of sites showing bleeding on probing; and
- radiographic evidence of bone loss are among the periodontitis enrollment criteria.

## 2.2. Exclusion criteria

- Individuals who had undergone periodontal treatment in the previous 6 months;
- history of medications [(antibiotics or anti-inflammatory drugs)] in the previous 6 months;
- smoking or use of tobacco in any form; history of alcoholism;
- betel nut users; history of known systemic diseases that would alter the healing response of the oral tissues;
- acute periodontal conditions, such as periodontal abscess and acute necrotizing gingivitis; and
- detection of periodontitis were all excluded.

## 2.3. Study design

The control group consisted of ten people of similar age, race, ethnicity, and sex who had fewer than 10% of sites with bleeding on probing, no sites with periodontal depth greater than or equal to 4 mm, no clinical attachment loss greater than 2 mm, and no radiographic evidence of bone loss visible in radiographs. [The control group in the abstract is healthy????? Also in the abstract 30 patients ??? Please check the study design ???](#)

Ten people with periodontitis and diabetes mellitus were chosen because they had not more than two teeth missing in each quadrant, had more than or equal to 30% of periodontal sites with periodontal depth greater than or equal to 4 mm, had more than or equal to 20% of periodontal sites with interproximal clinical attachment loss >2 mm, had more than or equal to 30% of sites with bleeding on probing, and had radiographic evidence.

## 2.4. Saliva collection

[Twelve 42](#) hours before saliva collection, participants were instructed to abstain from eating, drinking, and doing oral hygiene routines. Expectoration into sterile bulbs was used to collect whole unstimulated saliva from all patients. The samples were immediately placed on ice and transferred to the lab, where they were centrifuged for 10 minutes at 5,000 rpm and the clear supernatants were stored in aliquots at -70°C. Within three months of collection, the samples were thawed and the assay was carried out.

## 2.5. IL-1 $\beta$ analysis in saliva

A commercially available enzyme-linked immunosorbent assay kit that was specific for human IL-1 $\beta$  was used to assess IL-1 $\beta$  levels in saliva samples in duplicate [(Figure 1)]. The quantitative sandwich enzyme immunoassay approach was employed in this assay. The samples were diluted in a 1:4 ratio with the calibrator diluent included in the kit, and the assay was carried out according to the manufacturer's instructions. Each run included standards, and all findings were reported within the assay's linearity. The findings of the colorimetric reaction were read directly on the automatic microplate reader set to 450 nm as the optical density value. To achieve the actual concentration of IL-1 $\beta$ , the values obtained were

multiplied by the dilution factor. The results were reported as concentration of IL-1 $\beta$  in picograms per milliliter of sample.

## 2.6. Statistical Analysis

The mean  $\pm$  standard deviation of the triplicate analysis results of the experiments performed on control and treatment patients were expressed. The [data results](#) were statistically examined using Graph Pad Prism version 5's one-way analysis of variance [(ANOVA)] and the Newman-Keuls multiple comparison test to determine significant differences between the mean values. The results were considered statistically significant at the 0.05 level.



Fig 1: Human IL-1beta ELISA kit

## 3. RESULTS AND DISCUSSION

IL-1 $\beta$  level was found to be significantly higher ( $p < 0.05$ ) in periodontitis with diabetes mellitus ( $75 \pm 7$  pg/L) [when](#) compared with periodontitis [patients](#) only ( $45.06 \pm 7$  pg/L) and also when compared with healthy controls ( $36.37 \pm 5.6$  pg/L) (Table 1 and Figure 2). On the basis of age and sex, individuals with periodontal health were demographically similar to patients with periodontitis with and without diabetes mellitus,

although IL-1 $\beta$  levels were considerably different. All saliva samples from patients and controls had detectable amounts of IL-1 $\beta$ .

In this investigation, IL-1 $\beta$  was found in all samples from individuals with periodontitis, both those with and without diabetes [Table-1]. Miller et al and Gursikh et al reported similar results, detecting IL-1 $\beta$  in all saliva samples, including those from controls [33].


In a study conducted by Teles et al, patients with periodontitis had a three-fold greater level of IL-1 $\beta$  [34]. Salivary IL-1 $\beta$  is shown to be eight-fold higher in individuals with chronic periodontitis than in healthy controls, according to Rachna Kaushik et al. There could be differences in salivary IL-1 $\beta$  levels due to a variety of causes, including individual immunologic responses, the quality of local bacterial challenge, or the action of diverse combinations of inflammatory mediators, implying periodontal disease heterogeneity. Another explanation is that the local production of inflammatory mediators varies from site to site within the same patient [35].

IL-1 $\beta$  is a key pro-inflammatory cytokine that plays a role in periodontitis. Continuous bone loss may be produced by IL-1 $\beta$ , which is a potent activator of bone resorption. Increased IL-1 $\beta$  levels in periodontal patients with diabetes mellitus would lead to additional systemic diseases and problems, and individuals without diabetes but with periodontitis would be at risk of developing diabetes in the future [36]. Diabetes is a key risk factor for periodontitis; diabetics are three times more likely to develop periodontitis than non-diabetics. Salivary IL-1 $\beta$  level estimation can be employed as a non-invasive predictor for these diseases because there is a definite association between degree of hyperglycemia and severity of periodontitis.

The current study's shortcomings include a small sample size and regional restrictions. Variations in the study could be related to factors such as periodontitis stages and grading, saliva collection, and transportation. The study of salivary IL-1 $\beta$  in a wider population with various stages of periodontal disease and diabetes mellitus should be the focus of future research.

**Table- 1: comparison of salivary IL-1 $\beta$  levels among 3 groups (periodontitis patients-P , patients with periodontitis along with diabetes-P+DM, and patients with periodontal health-P). The values are expressed in pg /L**

Group	Periodontal health	P+DM	P	pvalue
IL-1 $\beta$ (pg/L)	36.37 $\pm$ 5.6	75 $\pm$ 7.0	45.06 $\pm$ 7.0	P<0.0001



**Fig 2.** Assessment of salivary caspase-9 concentration among periodontal health, periodontitis and periodontitis with diabetes mellitus. The levels of salivary caspase-9 were assessed by the Enzyme Linked Immunosorbent Assay (ELISA) method. Significance at  $p < 0.05$ , a- compared with the periodontal health group. b-compared with periodontitis with diabetes mellitus.

#### 4. CONCLUSION


Within the study limitations, we concluded that IL-1 $\beta$  level was significantly increased in periodontitis patients and periodontitis with diabetes patients compared to periodontally healthy individuals. In conclusion-, diabetes mellitus and periodontal diseases are intimately linked. IL-1 $\beta$  is potentially useful in distinguishing diabetes patients who are periodontally healthy and monitoring periodontal disease activity. Thus, salivary IL-1 $\beta$  level can be used as a biomarker in early diagnosis of periodontitis in patients with diabetes mellitus.

**COMPETING INTERESTS DISCLAIMER:**



Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

**REFERENCES: I suggest to remove all [Internet]..... No need to them. References should be written according to journal format.**

1. Delima AJ, Karatzas S, Amar S, Graves DT. Inflammation and Tissue Loss Caused by Periodontal Pathogens Is Reduced by Interleukin-1 Antagonists [\[Internet\]\(Internet\)](#). Vol. 186, The Journal of Infectious Diseases. 2002. p. 511–6. Available from: <http://dx.doi.org/10.1086/341778>
2. Dennison DK, Dyke TE. The acute inflammatory response and the role of phagocytic cells in periodontal health and disease [\[Internet\]\(Internet\)](#). Vol. 14, Periodontology 2000. 1997. p. 54–78. Available from: <http://dx.doi.org/10.1111/j.1600-0757.1997.tb00192.x>
3. Alexander MB, Damoulis PD. The role of cytokines in the pathogenesis of periodontal disease. Curr Opin Periodontol [\[Internet\]\(Internet\)](#). 1994;39–53. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/8032465>
4. Okada H, Murakami S. Cytokine Expression in Periodontal Health and Disease [\[Internet\]\(Internet\)](#). Vol. 9, Critical Reviews in Oral Biology & Medicine. 1998. p. 248–66. Available from: <http://dx.doi.org/10.1177/10454411980090030101>
5. Graves DT, Delima AJ, Assuma R, Amar S, Oates T, Cochran D. Interleukin-1 and Tumor Necrosis Factor Antagonists Inhibit the Progression of Inflammatory Cell Infiltration Toward Alveolar Bone in Experimental Periodontitis [\[Internet\]\(Internet\)](#). Vol. 69, Journal of Periodontology. 1998. p. 1419–25. Available from: <http://dx.doi.org/10.1902/jop.1998.69.12.1419>
6. Boch JA, Wara-aswapati N, Auron PE. CONCISE REVIEW Biological: Interleukin 1 Signal Transduction— Current Concepts and Relevance to Periodontitis [\[Internet\]\(Internet\)](#). Vol. 80, Journal of Dental Research. 2001. p. 400–7. Available from: <http://dx.doi.org/10.1177/00220345010800020101>
7. Masada MP, Persson R, Kenney JS, Lee SW, Page RC, Allison AC. Measurement of interleukin-1alpha and -1beta in gingival crevicular fluid: Implications for the pathogenesis of periodontal disease [\[Internet\]\(Internet\)](#). Vol. 25, Journal of Periodontal Research. 1990. p. 156–63. Available from: <http://dx.doi.org/10.1111/j.1600-0765.1990.tb01038.x>
8. Bunt M van de, van de Bunt M, Morán I, Ferrer J, McCarthy MI. Insights into Cell Biology and Type 2 Diabetes Pathogenesis from Studies of the Islet Transcriptome [\[Internet\]\(Internet\)](#). Frontiers in Diabetes. 2014. p. 111–21. Available from:

<http://dx.doi.org/10.1159/000362471>

9. Sun X, Yu W, Hu C. Genetics of Type 2 Diabetes: Insights into the Pathogenesis and Its Clinical Application [\[Internet\]\(Internet\)](#). Vol. 2014, BioMed Research International. 2014. p. 1–15. Available from: <http://dx.doi.org/10.1155/2014/926713>
10. Lamster IB. Diabetes Mellitus and Oral Health: An Interprofessional Approach [\[Internet\]\(Internet\)](#). John Wiley & Sons; 2014. 272 p. Available from: <https://play.google.com/store/books/details?id=FaY6AwAAQBAJ>
11. National Research Council, Institute of Medicine, Board on Health Care Services, Board on Children, Youth, and Families, Committee on Oral Health Access to Services. Improving Access to Oral Health Care for Vulnerable and Underserved Populations [\[Internet\]\(Internet\)](#). National Academies Press; 2012. 296 p. Available from: [https://books.google.com/books/about/Improving\\_Access\\_to\\_Oral\\_Health\\_Care\\_for.html?hl=&id=Uyl8jcuhs8IC](https://books.google.com/books/about/Improving_Access_to_Oral_Health_Care_for.html?hl=&id=Uyl8jcuhs8IC)
12. Hirschfeld J, Chapple ILC. Periodontitis and Systemic Diseases: Clinical Evidence and Biological Plausibility [\[Internet\]\(Internet\)](#). Quintessenz Verlag; 2021. 360 p. Available from: <https://play.google.com/store/books/details?id=T9UqEAAAQBAJ>
13. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. J Periodontol [\[Internet\]\(Internet\)](#). 2018 Oct;89[10]:1241–8. Available from: <http://dx.doi.org/10.1002/JPER.17-0445>
14. Paramasivam A, Priyadharsini JV, Raghunandhakumar S, Elumalai P. A novel COVID-19 and its effects on cardiovascular disease. Hypertens Res [\[Internet\]\(Internet\)](#). 2020 Jul;43[7]:729–30. Available from: <http://dx.doi.org/10.1038/s41440-020-0461-x>
15. S G, T G, K V, Faleh A A, Sukumaran A, P N S. Development of 3D scaffolds using nanochitosan/silk-fibroin/hyaluronic acid biomaterials for tissue engineering applications. Int J Biol Macromol [\[Internet\]\(Internet\)](#). 2018 Dec;120[Pt A]:876–85. Available from: <http://dx.doi.org/10.1016/j.ijbiomac.2018.08.149>
16. Del Fabbro M, Karanxha L, Panda S, Bucchi C, Nadathur Doraiswamy J, Sankari M, et al. Autologous platelet concentrates for treating periodontal infrabony defects. Cochrane Database Syst Rev [\[Internet\]\(Internet\)](#). 2018 Nov 26;11:CD011423. Available from: <http://dx.doi.org/10.1002/14651858.CD011423.pub2>
17. Paramasivam A, Vijayashree Priyadharsini J. MitomiRs: new emerging microRNAs in mitochondrial dysfunction and cardiovascular disease. Hypertens Res [\[Internet\]\(Internet\)](#). 2020 Aug;43[8]:851–3. Available from: <http://dx.doi.org/10.1038/s41440-020-0423-3>
18. Jayaseelan VP, Arumugam P. Dissecting the theranostic potential of exosomes in autoimmune disorders. Cell Mol Immunol [\[Internet\]\(Internet\)](#). 2019 Dec;16[12]:935–6. Available from: <http://dx.doi.org/10.1038/s41423-019-0310-5>



19. Vellappally S, Al Kheraif AA, Divakar DD, Basavarajappa S, Anil S, Fouad H. Tooth implant prosthesis using ultra low power and low cost crystalline carbon bio-tooth sensor with hybridized data acquisition algorithm. *Comput Commun* [\[Internet\]\(Internet\)](#). 2019 Dec 15;148:176–84. Available from: <https://www.sciencedirect.com/science/article/pii/S0140366419307017>
20. Vellappally S, Al Kheraif AA, Anil S, Assery MK, Kumar KA, Divakar DD. Analyzing Relationship between Patient and Doctor in Public Dental Health using Particle Memetic Multivariable Logistic Regression Analysis Approach [MLRA2]. *J Med Syst* [\[Internet\]\(Internet\)](#). 2018 Aug 29;42[10]:183. Available from: <http://dx.doi.org/10.1007/s10916-018-1037-z>
21. Varghese SS, Ramesh A, Veeraiyan DN. Blended Module-Based Teaching in Biostatistics and Research Methodology: A Retrospective Study with Postgraduate Dental Students. *J Dent Educ* [\[Internet\]\(Internet\)](#). 2019 Apr;83[4]:445–50. Available from: <http://dx.doi.org/10.21815/JDE.019.054>
22. Venkatesan J, Singh SK, Anil S, Kim S-K, Shim MS. Preparation, Characterization and Biological Applications of Biosynthesized Silver Nanoparticles with Chitosan-Fucoidan Coating. *Molecules* [\[Internet\]\(Internet\)](#). 2018 Jun 12;23[6]. Available from: <http://dx.doi.org/10.3390/molecules23061429>
23. Alsubait SA, Al Ajlan R, Mitwalli H, Aburaisi N, Mahmood A, Muthurangan M, et al. Cytotoxicity of Different Concentrations of Three Root Canal Sealers on Human Mesenchymal Stem Cells. *Biomolecules* [\[Internet\]\(Internet\)](#). 2018 Aug 1;8[3]. Available from: <http://dx.doi.org/10.3390/biom8030068>
24. Venkatesan J, Rekha PD, Anil S, Bhatnagar I, Sudha PN, Dechsakulwatana C, et al. Hydroxyapatite from Cuttlefish Bone: Isolation, Characterizations, and Applications. *Biotechnol Bioprocess Eng* [\[Internet\]\(Internet\)](#). 2018 Aug 1;23[4]:383–93. Available from: <https://doi.org/10.1007/s12257-018-0169-9>
25. Vellappally S, Al Kheraif AA, Anil S, Wahba AA. IoT medical tooth mounted sensor for monitoring teeth and food level using bacterial optimization along with adaptive deep learning neural network. *Measurement* [\[Internet\]\(Internet\)](#). 2019 Mar 1;135:672–7. Available from: <https://www.sciencedirect.com/science/article/pii/S0263224118311333>
26. PradeepKumar AR, Shemesh H, Nivedhitha MS, Hashir MMJ, Arockiam S, Uma Maheswari TN, et al. Diagnosis of Vertical Root Fractures by Cone-beam Computed Tomography in Root-filled Teeth with Confirmation by Direct Visualization: A Systematic Review and Meta-Analysis. *J Endod* [\[Internet\]\(Internet\)](#). 2021 Aug;47[8]:1198–214. Available from: <http://dx.doi.org/10.1016/j.joen.2021.04.022>
27. R H, Ramani P, Tilakaratne WM, Sukumaran G, Ramasubramanian A, Krishnan RP. Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris-A review. *Oral Dis* [\[Internet\]\(Internet\)](#). 2021 Jun 21; Available from: <http://dx.doi.org/10.1111/odi.13937>

28. Ezhilarasan D, Lakshmi T, Subha M, Deepak Nallasamy V, Raghunandhakumar S. The ambiguous role of sirtuins in head and neck squamous cell carcinoma. *Oral Dis* [\[Internet\]\(Internet\)](#). 2021 Feb 11; Available from: <http://dx.doi.org/10.1111/odi.13798>
29. Sarode SC, Gondivkar S, Sarode GS, Gadgail A, Yuwanati M. Hybrid oral potentially malignant disorder: A neglected fact in oral submucous fibrosis. *Oral Oncol* [\[Internet\]\(Internet\)](#). 2021 Jun 16;105390. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2021.105390>
30. Kavarthapu A, Gurumoorthy K. Linking chronic periodontitis and oral cancer: A review. *Oral Oncol* [\[Internet\]\(Internet\)](#). 2021 Jun 14;105375. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2021.105375>
31. Vellappally S, Abdullah Al-Kheraif A, Anil S, Basavarajappa S, Hassanein AS. Maintaining patient oral health by using a xeno-genetic spiking neural network. *J Ambient Intell Humaniz Comput* [\[Internet\]\(Internet\)](#). 2018 Dec 14; Available from: <https://doi.org/10.1007/s12652-018-1166-8>
32. Aldhuwayhi S, Mallineni SK, Sakhamuri S, Thakare AA, Mallineni S, Sajja R, et al. Covid-19 Knowledge and Perceptions Among Dental Specialists: A Cross-Sectional Online Questionnaire Survey. *Risk Manag Healthc Policy* [\[Internet\]\(Internet\)](#). 2021 Jul 7;14:2851–61. Available from: <http://dx.doi.org/10.2147/RMHP.S306880>
33. Miller CS, King CP, Chris Langub M, Kryscio RJ, Thomas MV. Salivary biomarkers of existing periodontal disease [\[Internet\]\(Internet\)](#). Vol. 137, *The Journal of the American Dental Association*. 2006. p. 322–9. Available from: <http://dx.doi.org/10.14219/jada.archive.2006.0181>
34. Teles RP, Likhari V, Socransky SS, Haffajee AD. Salivary cytokine levels in subjects with chronic periodontitis and in periodontally healthy individuals: a cross-sectional study [\[Internet\]\(Internet\)](#). Vol. 44, *Journal of Periodontal Research*. 2009. p. 411–7. Available from: <http://dx.doi.org/10.1111/j.1600-0765.2008.01119.x>
35. Kaushik R, Yeltiwar RK, Pushpanshu K. Salivary interleukin-1 $\beta$  levels in patients with chronic periodontitis before and after periodontal phase I therapy and healthy controls: a case-control study. *J Periodontol* [\[Internet\]\(Internet\)](#). 2011 Sep;82[9]:1353–9. Available from: <http://dx.doi.org/10.1902/jop.2011.100472>
36. Reinhardt RA, Masada MP, Johnson GK, DuBois LM, Seymour GJ, Allison AC. IL-1 in gingival crevicular fluid following closed root planing and papillary flap debridement [\[Internet\]\(Internet\)](#). Vol. 20, *Journal of Clinical Periodontology*. 1993. p. 514–9. Available from: <http://dx.doi.org/10.1111/j.1600-051x.1993.tb00400.x>