

Effect of extraction techniques and evaluation of antimicrobial activity of *Argemone Mexicana* leaves and roots extracts in different solvents

Abstract:

To estimate effect of extraction techniques and antimicrobial activity of different solvent extract of *Argemone mexicana* leaves and roots. *Argemone mexicana* leaves and roots used in this study was collected. Roots and leaves were shade dried, cut into small pieces of 2-3 inches and grounded in to powdered form using mixer grinder. Leaves and roots extracts were prepared using Soxhlet extraction and Microwave-assisted extraction techniques in solvents named acetone, methanol, aqueous. The antimicrobial activity of the roots and leaves extracts were evaluated against Gram +ve bacteria (*Xanthomonas campestris*, *Bacillus cereus*, *Staphylococcus aureus*) and fungal species (*Fusarium oxysporum*, *Macrophomina phaseolina* and *Candida albicans*) and their zones of inhibition in mm are measured by Agar well diffusion method. Soxhlet extraction technique gave better extract yield (g/100g) 12.19 and 8.54 while microwave-assisted extraction gave 8.88 and 6.94 for leaves and roots respectively. The methanolic root and leaves extracts exhibited higher antimicrobial activity followed by acetone and aqueous extracts. The result of the investigation showed that extraction techniques significantly affected extraction yield and antimicrobial activity. Soxhlet extraction is better one extraction method and methanolic extract of leaves was found to be good antimicrobial followed by acetone and aqueous.

Keywords: *Argemone mexicana*, Extraction, Roots, Leaves, Antimicrobial.

Introduction

Plants have long been known as approaching sources of special classes of chemical compounds, known as phytochemicals, having diverse biological and curative activities, which are effective in controlling or treating a variety of diseases. Plant-based traditional medicine system continues to play a vital role in the health care system with about 60 % of the world inhabitants relying mainly on traditional medicines for their primary health care (Khan and Bhadauria 2017). Modern knowledge on medicinal plant research still contains at least 25 % drugs and many others, which are synthetic analogues, built on prototype compounds isolated from medicinal plants. The ongoing growing recognition of medicinal plants is due to escalating faith in herbal medicine (S. Dutta *et al.*, 2014). The medicinal plant products, which are derived from plant parts such as stem, bark, leaves, fruits, roots and seeds have been part of phytomedicine that produce a definite physiological action on human body. Leaves and seeds are also reported to find application in maintaining normal blood circulation and cholesterol level in human body (Albuquerque *et al.*, 2007). These plant parts possess anti-venom property as well as antimicrobial property (Makhija and Khamar, 2010; Minu *et al.*, 2012). The most important of these natural bioactive constituents of plants are alkaloids, tannins, flavonoids and phenolic compounds (D. Chaudhuri *et al.*, 2012). *Argemone mexicana* Linn is an exotic weed has wide spread distribution in many tropical and sub-tropical regions in India. *Argemone mexicana* have many biologically important compounds, so it can be recommended as a plant of pharmaceutical importance (Khan *et al.*, 2019). This plant grows widely in all dried parts of subtropical India i.e. in Haryana, Madhya Pradesh, Uttar Pradesh, Punjab plains and North-Western part of India like Gujarat and Rajasthan. *Argemone mexicana* possess a wide-range of biological activities, such as antibacterial (Rahman *et al.*, 2011; Rahman M. *et al.*, 2006; Sahu *et al.*, 2012), antifungal (Kushtwar *et al.*, 2017; Singh *et al.*, 2009; More N. *et al.*, 2016; Andleeb S. *et al.*, 2020). Plant crude extracts were proved to be higher in antimicrobial activity as a synergistic effect than purified individual constituents. Plants screened for antimicrobial activities have provided modern medicine with abundance of drugs and treatments against various ailments (Mahesh *et al.*, 2008). Plants produce a host of antimicrobial agents, including a wide variety of natural defense compounds, such as phenolics, terpenoids, alkaloids, polyacetylenes and polypeptides. However, with the advent of modern antibiotic drugs mainly of bacterial, fungal and synthetic sources, many of these natural plant derived antibiotic compounds have been left unexplored. In the work herein, a comprehensive evaluation for effect of extraction techniques and evaluation

of antimicrobial activity of *Argemone Mexicana* leaves, roots extracts in different solvents that had not previously been assessed in the literature in Haryana.

Materials and methods

Plant materials - *Argemone Mexicana* roots and leaves were procured from the roadsides, dry region in Haryana in the end of the March 2021. Roots and leaves were brought to lab, cleaned using running tap water 2-3 times and then finally washed with distilled water followed by shade dried. After drying, roots and leaves were cut into small pieces of 2-3 inches and were ground in to powdered form using mixer grinder.

Chemicals and Reagents

HPLC grade solvents were used for extraction and evaluation of antimicrobial activity. All the solutions were prepared fresh and utilized on the same day of the assay and used for experimental procedures.

Extraction

For evaluation of antimicrobial activity extracts were prepared using different solvents named acetone, methanol and aqueous followed by two extraction methods.

Soxhlet Extraction

Four grams of powdered samples of *Argemone mexicana* roots and leaves were placed in a filter paper (Whatman No. 1) thimble in a classical Soxhlet apparatus fitted with a 250 mL round bottom flask. The acetone and methanol solvent were added up to one and a half siphons that are approximately 150 mL. After the completion of first extraction step of 5 h, residue in thimble was again extracted twice (each extraction time 2 and 1 h, respectively) with suitable amount of acetone and methanol solvent. Filtrates of acetone and methanol solvent from three extraction steps were pooled and their volumes were noted. But in the case of aqueous as a solvent it takes longer to extract through the siphon mechanism, requiring more time to complete the 7-8 cycles. longer. After extraction, the volume of each filtered solvent was measured. These extracts were filtered and used for evaluation of antimicrobial activity.

Microwave-assisted extraction

Extraction was carried out using a microwave oven (IFB, model : 2301) with output of 800W and operating frequency 2450 MHz. Eight gram of powdered samples of *Argemone mexicana* roots and leaves were placed in a 250 ml conical flask with solvents 100 ml each acetone, methanol and aqueous. Flasks were left overnight. Samples containing flasks were irradiated for 10 seconds in microwave oven at 40 per cent power and not allowed to boil. Then cooled to room temperature and irradiation step was repeated up to 12 times to complete the process. After extraction, the volume of each filtered solvent was measured and used for evaluation of antimicrobial activity.

Evaluation of Antimicrobial Activity

The antimicrobial activity of the roots and leaves extracts were evaluated by Agar well diffusion method described by Bayer *et al.*, (1966). Eighteen to 24 hrs single colonies on agar plates were used to prepare the bacterial suspension with the turbidity of 0.5 McFarland (equal to 1.5×10^8 colony-forming units (CFU)/ml). Turbidity of the bacterial suspension were measured at 600 nm. Agar plates were inoculated with 100 μ l of the test microorganisms and were spreaded uniformly with the help of spreader, then allowed to dry for 5 minutes. Mueller hinton agar plates and Potato dextrose agar were inoculated with bacterial strain and fungal strain respectively under aseptic conditions and wells (diameter=6mm) were filled with 50 μ l of the test samples and incubated at 37°C for 24 hours for bacteria and 72 hours for fungi. After the incubation period, the diameter of the growth inhibition zones was measured in mm. Zone around each well was measured after 24 h for bacteria and 72 h for fungi. All the experiments were performed in triplicate to reduce error. Cycloheximide was used as standard for fungi and for bacteria tetracycline was used. After incubation, zone of inhibition was measured in mm. The antimicrobial activity of root and leaves extracts obtained was tested against Gram +ve bacteria *Xanthomonas campestris*, *Bacillus cereus*, *Staphylococcus aureus* and fungal species *Fusarium oxysporum*, *Macrophomina phaseolina* and *Candida albicans* and their zones of inhibition in mm are measured.

Results and Discussion

Extract Yield:

Extract yield of *Argemone mexicana* roots, leaves prepared by soxhlet extraction and microwave-assisted extraction technique was given in Table 1. Among *Argemone mexicana* roots and leaves extracts prepared by two extraction technique, yield (g/100g) of extracts prepared by soxhlet extraction technique was highest than microwave-assisted extraction technique for the solvents aqueous followed by methanol and acetone. The results are in agreement with other researchers (Kaniya Mahour *et al.*, 2011; Datkhile Kailas, *et al.*, 2020). Among plant parts, extract yield of leaves was highest. Extraction yield is a measure of solvent and extraction method efficiency. Soxhlet extraction gave higher yield and results are in agreement with other researchers. Literature results showed that supercritical carbon dioxide(SC-CO₂) extraction give the lowest extraction yield as compared to soxhlet extraction for *Quercus infectoria galls* (Hasmida *et al.*, 2014). Soxhlet extraction was found to be 1.8 and 3 times higher than ultrasound extraction and maceration extraction but slightly (1.2 times) higher than microwave extraction in aerial parts of *Potentilla atrosanguinea* Lodd. Reported by kalia *et al.*, (2008).

Evaluation of Antimicrobial Activity

Extract obtained from soxhlet extraction technique suitable to better yield was used to evaluate antimicrobial activity. Among different solvent extracts, methanol extract showed good activity against bacterial culture *Bacillus sp.* and *Staphylococcus aureus* as well as antifungal activity against *Candida albicans*, *Fusarium oxysporum* and *Macrophomina phaseolina* as in Table 2,3 & Fig.1. Among leaves and roots extracts, leaves extracts showed better activity for antibacterial as well as antifungal in methanol. The antibacterial activity in terms of inhibition zone against *Bacillus sp.*, *Xanthomonas campestris* and *Staphylococcus aureus* was observed. The antifungal activity in terms of inhibition zone against *Candida albicans*, *Macrophomina phaseolina* and *Fusarium oxysporum* was observed. But the antifungal activity against *Fusarium oxysporum* was found to be nil in all leaves extracts.

Abdulkarim *et al.*, (2016) evaluated ethanol leaf extract that exhibit more potent antibacterial activity than methanol leaf extract in *Argemone mexicana*. Chloroform extract of seeds of *Argemone Mexicana* exhibited antibacterial activity with minimum inhibitory concentrations (MIC) of 2.0 -5.0 mg/ml, against both Gram-positive and Gram-negative bacteria reported by Singh *et al.*, (2009;). Bhattacharjee *et al.*, (2006) studied that methanol extracts of the leaves and seeds of the *A. mexicana* showed greater antibacterial activity than the corresponding water extracts. Methanol leaves extracts of *Argemone mexicana* at 100µl concentration showed better activity against two Gram positive (*Bacillus subtilis*, *Staphylococcus aureus*), four Gram negative bacteria (*Escherichia coli*, *Pseudomonas aeruginosa*, *Proteus vulgaris*, *Salmonella typhi*) and 4 fungi (*Aspergillus niger*, *Fusarium moniliforme*, *Candida albicans* and *Mucor plumbeus*) as reported by Shyam Prasad and Dhanapal, (2010). Nuclear magnetic resonance analysis of the root and leaf methanol fractions showed two main antibacterial compounds, chelerythrine and berberine and data highlight the importance of plants as an invaluable pharmaceutical resource at a time when antimicrobial and anticancer drug discovery had reported by Orozco *et al.*, (2021).

Research Content : The research content is original and has not been published.

Ethical Approval : Not Applicable

Data from other sources: Not applicable

Consent to publish : All the authors agree to publish the paper in Journal of Environmental Biology.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

- Abdul karim Kassem., YehiaAl Zomo., Ahmed Saif Moharem., Nahlah MN., Sallam Jameel Alafifi. (2016). Extraction, formulation and evaluation of *Argemone Mexicana* leaves as antimicrobial cream and ointment. *International journal of pharmaceuticals biosciences research*, **5** (4),37-51.
- Albuquerque, UP., Monteiro, JM., Ramosa, MA., Amorim, ELC.,(2007). Medicinal and magic plants from a public market in northeastern Brazil. *J Ethnopharmacol*, **110**,76-91.
- Andleeb, S., Alsalmeh A., Al-Zaqri N., Warad I., Alkahtani J., Bukhari SM. (2020). In-vitro antibacterial and antifungal properties of the organic solvent extract of *Argemone mexicana* L. *J King Saud Univ—Sci*,**32**,2053–2058.
- Aziz Mohammad Khan., Seema Bhadauria. ,(2019).Analysis of medicinally important phytochemicals from *Argemone mexicana*, *Journal of King Saud University Science*, **31**,**(4)**,1020-1026.
- A W Bauer, W M Kirby, J C Sherris, M Turck ; Antibiotic susceptibility testing by a standardized single disk method.*Am J Clin Pathol*,.Apr;**45**(4):493-6(1966).
- Bhattacharjee I., Chatterjee SK., Chatterjee S., Chandra G. (2006). Antibacterial potentiality of *Argemone mexicana* solvent extracts against some pathogenic bacteria. *Mem Inst Oswaldo Cruz*,. **101**, 645–648.
- Cowan MM. (1999). Plant Products as Antimicrobial Agents. *Clin Microbiol Rev*.**12**: 564–582 .
- Chaudhuri, D., Ghate, N. B., Sarkar, R., & Mandal, N. (2012). Phytochemical analysis and evaluation of antioxidant and free radical scavenging activity of *withania somnifera* root. *Asian Journal of Pharmaceutical and Clinical Research*, **5**(4), 193-199.
- Kailas Datkhile., Satish Patil., Madhavi Patil., Pratik Durgawale., Nilam Jagdale and Vinit Deshmukh.(2020). Studies on phytoconstituents, In vitro antioxidant, antibacterial, and cytotoxicity potential of *Argemone mexicana* L. *Journal of Natural Science, Biology and Medicine*, **11**(2).
- Hasmida, M.N.(2014). Effect of different extraction techniques on total phenolic content and antioxidant activity of *Quercus infectoria* galls. *International Food Research Journal*, **21**(**3**),1075–1079.
- Isha Singh., V.K. Madan., Satya Shree Jangra and Dushyant. (2018). Effect of Various Extraction Techniques and Solvents on Extract Yield and Sugars Content of Clove (*Syzygium aromaticum* L.) *Asian Journal of Chemistry*, **30** (**4**) 759-761.
- Kalpna Kalia., Kapil Sharma., Harsh Pratap Singh and Bikram Singh. (2008).Effects of Extraction Methods on Phenolic Contents and Antioxidant Activity in Aerial Parts of *Potentilla atrosanguinea* Lodd. and Quantification of Its Phenolic Constituents by RP-HPLC, *Journal of Agricultural and Food Chemistry* , **56** (**21**), 10129-10134.

Kanhiya Mahour., Ashok Kumar and V.S. Vihan. (2011).Effect on Physicochemical Characters and Bioactivity of *Argemone mexicana* in Different Plant Extraction Methods. *Journal of Advanced Laboratory Research in Biology*, 2, (4).

Khan, A. M and S. Bhadauria (2017). Antikeratinophilic activity of plant extracts: a review. *Int. J. Pharma. Biosci.* **8**, 726–734.

Aziz Mohammad Khan and Seema Bhadauria. (2019).Analysis of medicinally important phytochemicals from *Argemone Mexicana*. *Journal of King Saud University - Science*, 31, (4),1020-1026.

Kushtwar RS. (2017). Study on Antifungal Activity of Aerial Part of *Argemone mexicana* Linn. *World J Pharm Res.*1025–1031.

Mahesh, B., Satish, S. (2008).Antimicrobial activity of some important medicinal plants against plant and human pathogens. *World J Agric Sci*, **4**: 839-843 .

Makhija IK, Khamar D. (2010).Anti-snake venom properties of medicinal plants. *Der Pharmacia Lettre.* **2**:399-411.

M. Mashiar Rahman., Md. Jahangir Alam., Shamima Akhtar Sharmin., M. Mizanur Rahman1., Atiqur Rahman and M. F. Alam. 84 (2009). In Vitro Antibacterial Activity of *Argemone mexicana* L.(Papaveraceae). *Journal of Natural Sciences.* **8(1)** 77-80.

Minu V., Harsh V., Ravikant T., Paridhi J., Noopur S.(2012).Medicinal plants of Chhattisgarh with anti-snake venom property. *Int J Curr Pharm Rev Res.* **3**:1-10.

More N., Kharat A.(2016).Antifungal and Anticancer Potential of *Argemone mexicana* L. *Medicines*.**3**.28. pmid:28930138.

Orozco-Nunnally DA, Pruet J, Rios-Ibarra CP, Bocangel Gamarra EL, Lefeber T, Najdeska T. (2021). Characterizing the cytotoxic effects and several antimicrobial phytochemicals of *Argemone mexicana*. *PLoS ONE* **16(4)**: e0249704.

Rahman MdS., Salehin MdF., Mostofa Ja MdAH., Parvin A., Alam MdK. (2011). Antibacterial Activity of *Argemone mexicana* L. against Water Borne Microbes. *Res J Med Plant.* **5**, 621–626.

S. Dutta., P. Dey and T.K. Chaudhuri . (2014).*J. Pharmacogn. Phytochem.*, 3, 142.

Sahu MC., Debata NK., Padhy RN.(2012). Antibacterial activity of *Argemone mexicana* L. against multidrug resistant *Pseudomonas aeruginosa*, isolated from clinical samples. *Asian Pac J. Trop Biomed.*(**2**),800–807.

Santosh Kumar Singh., Vidya Dhar Pandey., Aradhana Singh., Chandan Singh. (2009). Antibacterial activity of seed extracts of *Argemone mexicana* L. on some pathogenic bacterial strains. *Afr. J. Biotechnol.* **8**,(24), 7077-7081.

Shyam Prasad G and Dhanapal R. (2010). Antibacterial and antifungal activity of methanolic extract of *Argemone mexicana* leaves. *International Journal of Phytopharmacology.* **1(2)**, 64-67.

Singh A., Singh S., Singh S., Singh T., Singh V., Pandey V. (2009) Fungal Spore Germination Inhibition by Alkaloids Dehydrocorydalmine and Oxyberberine. *J Plant Prot Res.* **49**.

Table 1. Extract yield (g/100g) of *Argemone Mexicana* roots, leaves prepared by two extraction technique

Plant & Location	Plant Parts Extraction Technique	Leaves			Roots		
		Aqueous	Methanol	Acetone	Aqueous	Methanol	Acetone
<i>Argemone Mexicana</i> & Hisar	Soxhlet	12.19	11.48	9.00	8.54	7.65	5.63
	Microwave	8.88	7.46	7.31	6.94	4.84	4.83
	Mean	10.5	9.47	8.15	7.74	6.24	5.23
	SE(m)	0.05	0.03	0.05	0.03	0.05	0.03
	CD at 5 %	0.16	0.19	0.16	0.12	0.16	0.12


Table 2. Antimicrobial activity of roots extracts of *Argemone Mexicana*

Plant extract	Antimicrobial activity (mm) roots					
	Antibacterial activity (mm)			Antifungal activity (mm)		
	<i>Bacillus</i> sp.	<i>Xanthomonas campestris</i>	<i>Staphylococcus aureus</i>	<i>Candida albicans</i>	<i>Fusarium oxysporum</i>	<i>Macrophomina phaseolina</i>
Methanol	13	18	18	34	30	32
Aqueous	12	08	12	28	28	18
Acetone	11	08	14	31	20	17
Tetracycline	21	16	18	-	-	-
Cycloheximide	--	--	--	13	15	10

Table 3. Antimicrobial activity of leaves extracts of *Argemone Mexicana*

Plant extract	Antimicrobial activity (mm) leaves					
	Antibacterial activity (mm)			Antifungal activity (mm)		
	<i>Bacillus</i> sp.	<i>Xanthomonas campestris</i>	<i>Staphylococcus aureus</i>	<i>Candida albicans</i>	<i>Fusarium oxysporum</i>	<i>Macrophomina phaseolina</i>
Methanol	21	16	18	38	-	32
Aqueous	11	11	11	9	-	11
Acetone	13	12	14	34	-	28
Tetracycline	21	16	18	-	-	-
Cycloheximide	--	--	--	13	15	10

Fig. 1. Antimicrobial activity of roots and leaves extracts of *Argemone Mexicana* - A) *Staphylococcus aureus* B) *Bacillus* sp C) *Xanthomonas campestris* D) *Macrophomina phaseolina* E) *Fusarium oxysporum* F) *Candida albicans*



UNDER PEER REVIEW