

Original Research Article

RADIONUCLIDE TRANSFER RATIO FROM SOIL TO CROPS IN SELECTED FERTILIZED FARMS AND ESTIMATION OF ITS HEALTH RISK IN RIVERS STATE OF NIGERIA.

Abstract

The activity concentration of Radionuclide in crops and soil of selected fertilized farmers of Rivers State and its transfer factor (TF) have been carried out using a well calibrated sodium iodide (NaI) detector. The activity concentrations of ^{40}K , ^{234}U , and ^{232}Th in crops and soils of the selected fertilized farmers of Rivers State have been determined. The radionuclide concentration in soil varies from 6.64 ± 0.56 – $164.11 \pm 4.51 \text{ Bqkg}^{-1}$ with control 3.91 ± 0.80 and 4.75 ± 1.24 , BDL– 25.58 ± 4.38 with control 2.58 ± 1.38 and 11.07 ± 0.81 and 0.29 ± 0.02 – 21.12 ± 1.20 and control 1.06 ± 0.22 and $0.38 \pm 0.06 \text{ Bqkg}^{-1}$ for ^{40}K , ^{234}U , and ^{232}Th and its control values respectively. The concentration in crops varies from 7.38 ± 0.70 – 494.41 ± 29.18 with control 89.22 ± 1.32 and 78.69 ± 9.37 , BDL– 46.36 ± 8.05 with control 10.48 ± 3.75 and $0.47 \pm 3.93 \text{ Bqkg}^{-1}$, and 0.22 ± 0.07 – $7.30 \pm 0.62 \text{ Bqkg}^{-1}$ with control of 0.40 ± 0.07 and $0.9 \pm 0.36 \text{ Bqkg}^{-1}$ for ^{40}K , ^{234}U , and ^{232}Th and its control values respectively. The activity concentration of radionuclides of ^{40}K , ^{234}U , and ^{232}Th were all higher than its respective control values, which show that the fertilized farms contain more radionuclide than the unfertilized farms (control) and this may be due to the long use of inorganic fertilizers during cultivation on the selected ADP farms. The transfer factor (TF) obtained from the selected fertilized farmers varies from 0.121 – 22.818 , BDL– 10.25 Bqkg^{-1} , and 0.023 – 5.03 Bq/kg for ^{40}K , ^{234}U and ^{232}Th respectively with mean of 7.60 Bqkg^{-1} , 1.62 Bqkg^{-1} and 1.913 Bqkg^{-1} . This variation may be due to differences in organic matter, soil content and soil of type. The mean transfer factor of ^{234}U (1.63) < ^{232}Th (1.91) < ^{40}K (7.600) show that although activity concentration of natural radio isotopes in the study area are high also the rate at which the radionuclides are transfer from soil to crops is also high. The higher value of transfer (TF) was obtained in ^{40}K from Gokana Local Government Area. This may be due to the long use of nitrogen–Phosphate–potassium (NPK) fertilizers within the farms and due to its high accumulation in soil and the higher uptake by plants within the study area.

Keywords: Transfer factor, Background, Activity concentration, ADP, Fertilized farm

Comment [U1]: The objective of the study is missing. It should be the first statement. e.g. A study was carried out to investigate in some selected local government in River state

Comment [U2]: The concentrations in the crops vary

Comment [U3]: concentrations

Comment [U4]: This statement is more of result and discussion. The statement of conclusion should be the last one

Comment [U5]: Replace Transfer factor with NPK

Comment [U6]: Remove "Background" and replace with Radionuclide transfer ratio

Comment [U7]: Remove ADP

1.

INTRODUCTION

Human environment contains various types of contamination which attracted the attention of research in our present society. Most of this contamination occur naturally or as a result of our daily activities here on earth. The crops obtained from this contaminated farms or environments accumulate radionuclides that could form a direct route of radiation exposure to human when ingested or through the consumption of the crops obtained from contaminate farms [9]. Natural radioactivity has always been present and broadly distributed in the earth's crust and the atmosphere, either as primordial radionuclides of uranium (^{238}U) and thorium(^{232}Th) decay series and radioactive potassium (^{40}K), or as cosmic radiations that are produced constantly in the atmosphere [11,5,4] Primordial radionuclides of ^{238}U and ^{232}Th decay series and ^{40}K which has extremely long half-lives which are of great concern in terms of radiation exposure due to their gamma ray emitting potentials. The uptake of radionuclides by plant roots constitutes the main pathways for the migration of radionuclides from the soil to plants and from plants to humans through consumption of these crops. Nigeria's population is on the creasing sides which in turn lead to an increase in crops production as a result of the high rate of crops (food) demand by the consumers, this increase in the crops demand (food production) as led to the consistence utilization of chemical fertilizers by farmers. The utilizations of phosphate fertilizer globally for the increased in crop productions and for land reclamation has risen to more than 30 million tons annually [3]. The major raw materials for the production of chemical fertilizers must therefore supply the essential nutrients necessary for plant growth and the plant transfer this chemical to the tubers. And these essential nutrients are Nitrogen, phosphorus and potassium. Natural radioactivity of mainly Uranium (^{238}U), Thorium (^{232}Th) and Potassium (^{40}K) seen in phosphate fertilizers emanate from the phosphate ore, (due to geological reasons) which is the main raw material used for phosphate fertilizer production. Transfer factor (TF) is a necessary parameter

Comment [U8]: Rephrase this and mind the punctuations. You may use grammar-correcting app to check the whole write-up

that encompasses influence of physicochemical properties of soil, environmental conditions, and types of radionuclides [2]. The migration of radionuclides in the soil to crops is complex and the transfer factor assessment models is commonly utilized to describe the translocation of radionuclides among different environmental matrices, it is defined as the ratio of the concentration of radionuclide in the destination matrix and that in the departure matrix. The transfer factor depends on chemical, physiological and ecological [17].

Rivers state environment is naturally blessed with different resources which has attracted industrials activities and farming activities. The industrials waste and operation of these industries could directly or indirectly contaminate the soil and crop from some of the local government area such as Khana, Gokana Eleme and Ahoada. Some of these contamination may be due to consistence agricultural cultivation on the farms, and the farms is gradually losing its yielding ability, which now led to the application of fertilizers by farmer in order to obtained high yield of crops. The utilization of chemical fertilizers increases the concentration of phosphate and uranium in the soil and also increases the concentration in nutrients. This could also have some harmful effect in the long run if not checked. The Consumption of crops containing high concentration of radionuclides may be dangerous to human health. The present of radionuclide in the consumed food will continues to irradiate the body as long as it remains radioactive and stays in the body (FAO, 1986). Researches have shown that any dose of radiation increases an individual risk of developing cancer. The concentration of radionuclide in the food chain and the continue consumption of such food adds to the cumulative risk of developing skin burn, leukemia, cancer and other related diseases [12].

The large data dispersion is also due to the fact that some of the ratios between two specific activities are in fact time-dependent, so that the assumption of an equilibrium model is not always valid. The research work aim at the determination of the radionuclide transfer factor (TF) from soil to crops in selected fertilized farms (ADP) of Rivers State, where crops are cultivated in large quantities for human consumption.

Comment [U9]: aims

Comment [U10]: indicate the full meaning of ADP

2. Study Area

This research work was carried out in selected fertilized farmers within six Local Government Area of Rivers State. figure1 shows the maps the study area. The following Local Government Area considered for research are; Ahoada, Emohua, Obio/Akpo, Eleme, Gokana and Khana Local Government Area and they are shown on the map with red colour. The samples collected from the selected Fertilized Famers were label with different code for proper identification during the analyze.

Comment [U11]: I don't think this makes sense. Do you mean Fertilized farm?

Comment [U12]: Areas

Comment [U13]:

Comment [U14]: the research were

Comment [U15]: analysis




Fig.1: Show the Map of Study Area

Comment [U16]: what does the circular red symbol represent?

3.1 Materials and Methods

3.1 Materials

Gamma-ray spectrometry system was use for analyzing the samples using Canberra 3 × 3 model 802. The Thallium-activated Sodium Iodide [NaI] detector and installed in a 100mm thick lead castle. The detector is connected to an amplifier linked to a computer program GENIE 2 K Window that correlated gamma energies to a number of possible isotopes. The sample was placed the marinelli beaker and then made to sit on the Sodium Iodide (NaI) detector. Shielding from background (environmental) radiation was achieved by counting in Canberra 100 mm thick lead castle. The energy resolution for the detector using Cs137 from International Atomic Energy Agency (IAEA) is 7.5% at 662KeV Cs-137 line. The standard and the sample were counted for a period of 36,000 seconds to acquire spectral data for a better counting statistics and evaluation. The activity concentration of ²³⁸U, ²³²Th and ⁴⁰K were determined after correction for background and inhomogeneity. The crops and soil samples were symmetrically placed on the top of the detector and the net area under the corresponding peaks in the energy spectrum was computed by subtracting counts due to Compton scattering of higher peaks and other background sources from the total area of the peaks. From the net area of a certain peak, the activity concentrations in the samples were obtained using the equation below

$$C(Bqkg - 1) = \frac{Cn}{\epsilon P \gamma M_s} \quad (1)$$

where C is the activity concentration of the radionuclide in the sample given in Bq kg⁻¹, Cn is the count rate under the corresponding peak, ϵ is the detector efficiency at the specific c-ray energy, P_c is the absolute transition probability of the specific c-ray, and M_s is the mass of the sample (kg).

3.2 Methods

3.2.1 Samples collection

The samples were collected from selected fertilized farms within three Senatorial district of Rivers State namely: Rivers-East Senatorial district, Rivers West Senatorial district and Rivers-South Senatorial district. Sixty-four (64) samples were collected for the purpose of the research work. Thirty-two samples were crops and the other thirty-two samples were soil. This samples were collected from six different fertilized farms of the selected Local Government Area within the Senatorial district. The following crops was collected from the selected fertilized farmers: Yam, Cassava, Coco-yam, Three leave yam, Maize and Potato.

Comment [U17]: districts

Comment [U18]: Areas

Comment [U19]: were

Comment [U20]: yam, cassava, cocoyam, three-leaf yam, maize and potato. Change the first letters to lower case

3.2.2 Sample Preparation

The 64 samples collected for the purpose of the study were properly washed with pipe-born water to remove mud and dust. The Crops were peeled and sun dried for twenty-eight days to get constant dry weight, and then grind with hand grinding machine and later packed into a new well label black polythene bags accordingly for proper identification. The maize samples collected from the selected fertilized farmers were detached from the cob, air dried for twenty-eight days to get a constant dry weight and then grind using hand grinding machine and packed into a new black polythene bags well label. The soil samples collected from the selected fertilized farms were filtered, sun dried and packed into a well label black polythene bags for proper identification.

Comment [U21]: pipe-borne

Comment [U22]: peeled

Comment [U23]: labelled

Comment [U24]: labelled

Comment [U25]: a proper identification

3.3 Radionuclide Uptake and Transfer factor

The earth contains varied degrees of radioactivity due to chain decay of ^{226}Ra and ^{232}Th (Saleh *et al.*, 2007). Natural radionuclides are in different concentrations in environmental media. Human activities like routine and accidental discharge of nuclear waste, production of energy, use of fertilizers and mining have altered their natural settings in the environment.

Comment [U26]: be consistent with your style of referencing

According to [13] absorption of radioisotopes is enhanced at the initial plant growth stage meaning that absorption varies with plant growth. The transfer factor depends also on the mass of plant. Equation 3 below expresses the dependence on mass by transfer factor.

$$TF(m) = TF(0) \left(\frac{m}{m_0} \right)^{\alpha-1} \quad (2)$$

Where m_0 is the initial plant mass, $TF(0)$ is the initial value of the transfer factor at $t = 0$, $m = m_0$, α is a function that determines the rate of decrease of transfer factor with increasing plant mass. Transfer factors can also be defined based on dry weight, as ratio of activity content ($\text{Bq}\cdot\text{kg}^{-1}$) in plant to activity content ($\text{Bq}\cdot\text{kg}^{-1}$) of soil or can be based on surface area of soil and expressed as $\text{Bq}\cdot\text{kg}^{-1}$ dry weight of plant to $\text{Bq}\cdot\text{m}^{-1}$ in soil [6] The recommended soil depth is 10 cm for grass and 20 cm for all other crops and trees. The radioisotope content at this depth is homogeneous. This transfer factor is expressed as:

$$TF = \frac{\text{activity concentration of plant (Bq.kg}^{-1}\text{ dry weight)}}{\text{activity concentration of soil (Bq.kg}^{-1}\text{ dry weight)}} \quad (3)$$

In this work, equation 2 was adopted because it was impossible to monitor the different plant growth stages to note mass of crops.

4. RESULTS AND DISCUSSION

Table 1: Specific Activity Concentration of ^{40}K , ^{234}U and ^{232}Th in Soil Samples from the selected Fertilizer Farms

S/N	LOCATION	SAMPLE CODE	^{40}K (Bq /Kg)	^{234}U (Bq/Kg)	^{232}Th (Bq/Kg)
1	Gokana	CVT1	6.64±0.56	9.73±2.06	2.43±0.24
2	"	CVT2	8.90± 0.23	2.52±0.44	2.77±0.20
3	"	CYMT	106.63±8.66	13.19±4.08	1.42±0.015
4	"	TLYMT1	124.73±9.76	10.45±2.27	2.42±0.23
5	"	TLYMT2	31.79±2.80	23.35±4.28	1.73±0.16
6	Khana	CON	56.53±5.02	16.22±3.22	1.15±0.12
7	"	YMT	115.07±9.30	14.14±2.79	21.2±0.22
8	"	TLYT	50.16±3.08	BDL	8.69±0.61
9	"	KPOT	171.01±14.59	23.43±5.09	10.00±0.90
10	"	KCVT1	55.53±4.80	BDL	4.32±0.38
11	"	KCVT2	27.56±1.68	BDL	0.41±0.03
12	Emohua	EMCVT	66.59±6.18	18.88±3.66	7.23±0.63
13	"	EMCYMT	168.59±12.73	4.04±0.87	0.68±0.07
14	"	EMCYMT2	81.88±6.56	8.94±1.84	3.08±0.28
15	Eleme	ELCVT1	161.15±9.69	3.22±0.49	0.29±0.02
16	"	ELCVT2	81.88±4.92	8.94±1.40	3.08±0.22
17	"	ELYMT	22.47±2.01	11.53±2.61	14.12±1.20
18	"	ELYMT2	99.38±8.63	15.21±3.09	5.34±0.50
19	Obio/Akpo	CVT1	72.63±6.31	22.99±4.44	6.86±0.59
20	"	CON	29.78±2.65	18.31±3.90	5.47±0.05
21	"	YMT1	30.9±2.80	25.58±4.38	3.06±0.28
22	"	YMT2	24.75±2.34	14.87±2.81	1.58±0.16
23	"	YMT3	169.64±6.71	5.03±0.57	1.58±0.09
24	Ahoada	CVT1	81.88±6.56	8.94±1.84	3.08±0.28
25	"	CVT2	164.11±4.51	4.32±0.59	0.89±0.01
26	"	CVT3	86.24±4.92	8.34±1.80	2.07±0.26
27	"	CYMT1	23.57±1.01	12.83±1.51	15.22±1.90
28	"	CYMT2	99.58±3.23	13.21±1.09	4.74±0.60
29	"	CON	82.23±2.30	24.19±2.24	6.86±0.59
30	"	TLY	29.38±1.35	19.41±1.90	7.40±0.08
	Mean		75.35±2.35	12.02±2.25	4.316±0.32
31	Control 1		3.91±0.80	2.58±1.38	1.06±0.22
32	Control 2		4.75±1.24	11.07±0.81	0.38±0.06




Fig .2: Concentration of ^{40}K in soil with standard





Fig .3: Concentration of ^{234}U in soil with standard



Comment [U27]: check "sampling points" and the concentrations as they are not well written. Correct for the others too

Fig. 4: Concentration of ^{234}U in soil with standard

Table 2: Specific Activity Concentration of ^{40}K , ^{238}U and ^{232}Th in Crops Samples from selected Fertilized Farms

Comment [U28]: correct this

S/N	LOCATION		^{40}K (Bq /Kg)	^{234}U (Bq/Kg)	^{232}Th (Bq/Kg)
1	Gokana	CVT1	183.55±11.00	BDL	2.17±0.16
2	"	CVT2	361.46±25.44	BDL	4.22±0.37
3	"	CYMT	494.41±32.39	6.85±1.40	1.61±0.15
4	"	TLYMT1	418.12±29.18	13.21±2.88	1.61±0.61
5	"	TLYMT2	212.24±15.32	16.58±3.75	0.80±0.07
6	Khana	CON	117.69±9.37	18.47±3.93	4.09±0.36
7	"	YMT	193.38±14.44	20.35±3.93	0.48±0.05
8	"	TLY	330.61±24.25	16.18±3.60	0.22±0.02
9	"	POT	447.60±29.24	10.57±2.31	21.74±1.90
10	"	CVT1	7.38±0.70	4.44±1.03	21.74±1.90
11	"	CVT2	248.87±18.18	16.58±3.75	0.80±0.07
12	Emohua	CVT	43.89±29.61	4.92±1.12	4.61±0.43
13	"	CYMT	74.10±6.78	28.11±5.58	4.41±0.40
14	"	CYMT2	369.42±24.13	7.57±1.65	5.06±0.43
15	Eleme	CVT1	19.51±1.75	BDL	6.02±0.52
16	"	CVT2	49.09±4.27	13.12±2.56	4.32±0.38
17	"	YMT	195.14±14.71	32.79±6.24	2.63±0.25
18	"	YMT2	293.73±22.01	15.96±3.25	1.33±0.16
19	Obio/Akpo	CVT1	171.34±13.47	26.91±4.82	0.71±0.08
20	"	CON	205.88±1.64	20.54±3.82	7.30±0.62
21	"	YMT1	238.74 ±18.15	46.36±8.05	1.27±0.02
22	"	YMT2	452.66±31.02	BDL	0.88±0.09
23	"	BYMT3	173.14±10.47	25.91±4.82	0.61±0.08
24	Ahoada	CVT1	204.66±1.14	22.14±3.82	6.40±0.62
25	"	CVT2	228.64 ±12.15	44.26±8.05	1.77±0.02
26	"	CVT3	451.33±11.02	26.14±3.05	1.08±0.09
27	"	CYMT1	418.12±29.18	13.21±2.88	1.61±0.61
28	"	CYMT2	218.14±12.02	17.28±1.65	2.80±0.07
29	"	CON	115.19±2.37	17.27±1.93	5.04±0.16
30	"	TLYMT	412.12±29.18	16.21±2.80	2.61±0.61
	Mean		218.14±23.28	17.28±3.11	2.80±0.31
31	Control 1		89.22±1.32	10.48±3.75	0.40±0.07
32	Control 2		78.69±9.37	0.47±3.93	1.09±0.36

Table 3: Specific Activity Concentration of Transfer Factor of ^{40}K , ^{238}U and ^{232}Th

S/N	Location	Sampling Code	^{40}K (Bq /Kg)	^{238}U (Bq/Kg)	^{232}Th (Bq/Kg)
1	Gokana	CVT1	27.64307	0	0.893004
2	"	CVT2	40.61124	0	1.523466
3	"	CYMT	4.636688	0.519333	1.133803
4	"	TYM1	3.352201	1.264115	0.665289
5	"	TYM2	6.676313	0.710064	0.462428
6	Khana	CON	2.081903	1.138718	3.556522
7	"	YMT	1.680542	1.43918	0.022642
8	"	TYT	6.591108	0	0.025316
9	"	POT	2.617544	0.451131	2.174
10	"	CVT1	0.132901	0	5.032407
11	"	CVT2	9.030116	0	1.95122
12	Emohua	CVT	0.659108	0.260593	0.637621
13	"	CYMT	0.439528	6.957921	6.488235
14	"	CYMT2	4.511724	0.846756	1.642857
15	"	CVT1	0.121067	0	20.75862
16	Eleme	CVT2	0.599536	1.467562	1.402597
17	"	EYMT	8.684468	2.843886	0.186261
18	"	YMT2	2.955625	1.04931	0.249064
19	Obio/Akpor	CVT1	2.35908	1.170509	0.103499
20	"	CON	6.913365	1.121791	1.334552
21	"	YMT1	7.726214	1.826635	0.415033
22	"	YMT2	18.28929	0	0.556962
23	"	YMT3	1.020632	5.151093	0.386076
24	AHOADA	CVT1	2.499511	2.47651	2.077922
25	"	CVT2	1.393212	10.24537	1.988764
26	"	AHCVT3	5.233418	3.134293	0.521739
27	"	CYMT1	17.7395	1.026418	0.105782
28	"	CYMT2	2.190601	1.3081	0.590717
29	"	CON	1.400827	0.713931	0.733624
30	"	TYMT	14.02723	0.835137	0.352703
	Mean		6.7939±2.21	1.5986±3.07	1.9324±2.10
31	Control 1		22.81841	4.062016	0.377358
32	Control 2		16.56632	0.042457	2.868421
	Control Mean		19.69237	2.052237	1.62289




Fig 8 Percentage of Transfer Factor of ^{40}K , ^{234}U and ^{232}Th

4.2 Discussion

The transfer factor (TF) obtained varies from 0.121-22.818, 0-10.25, and 0.023-5.03 Bq/kg for ^{40}K , ^{234}U and ^{232}Th respectively and mean value of 7.60, 1.62 and 1.913 Bq/kg. This variation may be due to differences in organic matter, soil content and soil of type. The higher value of transfer (TF) was obtained in ^{40}K from Gokana Local Government Area. This may be due to long use of nitrogen–Phosphate–potassium (NPK) fertilizers within the farms and also due to the high accumulation of radionuclide in soil which in turn lead to the high uptake by plants within the study area. Fig 8 showed the Percentage of radionuclide Transfer Factor of ^{40}K , ^{234}U and ^{232}Th , 68%, 17% and 15% respectively. The results obtained show that the Radionuclide within the soil samples of the selected fertilized farms are transfer to the crops. The lower value of transfer factor (TF) was recorded in ^{234}U and it did not agree with the work [1]. The result shows the opposite of this assumption for example, the average activity concentrate from Gokana cassava

Comment [U29]: why starting with an uppercase?

two was 8.90 Bqkg^{-1} with a transfer factor of 40.64. The percentage of Radionuclide is shown in Fig.8 and the higher percentage of transfer factor of Radionuclide was obtained in potassium 40 (^{40}K), Thorium 232 (^{232}Th) has the second higher value and Uranium (^{234}U) has the lowest value of transfer factor within the study area and it indicate that the Radionuclide within the fertilized farms are transfer to the tubers due to the application fertilizers during the cultivation period. The result of the transfer factor buttresses the fact that transfer factor are not linearly related to the soil concentration [10]. Several factors may affect the transfer factor such as the plant species, soil PH, types of plant, organic matter content, soil management practice and methods of application of fertilizers during the cultivation period. The result of transfer factor of the study area is higher than the transfer factors value reported by [8] potassium (^{40}K) has the highest mean value of transfer factor which properly may be due to its higher accumulation in soil and higher uptake by plant due to the consistence application of NPK fertilizers. The mean transfer factor of ^{234}U (1.63) < ^{232}Th (1.91) < ^{40}K (7.600) show that the activity concentration of natural radio isotopes in the study area are high and the rate at which the radionuclides are transfer from soil to crops is also high. According to [16] it is of great concern than care must be taken in the use of transfer factor to determine food safety for consumption. Transfer factor varies with location and plant type and it is assumed that plant concentration increases with an increase in soil concentration according transfer factor definition. The transfer factor (TF) result is higher than the value obtained by [16] [12] [7].

Comment [U30]: you don't have to repeat this

Comment [U31]: pH not PH

Comment [U32]: the methods

Comment [U33]: mind the punctuation

Comment [U34]: consistent

Comment [U35]: rephrase this

Comment [U36]: transfered

Comment [U37]: that

4.3 Conclusion

The transfer factor of the radionuclide (^{40}K , ^{234}U and ^{232}Th) within the selected government fertilized farms has been carried out and higher value of transfer (TF) was obtained in ^{40}K from Gokana Local Government Area. This high value of ^{40}K may be due to the long use of nitrogen – Phosphate–potassium (NPK) fertilizers within the fertilized farms (ADP) or due to its high accumulation in soil and the higher uptake by plants within the study area. The results obtained shows that the activity concentration of tuber samples were higher than the activity concentration of cereals samples, this may be as a result of the direct contact of the tubers with the soil, more so the tubers spent more time to mature than the cereal samples while the cereals samples does not have direct contact with the soil so the nutrient take time in traveling through the crops stems

Comment [U38]: rephrase

Comment [U39]: using NPK alone is ok

Comment [U40]: indicate the relevance of ADP to this study

Comment [U41]: show

before getting to the seed also the cereal does not spent much time within the fertilized farms and it mature faster than tubers crops.

Comment [U43]: rephrase and mind the punctuations

The percentage of Radionuclide is shown in Fig.8 and the higher percentage of transfer factor of Radionuclide was obtained in potassium 40 (^{40}K), Thorium 232 (^{232}Th) has the second higher value and Uranium (^{234}U) has the lowest value of transfer factor within the study area and it indicate that the Radionuclide within the fertilized farms are transfer to the crops. The result of the transfer factor buttresses the fact that transfer factor are not linearly related to the soil concentration.

Comment [U44]: start with lowercase

Comment [U45]: transferred

Comment [U46]: your conclusion is more of result and discussion. Rewrite and make it concise

References

Al-masri MS, Al-Akel B, Nashawani A, Amin Y, Khalifa KH, Al-Ain F (2008). Transfer of ^{40}K , ^{238}U , ^{210}Pb and ^{210}Po from soil to plant in various locations in South of Syria. *Journal of Environmental radioactivity*. 99(2):322-331.

Comment [U47]: choose a uniform font, spacing and the style of reference citing as required by the journal. Justify

Alsaffar, M. S., Jaafar, M. S., Kabir, N. A. and Ahmad, N (2015). Distribution of ^{226}Ra , ^{232}Th , and ^{40}K in rice plant components and physicochemical effects of soil on their transportation to grains. *Journal of Radiation Resources Applied. Science*, 8(3): 300-310.

Ei-Taher A, Abbady Adel GE.(2012). Natural radioactivity levels and associated radiation hazards in Nile river sediments from Aswan to El-minia, upper Egypt. *Indian Journal.*;50:224-230.

Ferdous J, Begum A, Islam A (2015) Radioactivity of soil at proposed Rooppur Nuclear Power Plant site in Bangladesh. *Int. J Radiat Res*, 13(2): 135-142.

Hasan MM, Ali M, Paul D, Haydar M, Islam S (2013) Measurement of Natural Radioactivity in Coal, Soil and Water Samples Collected from Barapukuria Coal Mine in Dinajpur District of Bangladesh. *Journal of Nuclear and Particle Physics*, 3(4): 63-71.

IAEA (International Atomic Energy Agency) (1994). The interception, initial and post deposition retention by vegetation of dry- and wet-deposited radionuclides, Vienna

Ilemona C. Okemea, Iyeh V. Suleb, Norbert N. Jibiric and Hamed O. Shittuc, (2016). Radioactivity Concentrations in Soil and Transfer Factors of Radionuclides (^{40}K , ^{226}Ra and ^{232}Th) from Soil to rice in Kogi state, Nigeria. *Archives of Applied Science Research*, 8 (6):34-38

John Actor Ocheje, Alexander A. Tyovenda (2020). Determination of the Transfer Factor and Dose Rate of radionuclides in Some Selected Crops in Kogi state, Nigeria. *IOSR Journal of Applied Physics (IOSR-JAP)* e-ISSN: 2278-4861. Volume 12,

- Leonid, L. Nkuba and Najat K., Mohammed (2014). Determination of radioactivity in maize and Mung beans grown in the neighborhood of Minjingu phosphate mine, Tanzania. *Tanz. J. Sci.* 40:51-59
- Ononugbo *et al.*, (2019). Carried out study on Radiological health risk due to consumption of cassava from Nigerian government farms (ADP)
- Samad MA, Haydar MA, Ali MI, Paul D, Bhuiyan MMR, Islam SMA (2012) A Study on the Radioactivity Level in Raw Materials, Final Products and Wastes of the Phosphate Fertilizer Industries in Bangladesh. *Journal of Environmental Protection*, 3(10): 1393.
- Svetlana, G. Gordana, V. Branislava M. Petrujkic, B (2010). Natural and Anthropogenic radioactivity of foodstuffs, mosses and soil in the Belgrade environment. *Arch. Biol. Sci.* 62(2):301-307
- Saeed, M.A., Yusof, S.S., Hossain, I., Ahmed, R., Abdullah, H.Y., Shahid, M., Ramli, A.T., (2012). Soil to rice transfer factor of the natural radionuclides in Malaysia. *Roman Journal of Physics*, 57 (9–10), 1417–1424
- Saleh, I. H., Hafe, A.F., Alanary, N.H., Motaoveh, H. A. and Naim, M.A. (2007). Radiological Study of Soils, Foodstuff and Fertilizers in the Alexandria region, Egypt, Turkey *Journal of Environmental Science*, 31, 9-17
- Sabbarese, C., Stellato, L., Cotrufo, M.F., D’Onofrio, A., Ermice, A., Lubritto, C., Terrasi, F., Alfieri, S. & Migliore, G. (2002b) Dependence of radionuclide transfer factor on plant growth stage. *Environmental Modeling and Software*, 17(6), 545-551
- Tchokossa P, Olomo J B, Balogun F A and Adesanmi C A (2013) Assessment of radioactivity contents of food in the oil and gas producing area in Delta State, Nigeria *International Journal of Science and Technology*. 3 245–5
- Whicker, F.W.; Hinton, T.G.; Orlandini, K.A.; Clark, S.B. (1999). Uptake of natural and anthropogenic actinides in vegetable crops grown on a contaminated lake bed. *J. Environ. Radioactivity*. 45: 1–12.