

# Periodontal status and glycaemic control among type 2 diabetic patients- a comparable study between 2 teaching hospitals in 2 geographical zones in Nigeria

## Abstract

**Background:** There is clinical evidence that periodontitis and diabetes mellitus (DM) are interconnected. Thus, periodontitis can promote systemic chronic inflammation that can exacerbate type 2 diabetes mellitus.

**Methodology:** Data was collected with self-administered questionnaire. Glycaemic control and periodontal status were evaluated by glycated Hb values and CPI respectively. Analysis was done using the Statistical Package for Social Sciences version 20.0 (IBM SPSS Statistics Armonk New York). Association of glycaemic index with periodontal status was explored by the  $\chi^2$  test and statistical significance was set at  $P < 0.05$ .

**Results:** One hundred and eighty-five participants with type 2 diabetes were recruited with a female predominance in both centers. Majority of participants were above the fifth decade. The mean duration of diabetes was  $8.97 \pm 7.14$ . 73.3 % of participants in UPTH and 84.7% in LASUTH were out of (p=0.001). One-fourth of participants in LASUTH and 14.2% of participants from UPTH had poor oral hygiene. Twice participants in LASUTH had good glycaemic control compared to those in UPTH (p=0.001). The periodontal status of majority of the participants in the two centers was between CPI score 2 and 4 (p=0.02). The association between good glycaemic control and gender and between good glycaemic control and age were statistically significant (p=0.014; p=0.001).

**Conclusion:** The periodontal status of participants did not worsen with poor glycaemic control. However, education was significantly associated with extent of control. Periodontal care should be incorporated into the management of the diabetics for improved quality of life.

**Key words:** Demographics, DM, Glycaemic control, Periodontal status.

## Introduction

Studies have reported an association between periodontal disease (periodontitis) and poor glycaemic control in diabetics.<sup>1-5</sup>

Periodontal disease (PD) is both a chronic infectious and inflammatory disease that is caused by the microorganisms that are found in dental plaque.<sup>1</sup> These microorganisms which are gram negative cause local inflammation that progresses from the gingiva to the alveolar bone resulting in destruction and loss of periodontal attachment through infiltration of inflammatory cells such as lymphocytes, macrophages and polymorphonuclear leukocytes into the periodontal tissues.<sup>2</sup>

These micro-organisms produce byproducts such as lipopolysaccharide (LPS) that activate macrophages and T-lymphocytes to produce pro-inflammatory molecules like interleukin-1 (IL-1), prostaglandin E2 (PGE2), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and lymphotoxin (LT) that has properties that are similar to those of TNF- $\alpha$ .<sup>2</sup> These cytokines are important in the mechanisms that produce periodontal tissue destruction through collagenolytic enzymes such as metalloproteinases (MMPs) that are activated by reactive oxygen leading to the resultant elevation of interstitial collagenase in inflamed gingival tissues.<sup>2,3</sup> The consequence is the resultant attachment loss with formation of periodontal pocket with millions of bacterial whose metabolism further worsens the initial destruction.<sup>4-7</sup>

Many recent studies have reported that the effect of PD might not only be in the oral cavity but involves the entire body system since the human body acts as a unity and biologic processes in one part of the body can affect other body areas.<sup>9-11</sup>

Diabetes mellitus (DM) on the other hand is a chronic disease that affects individuals in every age group . If poorly controlled, it can result in hyperglycaemia that can lead to a lot of complications in other organs of the body such as the heart, kidney and eyes.<sup>12</sup>

Increased glucose levels have been shown to cause non-enzymatic glycation and oxidation of proteins resulting in the accumulation of advanced glycation end products (AGEs) in diabetic tissues.<sup>13</sup> AGEs have receptors on cell surfaces [receptor for AGE (RAGE)], which they bind to causing various pathological changes.<sup>14</sup> The AGE-RAGE interaction on the macrophages surfaces induces the production of pro-inflammatory cytokines such as Tumour Necrosis Factor alpha (TNF- $\alpha$ ) and Interleukin -1 beta (IL-1 $\beta$ ).<sup>14,15</sup>

Diabetes is recognized presently as one of the two true risk factors (the other is smoking) for periodontal and has been recategorized into the new classification of periodontal diseases by the American Academy of Periodontology (AAP) and the European Federation of

Periodontology (EFP).<sup>16-17</sup> Furthermore, a bidirectional relationship has been established between PD and DM by several studies that concluded that periodontitis resulting from periodontal inflammation may be a complication of diabetes creating a negative effect on their glycaemic control by raising the blood glucose levels.<sup>18-20</sup>

This study therefore assessed the periodontal status and glycaemic control of diabetic patients.

## Methodology

This was a descriptive cross-sectional study done between July and September 2019 at two outpatient diabetic clinics of two Teaching Hospital in two zones in Nigeria (University of Port Harcourt Teaching Hospital (UPTH) in South-South zone of Nigeria and Lagos State University Teaching Hospital (LASUTH) in Lagos, South West zone in Nigeria).

The included subjects were patients with type 2 diabetes aged 18years and above who had been diagnosed with diabetes for at least 1 year before the study; those who gave consent to be part of the study and resided in the study locations. Excluded were diabetic patients with physical or mental challenges, those with chronic systemic diseases such as asthma and epilepsy, those on xerostomia causing drugs (anti-hypertensives, anti-depressants, diuretics and radiation therapy) and those who smoked or consumed alcohol. One hundred and eighty-five diabetics were recruited for this study; 105 from UPTH and 80 from LASUTH

Self-administered questionnaire was used to collect data on demographics. The oral cleanliness and periodontal status were assessed with Simplified Oral hygiene index by Greene and Vermillion and CPI index respectively. Values of glycated heamoglobin levels in the blood (HbA1c) values were retrieved from patients records and confirmed with laboratory

reports. Using the American Diabetes Association (ADA) guidelines,  $HbA1c < 7\%$   $HbA1c \geq 7\%$  were taken to indicate good and poor glycemic control respectively.<sup>21</sup>

### **Simplified Oral Hygiene Index (OHI-S).**<sup>22</sup>

OHI-S, a composite index that scores both debris and calculus deposition on selected teeth was developed by Greene and Vermillion in 1964. It is the sum of the mean debris index (DI-S) and calculus index (CI-S) of the examined teeth. The OHI-S is interpreted as good, fair or poor. (Score 1 (good oral hygiene) = 0.0 – 1.2, Score 2 (fair oral hygiene) = 1.3 – 3.0, Score 3 (poor oral hygiene) = 3.1 – 6.0).

The periodontal status was measured using a Community Periodontal Index according to the World Health Organization (WHO) recommendation for oral health surveys.<sup>23</sup> The scoring for the community periodontal index (CPI) are as follows:

**Code-0-** Coloured band of the probe remains completely visible in the deepest sulcus of the sextant-healthy.

**Code-1-** Coloured band of the probe remains completely visible in the deepest sulcus of the sextant, some bleeding after gentle probing.

**Code-2-** Coloured band of the probe still completely visible, but there is bleeding on probing, supragingival or subgingival calculus and/or defective margins.

**Code-3-** The coloured band is partially submerged. Pocket 4-5 mm deep.

**Code-4-** The coloured band completely disappears in the pocket, indicating a depth greater than 5.5 mm and a loss of attachment of 3mm or more.

Data analysis was done using the Statistical Package for Social Sciences version 20.0 (IBM SPSS Statistics Armonk New York). Continuous variables were described with mean and

standard deviation and nominal variables with frequencies. Association of glycaemic index with periodontal status was explored by the  $\chi^2$  test and statistical significance was set at  $P < 0.05$ .

## Results

The study population was one hundred and eighty-five patients with type 2 diabetes (105 from UPTH and 80 from LASUTH). Mean age was  $57.11 \pm 13.45$  year and mean DM duration was  $8.97 \pm 7.14$  years. There was a female predominance in both centers; UPTH (F:M of 1.76:1), LASUTH (F:M of 2.64:1). Table 1a.

Table 1b shows the mean parameters of the study participants

Figure 1 shows the clustered count bar of Hb1Ac by CPI scores. The CPI score among the two groups was majorly 2

Participants glycaemic control and periodontal status showed that twice participants in LASUTH had glycaemic control than those in UPTH (51.3% in LASUCOM vs 26.7 % of participants in UPTH had good glycaemic control); ( $p = 0.001$ ). One-fourth of participants in LASUTH had poor oral hygiene, while 14.2% of participants from UPTH had poor oral hygiene. More participants in UPTH had poor glycaemic control (73.3%). The periodontal status of majority of the participants in the two centers were CPI score between 2 and 4. Statistical analysis showed a statistical significance ( $p = 0.02$ ). Table 2

The association between participants' glycaemic control and periodontal status showed that the oral hygiene status of about two-third of participants with or without glycaemic control in

both centers was fair and the periodontal status showed a CPITN score of between 2 and 3 majorly. The CPITN score of participants with good glycaemic control showed a statistical significance ( $p=0.03$ ). Table 3.

Association between gender and good glycaemic control was statistically significant ( $p=0.014$ ). Likewise, the association between age group and good glycaemic control ( $p=0.001$ ) and poor glycaemic control (0.02). Table 4a.

The associations between glycaemic control and occupation ( $p=0.10$ ;  $p=0.16$ ) and between glycaemic control and education ( $p=0.10$ ,  $p=0.11$ ) showed no statistical significance. Table 4b.

## Discussion

The bidirectional relationship between PD and DM is well documented, though it is still unclear if it is a causal one or due to their common risk factors.<sup>17,24-27</sup> The interaction could be because diabetes may directly influence the oral microbes leading to dysbiosis or the common inflammatory pathways as inflammatory markers have been reported to be elevated in these two comorbidities. As such, studies have reported a positive effect on glycated haemoglobin levels in the blood (HbA1c) when periodontal therapy is done as it reduces the periodontal inflammatory load.<sup>11,17,25,26</sup> This study reported a good glycaemic control among a quarter and half of participants in UPTH and LASUTH respectively.

The mean age of the study population was  $57.11 \pm 13.45$  years and comparable to other studies done among diabetes.<sup>28</sup> More female predominance in this study comparable to some other studies.<sup>28-31</sup>

HbA1c, the glycosylated haemoglobin assay, is an indicator of blood glucose levels and a suitable prognostic marker widely accepted since the 1980s as the laboratory test of choice for determining glycaemic control in the diabetics.<sup>32</sup> It is still widely in use today and measurement is done using a number of different methods based on internationally adopted standards such as the Diabetes Control and Complications Trial (DCCT) or the International Federation of Clinical Chemistry (IFCC).<sup>33,34</sup> The latter consistently gives lower values (non-diabetic reference range is about 3% to 5% IFCC and 4% to 6% DCCT) with good control in diabetic groups taken as 5% IFCC and 7% DCCT. The American Diabetes Association (ADA) guidelines has however endorsed  $\text{HbA1c} < 7\%$  and  $\text{HbA1c} \geq 7\%$  as indicates of good and poor glycemic control respectively.<sup>21</sup>

The participants in this study based on Glycated hemoglobin were classified into three groups. These are well controlled (Glycated hemoglobin <7.0%) and poorly controlled (Glycated hemoglobin  $\geq$ 7.0%). Our study showed no association between duration of diabetes, glycated haemoglobin and periodontal disease severity similar to the findings in other studies.<sup>35,36</sup> This contrasted with other studies that reported association between higher glycated haemoglobin and severe periodontitis.<sup>37-42</sup>

In our study, participants with poor glycaemic control were more than those with good glycaemic control. This correlated with the findings of other studies.<sup>43-45</sup>

The Simplified oral hygiene index (OHI-S) measures the cleanliness of the mouth and is used to classify individuals into good, fair or poor oral hygiene. Majority of our participants had fair oral hygiene. This compares to the study done among T2DM in Lucknow, India that reported that 68.8% of their participants had fair oral hygiene.

The periodontal status of the participants in this study did not worsen with poor glycaemic control. This compares to the study done among diabetics in Harvard Medical School, Boston.<sup>46</sup> The means of participants parameters examined in this study are comparable to those reported by other studies.<sup>46,47</sup>

Community periodontal index is used to detect periodontal diseases. It scores the presence and absence of supra and sub gingival plaque and calculus as well as pocket depth correlating it with the extent and severity of the disease. Two-fifth and about half of those with well controlled and poorly controlled DM respectively had CPI score 2. However, a third of all participants irrespective of glycaemic control had CPI scores 3 and 4. This contrast with another study that recorded that two-fifth of their participants with poorly controlled DM had CPI code 3 and three-fifth had CPI code 4.



## **Conclusion**

The periodontal status of participants did not increase with poor glycaemic index as participants had CPI scores of 0-4 irrespective of Hb1Ac. However, the resultant increase in systemic inflammation shows the adverse effect of periodontal infections on diabetes mellitus and this can contribute to insulin resistance in those diagnosed with it.

## **Ethical Approval**

The ethical approval for the study was gotten from the ethics committees of the two hospitals (UPTH and LASUTH).

## **Consent**

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

## **Recommendation**

There is the need to increase patients' awareness of the link between diabetes mellitus and periodontitis and encourage collaboration between medical and dental professionals for the management of affected individuals.

## **Limitation**

This study did not assess gingival recession.

## **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement

of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## References

1. Negrato CA, Tarzia O, Jovanović L, Chinellato LE. Periodontal disease and diabetes mellitus. *J Appl Oral Sci.* 2013;21(1):1-12.
2. Sorsa T, Ingman T, Suomalainen K, Haapasalo M, Kontinen YT, Lindy O, et al. Identification of proteases from periodontopathogenic bacteria as activators of latent human neutrophils and fibroblast-type interstitial collagenases. *Infect Immun.* 1992; 60:4491–4495.
3. Lee W, Aitken S, Sodek J, McCulloch CA. Evidence of a direct relationship between neutrophil collagenase activity and periodontal tissue destruction in vivo: role of active enzyme in human periodontitis. *J Periodontal Res.* 1995; 30:23–33.
4. Geerts SO, Legrand V, Charpentier J, Albert A, Rompen EH. Further evidence of the association between periodontal conditions and coronary artery disease. *J Periodontol.* 2004; 75:1274–1280.
5. Kornman KS, Page RC, Tonetti MS. The host response to the microbial challenge in periodontitis: assembling the players. *Periodontol 2000.* 1997; 14:33–53.
6. Friedewald VE, Kornman KS, Beck JD, Genco R, Goldfine A, Libby P, et al. The American Journal of Cardiology and Journal of Periodontology editor's consensus:

- periodontitis and atherosclerotic cardiovascular disease. *J Periodontol.* 2009; 80:1021–1032.
7. Amar S, Han X. The impact of periodontal infection on systemic diseases. *Med Sci Monit.* 2003;9:RA291–RA299.
  8. Kapila YL. Oral health's inextricable connection to systemic health: Special populations bring to bear multimodal relationships and factors connecting periodontal disease to systemic diseases and conditions. *Periodontol 2000.* 2021;87(1):11-16.
  9. Abusleme L, Morandini AC, Hashizume-Takizawa T, Sahingur SE. Editorial: Oral Microbiome and Inflammation Connection to Systemic Health. *Front Cell Infect Microbiol.* 2021; 11:780182.
  10. Nazir MA, Izhar F, Akhtar K, Almas K. Dentists' awareness about the link between oral and systemic health. *J Family Community Med.* 2019 Sep-Dec;26(3):206-212.
  11. Borgnakke WS, Ylöstalo PV, Taylor GW, Genco RJ. Effect of periodontal disease on diabetes: Systematic review of epidemiological observational evidence *J Periodontol* 2013;84(4): S135–S152.
  12. Caton JG, Armitage G, Berglundh T, Chapple ILC, Jepsen S, Kornman KS, et al. A new classification scheme for periodontal and peri-implant diseases and conditions—Introduction and key changes from the 1999 classification *J Periodontol* 2018;89(1): S1–S8.
  13. Schmidt AM, Weidman E, Lalla E, Yan SD, Hori O, Cao R, et al. Advanced glycation end products (AGEs) induce oxidant stress in the gingiva: a potential mechanism underlying accelerated periodontal disease associated with diabetes. *J Periodontal Res.* 1996; 31:508–15.
  14. Lalla E, Lamster IB, Drury S, Fu C, Schmidt AM. Hyperglycemia, glycoxidation and receptor for advanced glycation end products: potential mechanisms underlying

- diabetic complications, including diabetes-associated periodontitis. *Periodontol* 2000. 2000; 23:50–62.
15. Lalla E, Lamster IB, Schmidt AM. Enhanced interaction of advanced glycation end products with their cellular receptor RAGE. Implications for the pathogenesis of accelerated periodontal disease. *Ann Periodontol*. 1998; 3:13–19.
16. Tonetti MS, Greenwell H, Kornman KS. Staging and grading of periodontitis: Framework and proposal of a new classification and case definition *J Periodontol* 2018;89(1): S159–S172.
17. Genco RJ, Graziani F, Hasturk H. Effects of periodontal disease on glycemic control, complications, and incidence of diabetes mellitus. *Periodontol* 2000. 2020; 83(1):59–65.
18. Winning L, Linden GJ Periodontitis and systemic disease: Association or causality? *Curr Oral Health Rep* 2017; 4:1–7.
19. Taylor GW. Bidirectional interrelationships between diabetes and periodontal diseases: an epidemiological perspective *Ann Periodontol* 2001;6(1):99–112.
20. Borgnakke WS. IDF Diabetes Atlas: Diabetes and oral health—A two-way relationship of clinical importance *Diabetes Res Clin Pract* 2019; 157:107839.
21. American Diabetes Association. Glycemic Targets: Standards of Medical Care in Diabetes. *Diabetes Care* 2018; 41(1): S55–S64.
22. Green JC, Vermillon JR. The simplified oral hygiene index. *JADA*.1964;68:7–13.
23. World Health Organization. Oral Health Surveys, Basic Methods. 4th ed. Geneva, Switzerland: World Health Organization; 1998:1–67.
24. Sanz M, Ceriello A, Buysschaert M, Chapple I, Demmer RT, Graciani F, et al. Scientific evidence on the links between periodontal diseases and diabetes: Consensus report and guidelines of the joint workshop on periodontal diseases and diabetes by the

International Diabetes Federation and the European Federation of Periodontology J  
Clin Periodontol 2018;45:138–149.

25. Genco RJ, Sanz M Clinical and public health implications of periodontal and systemic diseases: An overview. *Periodontol 2000* 2020;83:7–13.
26. Polak D, Saunui T, Nishimura F, Shapira L Diabetes as a risk factor for periodontal disease—plausible mechanisms. *Periodontol 2000* 2020;83(1):46–58.
27. Lakschevitz F, Aboodi G, Tenenbaum H, Glogauer M Diabetes and periodontal diseases: Interplay and links *Curr Diabetes Rev* 2011;7:433–439.
28. Rapone B, Ferrara E, Corsalini M, Qorri E, Converti I, Lorusso F, et al. Inflammatory Status and Glycemic Control Level of Patients with Type 2 Diabetes and Periodontitis: A Randomized Clinical Trial. *Int J Environ Res Public Health*. 2021;18(6):3018.
29. Grossi SG, Skrepinski FB, DeCaro T, Robertson DC, Ho AW, Dunford RG, et al. Treatment of periodontal disease in diabetics reduces glycated hemoglobin. *J Periodontol* 1997;68(8):713–719.
30. Telgi RL, Tandon V, Tangade PS, Tirth A, Kumar S, Yadav V. Efficacy of nonsurgical periodontal therapy on glycaemic control in type II diabetic patients: a randomized controlled clinical trial. *J Periodontal Implant Sci*. 2013;43(4):177–82
31. Santos VR, Lima JA, Miranda TS, Gonçalves TE, Figueiredo LC, Faveri M. Full-mouth disinfection as a therapeutic protocol for type-2 diabetic subjects with chronic periodontitis: Twelve-month clinical outcomes. A randomized controlled clinical trial. *Journal of Clinical Periodontology* 2013;40(2):155–162.
32. Simpson TC, Weldon JC, Worthington HV, Needleman I, Wild SH, Moles DR, et al. Treatment of periodontal disease for glycaemic control in people with diabetes mellitus. *Cochrane Database Syst Rev*. 2015;2015(11):CD004714.

33. Florkowski C. HbA1c standardization issues: should New Zealand follow the DCCT or the IFCC position? *N Z Med J* 2003;116(1171): U395.
34. Hanas R, John G. International HBA1c Consensus Committee. 2010 consensus statement on the worldwide standardization of the hemoglobin A1C measurement. *Diabetes Care* 2010;33(8):1903-1904.
35. Engebretson SP, Hyman LG, Michalowicz BS, Schoenfeld ER, Gelato MC, Hou W, et al. The effect of nonsurgical periodontal therapy on hemoglobin A1c levels in persons with type 2 diabetes and chronic periodontitis: a randomized clinical trial. *JAMA*. 2013 Dec 18;310(23):2523-32.
36. Kebede TG, Pink C, Rathmann W, Kowall B, Völzke H, Petersmann A, et al. Does periodontitis affect diabetes incidence and haemoglobin A1c change? An 11-year follow-up study. *Diabetes Metab*. 2018; 44(3):243-249.
37. Chiu SY, Lai H, Yen AM, Fann JC, Chen LS, Chen HH. Temporal sequence of the bidirectional relationship between hyperglycemia and periodontal disease: a community-based study of 5,885 Taiwanese aged 35-44 years (KCIS No. 32). *Acta Diabetol*. 2015 ;52(1):123-131.
38. Demmer, RT, Jacobs DR. Jr, Desvarieux M. Periodontal disease and incident type 2 diabetes: results from the First National Health and Nutrition Examination Survey and its epidemiologic follow-up study. *Diabetes Care* 2008; 31: 1373– 1379.
39. Ide R, Hoshuyama T, Wilson D, Takahashi K, Higashi T. Periodontal disease and incident diabetes: a seven-year study. *Journal of Dental Research* 2011; 90: 41– 46.
40. Morita T, Yamazaki Y, Mita A, Takada K, Seto M, Nishinoue N, et al. A cohort study on the association between periodontal disease and the development of metabolic syndrome. *J Periodontol* 2010; 81: 512– 519.

41. Saito T, Shimazaki Y, Kiyohara Y, Kato I, Kubo M, Iida M, et al. The severity of periodontal disease is associated with the development of glucose intolerance in non-diabetics: the Hisayama study. *J Dent Res* 2004; 83: 485– 490.
42. Xiong X, Elkind-Hirsch KE, Vastardis S, Delarosa RL, Pridjian G, Buekens P. Periodontal disease is associated with gestational diabetes mellitus: a case-control study. *J Periodontol* 2009; 80: 1742– 1749.
43. Preshaw PM, Alba AL, Herrera D, Jepsen S, Konstantinidis A, Makrilakis K, et al. Periodontitis and diabetes: A two-way relationship. *Diabetologia*. 2012; 55:21–31.
44. Winning L, Patterson CC, Neville CE, Kee F, Linden GJ. Periodontitis and incident type 2 diabetes: A prospective cohort study. *J Clin Periodontol*. 2017; 44:266–274.
45. Stoicescu M, Calniceanu H, Țig I, Nemeth S, Tent A, Popa A, et al. Significant aspects and correlation between glycemic control and generalized chronic periodontitis in type 2 diabetes mellitus patients. *Exp Ther Med*. 2021;22(1):671.
46. Shinjo T, Ishikado A, Hasturk H, Pober DM, Paniagua SM, Shah H, et al. Characterization of periodontitis in people with type 1 diabetes of 50 years or longer duration. *J Periodontol*. 2019 ;90(6):565-575. doi: 10.1002/JPER.18-0735.
47. Tinsley LJ, Kupelian V, D'Eon SA, Pober D, Sun JK, King GL, et al. Association of Glycemic Control with Reduced Risk for Large-Vessel Disease After More Than 50 Years of Type 1 Diabetes. *J Clin Endocrinol Metab*. 2017 Oct 1;102(10):3704-3711.

## Tables

**Table 1a. Participants' demographics**

Variables	Teaching Hospitals						$\chi^2$	p-value
	UPTH		LASUTH		Total			
	Freq	%	Freq	%	Freq	%		
<b>Sex</b>							<b>1.57</b>	<b>0.21</b>
Female	67	63.8	58	72.5	125	67.6		
Male	38	36.2	22	27.5	60	32.4		
<b>Age group</b>							<b>29.82</b>	<b>&lt;0.0001*</b>
20-29	2	1.9	2	2.5	4	2.2		
30-39	11	10.4	4	5.0	15	8.1		
40-49	30	28.6	3	3.7	33	17.7		
50-59	30	28.6	21	26.3	51	27.6		
60-69	19	18.1	22	27.5	41	22.2		
>70	13	12.4	28	35.0	41	22.2		
<b>Tribe</b>							<b>78.29</b>	<b>&lt;0.0001*</b>
Hausa	7	6.7	0	0.0	7	3.8		
Igbo	34	32.4	20	25.0	54	29.2		
Yoruba	10	9.5	53	66.2	63	34.0		
Rivers	34	32.4	0	0.0	34	18.4		
Others	20	19.0	7	8.8	27	14.6		
<b>Education</b>							<b>9.51</b>	<b>0.02*</b>
Informal	18	17.1	3	3.8	21	11.4		
Primary	19	18.1	22	27.5	41	22.2		
Secondary	27	25.7	25	31.2	52	28.1		
Tertiary	41	39.1	30	37.5	71	38.4		




<b>Occupation</b>							<b>15.09</b>	<b>0.005*</b>
Civil servant	26	24.8	15	18.7	41	22.2		
Retired	14	13.3	28	35.0	42	22.7		
Farmer	4	3.8	1	1.3	5	2.7		
Self-employed	56	53.3	36	45.0	92	49.7		
Professionals	5	4.8	0	0.0	5	2.7		
<b>Duration of diagnosis (years)</b>							<b>5.13</b>	<b>0.53</b>
1-5	43	41.0	31	38.7	74	40.0		
6-10	34	32.4	20	25.0	54	29.2		
11-15	17	16.2	14	17.5	31	16.8		
16-20	8	7.7	10	12.5	18	9.7		
21-25	1	0.9	3	3.8	4	2.2		
26-30	1	0.9	2	2.5	3	1.6		
>30	1	0.9	0	0.0	1	0.5		
<b>Total</b>	<b>105</b>	<b>100.0</b>	<b>80</b>	<b>100.0</b>	<b>185</b>	<b>100.0</b>		

Mean age = 57.11±13.45 years; Mean DM duration 8.97±7.14years

**Table 1b. Participants' characteristics**

Variables	Mean $\pm$ SD; %	Mean $\pm$ SD; %	Total
	Well controlled (Hb1AC <7%)	Poorly controlled (Hb1AC $\geq$ 7%)	
Age (years)	58.1 $\pm$ 16.0	56.52 $\pm$ 11.7	57.11 $\pm$ 13.46
Duration (years)	9.0 $\pm$ 7.8	8.9 $\pm$ 6.8	8.97 $\pm$ 7.17
BMI (kg/m <sup>2</sup> )	26.9 $\pm$ 5.1	27.6 $\pm$ 4.9	27.34 $\pm$ 5.00
Hb1Ac (%)	6.1 $\pm$ 0.6	9.5 $\pm$ 2.3	8.22 $\pm$ 2.49
Total Cholesterol	2.8 $\pm$ 1.1	3.1 $\pm$ 1.6	2.95 $\pm$ 1.44
LDL (mmol/L)	1.8 $\pm$ 1.0	2.1 $\pm$ 1.3	1.31 $\pm$ 1.99
HDL (mmol/L)	0.8 $\pm$ 0.6	0.9 $\pm$ 0.7	0.84 $\pm$ 0.62
Triglycerides (mmol/L)	1.5 $\pm$ 1.0	2.3 $\pm$ 1.7	1.99 $\pm$ 1.54
OHI-S	2.2 $\pm$ 1.1	2.3 $\pm$ 1.2	2.25 $\pm$ 1.12
CPI	2.2 $\pm$ 1.0	2.2 $\pm$ 0.9	2.19 $\pm$ 0.92



**Figure1. Community periodontal index scores with glycated haemoglobin**

**Table 2. Participants glycaemic and periodontal status**

Variables	Teaching Hospitals						$\chi^2$	p-value
	UPTH		LASUTH		Total			
	Freq	%	Freq	%	Freq	%		
<b>HbA1c</b>							<b>11.73</b>	<b>0.001*</b>
<7% (well controlled diabetes)	28	26.7	41	51.3	69	37.3		
≥7% (poorly controlled diabetes)	77	73.3	39	48.7	116	62.7		
<b>OHI-S</b>							<b>3.59</b>	<b>0.17</b>
Good	24	22.9	18	22.5	42	22.7		
Fair	66	62.9	42	52.5	108	58.4		
Poor	15	14.2	20	25.0	35	18.9		
<b>CPI scores</b>							<b>11.46</b>	<b>0.02*</b>
0	0	0.0	7	8.7	7	3.8		
1	14	13.3	14	17.5	28	15.1		
2	51	48.6	36	45.0	87	47.0		

3	30	28.6	19	23.8	49	26.5		
4	10	9.5	4	5.0	14	7.6		
<b>Total</b>	<b>105</b>	<b>100.0</b>	<b>80</b>	<b>100.0</b>	<b>185</b>	<b>100.0</b>		

UNDER PEER REVIEW

**Table 3. Association between glycaemic control and periodontal status**

Variables		Teaching Hospitals						$\chi^2$	p-value
		UPTH		LASUTH		TOTAL			
HbA1c		Freq	%	Freq	%	Freq	%		
<7%	OHI-S							1.07	0.59
Good	7	25.0	12	29.2	19	27.6			
Fair	17	60.7	20	48.8	37	53.6			
Poor	4	14.3	9	22.0	13	18.8			
Total	28	100.0	41	100.0	69	100.0			
≥7%	OHI-S							3.45	0.18
Good	17	22.1	6	15.4	23	91.8			
Fair	49	63.6	22	56.4	71	61.2			
Poor	11	14.3	11	28.2	22	19.0			
Total	77	100.0	39	100.0	116	100.0			
<7%	CPI scores							10.63	0.03*
0	0	0.0	3	7.3	3	4.4			
1	1	3.6	10	24.4	11	15.9			
2	12	42.8	18	43.9	30	43.5			
3	11	39.3	8	19.5	19	27.5			
4	4	14.3	2	4.9	6	8.7			
Total	28	100.0	41	100.0	69	100.0			
≥7%	CPI Scores							9.17	0.06
0	0	0.0	4	10.3	4	3.4			
1	13	16.9	4	10.3	17	14.7			

2	39	50.6	18	64.2	57	49.1			
3	19	24.7	11	28.2	30	25.9			
4	6	7.8	2	5.1	8	6.9			
<b>Total</b>	<b>77</b>	<b>100.0</b>	<b>39</b>	<b>100.0</b>	<b>116</b>	<b>100.0</b>			

UNDER PEER REVIEW

**Table4a. Association between participants glycaemic control and some demographics**

Variables	Teaching Hospitals						$\chi^2$	p-value
	UPTH		LASUTH		TOTAL			
HbA1c	Freq	%	Freq	%	Freq	%		
<7%	Gender						6.97	0.014*
Female	15	53.6	34	82.9	49	71.0		
Male	13	46.4	7	17.1	20	29.0		
Total	28	100.0	41	100.0	69	100.0		
≥7%	Gender							
Female	52	67.5	24	61.5	76	65.5	0.41	0.52
Male	25	32.5	15	38.5	40	34.5		
Total	77	100.0	39	100.0	116	100.0		
<7%	Age group (years)						19.64	0.001*
20-29	2	7.1	1	2.4	3	4.3		
30-39	5	17.9	3	7.3	8	11.7		
40-49	8	28.6	1	2.4	9	13.0		
50-59	7	25.0	7	17.1	14	20.3		
60-69	2	7.1	10	24.5	12	17.4		
>70	4	14.3	19	46.3	23	33.3		
Total	28	100.0	41	100.0	69	100.0		
Mean age± SD	58.10±16.02 years							
≥7%	Age group (years)						13.27	0.021*
20-29	0	0.0	1	2.6	1	0.9		
30-39	6	7.8	1	2.6	7	6.0		



40-49	22	28.6	2	5.1	24	20.7		
50-59	23	29.9	14	35.8	37	31.9		
60-69	17	22.0	12	30.8	29	25.0		
>70	9	11.7	9	23.1	18	15.5		
<b>Total</b>	<b>77</b>	<b>100.0</b>	<b>39</b>	<b>100.0</b>	<b>116</b>	<b>100.0</b>		
<b>Mean age± SD</b>	56.52±11.70years							
<b>&lt;7%</b>	<b>Year of diagnosis (years)</b>						<b>3.01</b>	<b>0.08</b>
0-5	16	57.2	17	41.5	33	47.9		
6-10	6	21.4	7	17.1	13	18.8		
11-15	4	14.3	9	21.9	13	18.8		
16-20	2	7.1	4	9.8	6	8.7		
21-25	0	0.0	3	7.3	3	4.4		
26-30	0	0.0	1	2.4	1	1.4		
<b>Total</b>	<b>28</b>	<b>100.0</b>	<b>41</b>	<b>100.0</b>	<b>69</b>	<b>100.0</b>		
<b>Mean duration± SD</b>	6.10± 0.57years							
<b>≥7%</b>	<b>Year of diagnosis (years)</b>						<b>1.25</b>	<b>0.54</b>
0-5	27	35.1	14	35.9	41	35.3		
6-10	28	36.3	13	33.3	41	35.3		
11-15	13	16.9	5	12.8	18	15.6		
16-20	6	7.8	6	15.4	12	10.3		
21-25	1	1.3	0	0.0	1	0.9		
26-30	1	1.3	1	2.6	2	1.7		
>30	1	1.3	0	0.0	1	0.9		
<b>Total</b>	<b>77</b>	<b>100.0</b>	<b>39</b>	<b>100.0</b>	<b>116</b>	<b>100.0</b>		

<b>Mean duration± SD</b>	9.48±2.34years		
--------------------------	----------------	--	--

UNDER PEER REVIEW

**Table4b. Association between participants glycaemic control and some demographics**

Variables	Teaching Hospitals						$\chi^2$	p-value
	UPTH		LASUTH		TOTAL			
HbA1c	Freq	%	Freq	%	Freq	%		
<7%	Education						6.24	0.10
Informal	3	10.7	1	2.4	4	57.9		
Primary	3	10.7	14	34.1	17	24.6		
Secondary	8	28.6	9	22.0	17	24.6		
Tertiary	14	50.0	17	41.5	31	44.9		
Total	28	100.0	41	100.0	69	100.0		
≥7%	Education						5.96	0.11
Informal	15	19.5	2	5.1	17	14.7		
Primary	16	20.8	8	20.5	24	20.7		
Secondary	19	24.7	16	41.0	35	30.2		
Tertiary	27	35.1	13	33.3	40	34.5		
Total	77	100.0	39	100.0	116	100.0		
<7%	Occupation						7.50	0.11
Civil servant	5	17.9	7	17.1	12	17.4		
Retired	5	17.9	18	43.9	23	33.3		
Farmer	1	3.5	1	2.4	2	2.9		
Self-employed	15	53.6	15	36.6	30	43.4		
Professionals	2	7.1	0	0.0	2	2.9		
Total	28	100.0	41	100.0	69	100.0		
≥7%	Occupation						6.59	0.16

Civil servant	21	27.3	8	20.5	29	25.0		
Retired	9	11.7	10	25.6	19	16.4		
Farmer	3	3.9	0	0.0	3	2.6		
Self-employed	41	53.2	21	33.9	62	53.4		
Professionals	3	3.9	0	0.0	3	2.6		
<b>Total</b>	<b>77</b>	<b>100.0</b>	<b>39</b>	<b>100.0</b>	<b>116</b>	<b>100.0</b>		

UNDER PEER REVIEW