

# Impact Assessment of Front-Line Demonstration (FLDs) on The yield of Green gram

## ABSTRACT

The green gram cluster frontline demonstrations were conducted by Krishi Vigyan Kendra, Jalgaon Jamod during 2016-17, 2017-18 and 2018-19 in Kharif season and covered 30 ha of land with 75 demonstrations in total 15 villages in 2 clusters of Buldana District of Maharashtra. In front-line demonstrations, improved quality seed variety, seed treatment, recommended fertilizers doses, Rhizobium culture, PSB bio-fertilizers and crop protection management approaches were exhibited in farmers' fields. The results showed that the highest grain yield was obtained in the demonstration plot with an average of 6.15 q/ha compared to 4.66 q/ha in the farmers' practice. A higher average net return (Rs.19,712/ha) was obtained in the demonstration plots compared to the farmers' plot (Rs.13,250/ha). The average B: C ratio was 2.10 in demonstrated plot compared to 1.80 in farmers plot. The average increase in demonstration yield over farmers practice was 32 percent and the increase in net return over farmer practice was 49 %. This is due to both improved technology and improved varieties. The level of return was significantly low for local practices due to a significant difference in the extent of adoption of recommended practices versus the level of risk in terms of cost-benefit, skill and knowledge of the practices involved.

**KEYWORDS:** Greengram, FLD, Technology and Extension gap, Technology Index and Economic return

---

## INTRODUCTION

'Green gram (*Vigna radiata* L) is the third most important pulse crop in India. Green gram has strong root system and capacity to fix the atmospheric nitrogen into the soil and improves soil health and contributes significantly to enhancing the yield of subsequent crops' (Meena et al, 2012). Its seeds are transferred more nutritious, more digestible and non-flatulent than other legumes grown in the country. 'The Indian Institute of Pulses Research, Kanpur, has projected the country's demand for pulses to reach at 39 million tonnes by 2050, which will require pulses

production to grow at an annual rate of 2.2%' (IIPR, 2015). To meet the rising demand, the country must both produce enough pulses and competitiveness to defend domestic production. To achieve this, more efficient crop production technologies, as well as favorable policies and market support, must be developed and implemented to encourage farmers to commit more acreage to legumes. In light of the foregoing, an extensive intervention, including cluster front-line demonstrations have been organized in the Buldana district of Maharashtra to transfer the farm technology developed to enhance productivity and profitability. The yield gaps of technological extension in legumes in this comprehensive study are also presented in this paper appropriate extension strategy for effective technology transfer to target farmers in Buldana district to introduce and disseminate improved varieties of green gram (BM-2003-2).

## MATERIAL AND METHODS

. Krishi Vigyan Kendra, Jalgaon (Ja.) Dist. Buldana, Maharashtra, in 2016-17, 2017-18 and 2018-19 conducted a Cluster Frontline Demonstration (CFLD) on improved production technologies in 12 villages of two blocks of Sagrampur Tahsil, Buldana dist, Maharashtra for cultivating green gram (*Vigna radiata* L) through recommended practice packages; whereas in control plots, crops were grown according to agricultural practice generally adopted by individual farmers. Soils of the CFLD plots were poor in N and moderate in P and K availability. The primary data on grain yield farmers' practices were collected from beneficiary farmers through the crop cutting method followed by face-to-face interviews. Farmers were educated on preferred technology for a demonstration before the key decision was taken. Improved quality seeds of green gram (BM-2003-2), seed treatment, appropriate fertilizer dose, use of Rhizobium and PSB bio-fertilizers and crop protection management techniques were demonstrated in the farmers' fields through front-line demonstrations at various locations. The usual practices were maintained for the on-site inspections. All major farm operations were conducted under the supervision of KVK scientists through regular visits. The yield increase in demonstrations over farmers' practice was calculated by using the following formula:

Extension Gap (q/ha) = Demonstration Yield – Check Yield

Technology Gap (q/ha) = Potential Yield – Demonstration Yield

Technology Index (%) = Technology Gap / Potential Yield X 100

## RESULTS AND DISCUSSION

Technologies implemented in the **Front-Line** Demonstration cluster and practices adopted by farmers are presented in Table 1

### Grain Yield

Grains of green gram under improved practices and general farming practices are presented in Table 2 and Table 3, which clearly show that the use of high-yielding variety helps to increase the productivity of green gram under rainfed conditions. The average grain yield of green gram under improved practices ranged from 4.31 to 8.80 q/ha which is 23.47 to 29.07 percent higher than farmers' practices. In frontline demonstration plots, there was an average yield increase of 26.58 % in farmers' practices. The findings support those of Singh et.al. (2012), Kaur et al. (2014), Lalit et al (2015) and Kumar & Kispotta (2017), who all observed an increase in green gram grain yield in front line demonstration plots.

### Technology gap

The average green gram technology gap ranged from 1.20 to 5.69 q/ha. The average technology gap of 3.85 q/ha for green gram was reported in the present study. The larger technological gap can be mainly attributed to the uneven distribution of rainfall, differences in soil fertility and peripheral cultivation and locally specific crop management problems faced to exploit the yield potential of certain cultivars on demonstration plots.

### Extension gap & Technology index

The extension gap varied from 0.82 q/ha to 2.44 q/ha during the study period. The average extension gap was 1.49 q/ha the technology index varied from 12.00 to 56.90 showing the feasibility of the farmer field's developed technology. The lower value of **the** technology index, the more the feasibility is proven where as poor field establishment in **the** early vegetative stage due to water stress in rainfed agriculture with uneven distribution of rainfall, long dry period and increasing pressure from diseases and insect pests are the possible reason for poor yields causing higher technology index. Similar results were also reported by was also reported by Lalit et al. (2015) and Singh et al. (2017).

## Economic and Front-Line Demonstration

Gross return and net return costs of cultivation depicted in Table 3 and show that cultivation costs range from Rs. 13,466 to Rs 20,742 / ha with average Rs. 17,753 per hectare under FLDs, but farmer conditions with an average of Rs 16,292 per hectare. During 2016-17, the maximum net return of Rs 25,823 per hectare was achieved under the frontline demonstration. The suggested practices had a greater average benefit-cost-ratio (2.10) than a general farmer practice (1.80) was obtained. The findings are consistent with Raj et al (2013) and Kumar et al (2015) research.

## CONCLUSION

Green gram grain yield was higher under improved techniques than under farmer practices, increasing yield per unit area while also increasing farmers' income. However, due to technology and extension gaps there is a large gap in potential yields, demonstration yields and farmers' plot yields. This suggests for the need for appropriate dissemination of site-specific technologies embedded in high yielding varieties to increase productivity and profitability of rainfed agriculture to improve the agriculture of Maharashtra.

## REFERENCES

- IIPR.2015 Vision. Indian Institute of Pulses Research Kanpur, Uttar Pradesh, India;2050:1-50.
- Kaur, P., Kaur, A., Kaur, B. and Singh, K. 2014. Performance of front line demonstration on summer moong in Jalandhar district. *J. Krishi Vigyan*, **3** (1): 58-61.
- Kumar, P., Singh, K. and Kaur, P. 2015. Economic impact of frontline demonstration on pulses in Punjab a step towards diversification. *Indian J. Econo. & Develop.*, **11** (1): 111-116.
- Kumar, S. and Kispotta, W. 2017. Impact of front line demonstration on summer moong through improved technology in Bhadohi district of Uttar Pradesh, India. *Plant Archives*, **17** (1): 423-426.
- Lalit, M., Patil, D.J., Modi, H.M., Vasava and Gomkale, S.R. 2015. Evaluation of Front line demonstration programme on green gram variety Meha (IPM 99 – 125) in Bharuch district of Gujarat. *J. Agric. & Vet. Sci*, **8** (9): 01-03.
- Meena O P, Sharma K C, Meena R H and Mitharwal B S 2012. Technology transfer through FLDs on mung bean in semi-arid region of Rajasthan. *Rajasthan J Ext Edu* **20**:182-186.

- Raj, A.D., Yadav, V. and Rathod, J.H. (2013). Impact of front line demonstrations on the yield of pulses. *Interernat. J. Sci. & Res. Public.*, **3** (9): 1-4.
- Singh, D., Singh, K.B., Gill, N.S. and Grewal, I.S. 2017. Impact analysis of front line demonstration on pulses in Punjab. *Internat. J. Farm Sci.* , **7** (1):190-194.
- Singh, G., Sharma, K., Dhaliwal, N.S. and Singh, J. 2012. Boosting moong productivity through front line demonstration. *Rajasthan J. Extn. Edu.*, **20**: 32-34.

**Table 1 : Difference between technological intervention & farmers practices for Green gram**

<b>Sr. No.</b>	<b>Particular</b>	<b>Front-line demonstration practices</b>	<b>Farmers practices</b>
1	Soil testing	Soil tested	Not adopted
2	Variety	BM-2003-2	Local variety
3	Seed rate	10 kg	12 kg

4	Seed treatment	seed treatment with Rhizobium culture @ 250 ml/10 kg seed + PSB @ 250 ml/10 kg seed and Tricoderma 1.25kg /10 kg seed	No seed treatment
5	Time of Sowing	Last week of June to First fortnight of July	As per monsoon
6	Fertilizer dose	Urea 62.50 kg, SSP 250 kg and Zink sulfate 10 kg (based on soil testing report) per ha in demo plot	Irrational use of nitrogenous fertilizers and graded fertilizer
7	Method of fertilizer application	Fertilizer drilled at the time of sowing	Broadcasting
8	Management of insect-pests	Insecticide spraying based on need at the economic threshold level (ETL)	Excess doses/ unrecommended brand of insecticide

**Table 2 : Grain yield and gap analysis of front line demonstration on Green gram**

Year	Area (ha)	No. of farmers	Yield q/ha			% increase over farmer practices	Technology gap (q/ha)	Extension Gap (q/ha)	Technology Index (%)
			Potential	FLD plots	Farmer practices				
2016-2017	10	25	10	8.80	6.36	27.20	1.20	2.44	12.00

2017-2018	10	25	10	4.31	3.49	23.47	5.69	0.82	56.90
2018-2019	10	25	10	5.35	4.14	29.07	4.65	1.21	46.50

**Mean      6.15          4.66          26.58          3.85          1.49          38.47**

**Table : 3 Economics analysis of demonstrated plots and farmers practices of Green gram**

Year	Av. Cost of Inputs (Rs/ha)		Av. Gross return (Rs/ha)		Average net return (Rs/ha)		B;C ratio	
	Demo. Plots	farmers practices	Demo. Plots	farmers practices	Demo. Plots	farmers practices	Demo. Plots	farmers practices
2016-2017	20742	18885	46566	36609	25823	17725	2.24	1.94
2017-2018	13466	12869	27512	22326	14041	9457	2.05	1.73
2018-2019	19051	17122	38324	29682	19272	12569	2.01	1.73

**Mean    17753    16292    37467    29539    19712    13250    2.10    1.80**