

A Study on the Sum of the Squares of Generalized p-Oresme Numbers: The Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$

Abstract. In this paper, closed forms of the sum formulas $\sum_{k=0}^n x^k W_{mk+j}^2$ for generalized Oresme numbers are presented. As special cases, we give sum formulas of Modified p-Oresme, p-Oresme-Lucas and p-Oresme numbers. We present the proofs to indicate how these formulas, in general, were discovered. Of course, all the listed formulas may be proved by induction, but that method of proof gives no clue about their discovery.

2020 Mathematics Subject Classification. 11B37, 11B39, 11B83.

Keywords. Oresme numbers, Modified p-Oresme numbers, p-Oresme-Lucas numbers, p-Oresme numbers, sum formulas.

1. Introduction

p-Oresme numbers (see, for example, [3]) are defined by the recurrence relation

$$O_{n+2} = O_{n+1} - \frac{1}{p^2} O_n, \quad O_0 = 0, O_1 = \frac{1}{p}, \quad p \neq 0.$$

The case $p = 2$, which is called the Oresme sequence, $\{O_n\}_{n \geq 0}$, was introduced by Nicole Oresme (1320–1382) in the 14-th century. Oresme obtained the sum of the rational numbers formed by the terms $0, \frac{1}{2}, \frac{2}{4}, \frac{3}{8}, \frac{4}{16}, \frac{5}{32}, \frac{6}{64}, \dots, \frac{n}{2^n}$. These numbers form a second order sequence and are defined by the recurrence relation

$$O_{n+2} = O_{n+1} - \frac{1}{4} O_n, \quad O_0 = 0, O_1 = \frac{1}{2}.$$

In [4], Horadam presented a history and obtained an abundance of properties of these numbers. Oresme numbers have many interesting properties and applications in many fields of science (see, for example, [1,2,3,4,5,6]).

For $0 \neq p \in \mathbb{R}$, a generalized p-Oresme sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1)\}_{n \geq 0}$ is defined by the second-order recurrence relations

$$W_n = W_{n-1} - \frac{1}{p^2} W_{n-2} \quad (1.1)$$

with the initial values $W_0 = c_0, W_1 = c_1$ not all being zero.

The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = p^2 W_{-(n-1)} - p^2 W_{-(n-2)}$$

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1.1) holds for all integer n .

Modified p-Oresme sequence $\{G_n\}_{n \geq 0}$, p-Oresme-Lucas sequence $\{H_n\}_{n \geq 0}$ and p-Oresme sequence $\{O_n\}_{n \geq 0}$ are defined, respectively, by the second-order recurrence relations

$$G_{n+2} = G_{n+1} - \frac{1}{p^2} G_n, \quad G_0 = 0, G_1 = 1, \quad (1.2)$$

$$H_{n+2} = H_{n+1} - \frac{1}{p^2} H_n, \quad H_0 = 2, H_1 = 1, \quad (1.3)$$

$$O_{n+2} = O_{n+1} - \frac{1}{p^2} O_n, \quad O_0 = 0, O_1 = \frac{1}{p}. \quad (1.4)$$

The sequences $\{G_n\}_{n \geq 0}$, $\{H_n\}_{n \geq 0}$ and $\{O_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$G_{-n} = p^2 G_{-(n-1)} - p^2 G_{-(n-2)},$$

$$H_{-n} = p^2 H_{-(n-1)} - p^2 H_{-(n-2)},$$

$$O_{-n} = p^2 O_{-(n-1)} - p^2 O_{-(n-2)},$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrences (1.2)-(1.4) hold for all integer n . For more information on generalized Oresme numbers, see Soykan [7].

In the next section, closed forms of the sum formulas $\sum_{k=0}^n x^k W_{mk+j}^2$ for generalized Oresme numbers are presented. Sum of the squares of linear recurrence relations have been studied by many authors and more detail can be found in the extensive literature, see for example, [9,10,11,12,13,14,15,16].

2. The Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$

The following theorem presents sum formulas of generalized p-Oresme numbers (the case $r = 1, s = -\frac{1}{p^2}$).

THEOREM 1. Suppose that $p^2 \neq 4$ so that $p^2 - 4 \neq 0$. Let x be a real (or complex) number. For all integers m and j , for generalized p-Oresme numbers (the case $r = 1, s = -\frac{1}{p^2}$) we have the following sum formulas:

(a): If $(1 + \frac{1}{p^{4m}} x^2 - x H_{2m})(\frac{1}{p^{2m}} x - 1) = (x^2 - p^{4m} H_{2m} x + p^{4m})(x - p^{2m}) p^{-6m} \neq 0$ then

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_1}{(p-2)(p+2)(x-p^{2m})(x^2-p^{4m}H_{2m}x+p^{4m})} \quad (2.1)$$

where

$$\begin{aligned} \Omega_1 = & (x - p^{4m} H_{2m})(x - p^{2m})(p - 2)(p + 2)x^{n+1}W_{mn+j}^2 + (x - p^{2m})(p - 2)(p + 2)x^{n+1}W_{mn-m+j}^2 + \\ & (x - p^{2m})(p - 2)p^{4m}(p + 2)W_j^2 + (p^{2m} - x)(p - 2)(p + 2)xW_{j-m}^2 + 2p^{-2mn+4m-2j+2}(H_{2m}p^{2m} - \\ & 2)(x^n - p^{2mn})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x \end{aligned}$$

(b): If $(x^2 - p^{4m}H_{2m}x + p^{4m})(x - p^{2m}) = (x - a)(x - b)(x - c) = 0$ for some

$$a = \frac{p^{4m}H_{2m} + \sqrt{p^{8m}H_{2m}^2 - 4p^{4m}}}{2}, b = \frac{p^{4m}H_{2m} - \sqrt{p^{8m}H_{2m}^2 - 4p^{4m}}}{2}, c = p^{2m} \in \mathbb{C}$$

and $a \neq b \neq c$, i.e., $x = a$ or $x = b$ or $x = c$, then,

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_2}{(p-2)(p+2)(3x^2 - 2p^{2m}(p^{2m}H_{2m} + 1)x + p^{4m} + p^{6m}H_{2m})}$$

where

$$\begin{aligned} \Omega_2 = & (p^2 - 4)((x - p^{4m}H_{2m})x^{n+1} + (x - p^{2m})((n+2)x - p^{4m}(n+1)H_{2m})x^n)W_{mn+j}^2 + (p^2 - \\ & 4)((n+2)x - p^{2m}(n+1))x^nW_{mn-m+j}^2 + p^{4m}(p^2 - 4)W_j^2 - (p^2 - 4)(2x - p^{2m})W_{j-m}^2 + 2p^{-2mn+4m-2j+2}(p^{2m}H_{2m} - \\ & 2)(x^n + nx^n - p^{2mn})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0) \end{aligned}$$

(c): If $p^{8m}H_{2m}^2 - 4p^{4m} = 0$ and $(x^2 - p^{4m}xH_{2m} + p^{4m})(x - p^{2m}) = (x - a)^2(x - c) = 0$ for some

$$a = \frac{p^{4m}H_{2m}}{2}, c = p^{2m} \in \mathbb{C} \text{ and } a \neq c \text{ then if } x = c \text{ then}$$

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_3}{(p-2)(p+2)(3x^2 - 2p^{2m}(p^{2m}H_{2m} + 1)x + p^{4m} + p^{6m}H_{2m})}$$

where

$$\begin{aligned} \Omega_3 = & (p^2 - 4)((x - p^{4m}H_{2m})x^{n+1} + (x - p^{2m})((n+2)x - p^{4m}(n+1)H_{2m})x^n)W_{mn+j}^2 + (p^2 - \\ & 4)((n+2)x - p^{2m}(n+1))x^nW_{mn-m+j}^2 + p^{4m}(p^2 - 4)W_j^2 - (p^2 - 4)(2x - p^{2m})W_{j-m}^2 + 2p^{-2mn+4m-2j+2}(p^{2m}H_{2m} - \\ & 2)(x^n + nx^n - p^{2mn})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0) \end{aligned}$$

and if $x = a$ then

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_4}{2(p-2)(p+2)(3x - p^{2m} - p^{4m}H_{2m})}$$

where

$$\begin{aligned} \Omega_4 = & (p^2 - 4)((n+3)(n+2)x^2 - xp^{2m}(n+2)(n+1)(p^{2m}H_{2m} + 1) + p^{6m}n(n+1)H_{2m})x^{n-1}W_{mn+j}^2 + \\ & (p^2 - 4)(n+1)((2+n)x^n - p^{2m}nx^{n-1})W_{mn-m+j}^2 - 2(p^2 - 4)W_{j-m}^2 + 2n(n+1)p^{-2mn+4m-2j+2} \\ & (p^{2m}H_{2m} - 2)(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x^{n-1} \end{aligned}$$

Proof. Take $r = 1, s = -\frac{1}{p^2}$ and $H_n = H_n$ in Soykan [11, Theorem 2.1.]. Note that $(x^2 - p^{4m}xH_{2m} + p^{4m})(x - p^{2m}) = (x - a)^3 = 0$ can not be possible for some $a = p^{2m} \in \mathbb{C}$, $m \neq 0$, i.e., $x = a$, because

otherwise we must have $p^{8m}H_{2m}^2 - 4p^{4m} = 0$, $\frac{p^{4m}H_{2m}}{2} = p^{2m}$ and

$$\begin{aligned} \frac{p^{4m}H_{2m}}{2} - p^{2m} &= \frac{1}{2}p^{2m}(p^{2m}H_{2m} - 2) = 0 \Rightarrow p^{2m}H_{2m} - 2 = 0 \\ &\Rightarrow p^{2m} \times 2 \left(\frac{1}{2}\right)^{2m} - 2 = 0 \\ &\Rightarrow \left(\frac{p}{2}\right)^{2m} - 1 = 0 \end{aligned}$$

in which the last equality doesn't have any integer solutions m other than $m = 0$ (so $x = p^{2m} = 1$) for the case $p \neq 2$. Here we used

$$\begin{aligned} \alpha &= \frac{r}{2}, \\ H_{2m} &= 2\alpha^{2m} = 2 \left(\frac{1}{2}\right)^{2m}. \end{aligned}$$

In this case (i.e. $m = 0, x = 1$), we have $\sum_{k=0}^n W_j^2 = (n+1)W_j^2$. $\square$

Note that in the last theorem, the case $p^2 \neq 4$ so that $p^2 - 4 \neq 0$ is considered. The sum formulas for the case $p^2 = 4$ so that $p^2 - 4 = 0$ is given in Soykan [8, Theorem 2.1].

Note also that (2.1) can be written in the following form

$$\sum_{k=1}^n x^k W_{mk+j}^2 = \frac{\Omega_6}{(p-2)(p+2)(x-p^{2m})(x^2-p^{4m}H_{2m}x+p^{4m})}$$

where

$$\begin{aligned} \Omega_6 &= (p^2-4)(x-p^{2m})(x-p^{4m}H_{2m})x^{n+1}W_{mn+j}^2 + (p^2-4)(x-p^{2m})x^{n+1}W_{mn-m+j}^2 - (p^2-4)(x-p^{2m})(x- \\ &p^{4m}H_{2m})xW_j^2 - (p^2-4)(x-p^{2m})W_{j-m}^2x + 2p^{-2mn+4m-2j+2}(x^n-p^{2mn})(p^{2m}H_{2m}-2)(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x. \end{aligned}$$

As special cases of m and j in the last Theorem, we obtain the following proposition.

PROPOSITION 2. *Suppose that $p^2 \neq 4$ so that $p^2 - 4 \neq 0$. For generalized p -Oresme numbers (the case $r = 1, s = -\frac{1}{p^2}$) we have the following sum formulas for $n \geq 0$:*

(a): ($m = 1, j = 0$)

If $(x-p^2)(x^2-p^2(p^2-2)x+p^4) \neq 0$, i.e.,

$$x \neq \frac{p^2(p^2-2) + \sqrt{p^6(p^2-4)}}{2}, x \neq \frac{p^2(p^2-2) - \sqrt{p^6(p^2-4)}}{2}, x \neq p^2,$$

then

$$\sum_{k=0}^n x^k W_k^2 = \frac{\Phi}{(p^2-4)(x-p^2)(x^2-p^2(p^2-2)x+p^4)}$$

where

$$\begin{aligned} \Phi &= (x-p^2(p^2-2))(x-p^2)(p^2-4)x^{n+1}W_n^2 + (x-p^2)(p^2-4)x^{n+1}W_{n-1}^2 + (x-p^2)(p^2- \\ &4)p^4W_0^2 + p^4(p^2-x)(p^2-4)(W_0-W_1)^2x + 2p^{6-2n}((p^2-2)-2)(x^n-p^{2n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x, \end{aligned}$$

and

if $(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4) = 0$, i.e.,

$$x = \frac{p^2(p^2 - 2) + \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = \frac{p^2(p^2 - 2) - \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = p^2,$$

then

$$\sum_{k=0}^n x^k W_k^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^2(p^2 - 1)x + p^4(p^2 - 1))}$$

where

$$\begin{aligned} \Lambda = & (p^2 - 4)((x - p^2(p^2 - 2))x^{n+1} + (x - p^2)((n + 2)x - (n + 1)p^2(p^2 - 2))x^n)W_n^2 + (p^2 - 4)((n + 2)x - p^2(n + 1))x^n W_{n-1}^2 \\ & + p^4(p^2 - 4)W_0^2 - p^4(2x - p^2)(p^2 - 4)(W_0 - W_1)^2 + 2p^{6-2n}((p^2 - 2) - 2)(x^n + nx^n - p^{2n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0). \end{aligned}$$

(b): $(m = 2, j = 0)$

If $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) \neq 0$, i.e.,

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k W_{2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)}$$

where

$$\begin{aligned} \Phi = & (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}W_{2n}^2 + (x - p^4)(p^2 - 4)x^{n+1}W_{2n-2}^2 + (x - p^4)(p^2 - 4)p^8W_0^2 \\ & + (p^4 - x)(p^2 - 4)p^4(p^2W_0 - W_0 - p^2W_1)^2x + 2p^{12-4n}(p^2 - 4)(x^n - p^{4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x, \end{aligned}$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = 0$ and $p^2 \neq 2$ i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k W_{2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))}$$

where

$$\begin{aligned} \Lambda = & (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n + 2)x - p^4(p^4 - 4p^2 + 2)(n + 1))x^n)W_{2n}^2 + \\ & (p^2 - 4)((n + 2)x - p^4(n + 1))x^n W_{2n-2}^2 + p^8(p^2 - 4)W_0^2 - (p^2 - 4)(2x - p^4)p^4(p^2W_0 - W_0 - p^2W_1)^2 + \\ & 2p^{12-4n}(p^2 - 4)(x^n + nx^n - p^{4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0), \end{aligned}$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = (x - 4)(x + 4)^2 = 0$, $x \neq p^4 = 4$ and $p^2 = 2$

i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k W_{2k}^2 = \frac{\Theta}{32}$$

where

$$\Theta = 64(-4)^{n-1} (n^2 - 2) W_{2n}^2 + (-4)^{n+1} (n+1)^2 W_{2n-2}^2 + 16(W_0 - 2W_1)^2 - n(n+1) (-4)^{n-1} 2^{8-2n} (W_1^2 + \frac{1}{2}W_0^2 - W_1W_0).$$

(c): ($m = 2, j = 1$)

$$\text{If } (x - p^4) (x^2 - p^4 (p^4 - 4p^2 + 2) x + p^8) \neq 0, \text{ i.e.,}$$

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k W_{2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)},$$

where

$$\Phi = (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}W_{2n+1}^2 + (x - p^4)(p^2 - 4)x^{n+1}W_{2n-1}^2 + (x - p^4)(p^2 - 4)p^8W_1^2 + (p^4 - x)(p^2 - 4)p^4(W_0 - W_1)^2x + 2p^{10-4n}(p^2 - 4)(x^n - p^{4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x,$$

$$\text{and if } (x - p^4) (x^2 - p^4 (p^4 - 4p^2 + 2) x + p^8) = 0 \text{ and } p^2 \neq 2 \text{ i.e.,}$$

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k W_{2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))},$$

where

$$\Lambda = (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n+2)x - p^4(p^4 - 4p^2 + 2)(n+1))x^n)W_{2n+1}^2 + (p^2 - 4)((n+2)x - p^4(n+1))x^nW_{2n-1}^2 + p^8(p^2 - 4)W_1^2 - (p^2 - 4)(2x - p^4)p^4(W_0 - W_1)^2 + 2p^{10-4n}(p^2 - 4)(x^n + nx^n - p^{4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0),$$

$$\text{and if } (x - p^4) (x^2 - p^4 (p^4 - 4p^2 + 2) x + p^8) = (x - 4)(x + 4)^2 = 0, x \neq p^4 = 4 \text{ and } p^2 = 2$$

i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k W_{2k+1}^2 = \frac{\Theta}{32},$$

where

$$\Theta = 64(-4)^{n-1} (n^2 - 2) W_{2n+1}^2 + (-4)^{n+1} (n+1)^2 W_{2n-1}^2 + 16(W_0 - W_1)^2 - n(n+1) (-4)^{n-1} 2^{7-2n} (W_1^2 + \frac{1}{2}W_0^2 - W_1W_0).$$

(d): ($m = -1, j = 0$)

$$\text{If } (p^2x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) \neq 0, \text{ i.e.,}$$

$$x \neq \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq p^{-2},$$

then

$$\sum_{k=0}^n x^k W_{-k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-2})(x^2 - (p^2 - 2)p^{-2}x + p^{-4})},$$

where

$$\Phi = (x - p^{2m})(p^2 - 4)x^{n+1}W_{-n+1}^2 + (x - (p^2 - 2)p^{-2})(x - p^{-2})(p^2 - 4)x^{n+1}W_{-n}^2 + (x - p^{-2})(p^2 - 4)p^{-4}W_0^2 + (p^{-2} - x)(p^2 - 4)xW_1^2 + 2p^{2n-2}(p^2 - 4)(x^n - p^{-2n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x$$

and if $(p^2x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) = 0$, i.e.,

$$x = \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = p^{-2},$$

then

$$\sum_{k=0}^n x^k W_{-k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)p^{-2}x + (p^2 - 1)p^{-4})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-2}(n + 1))x^n W_{-n+1}^2 + (p^2 - 4)(p^4(n + 3)x^2 - p^2x(p^2 - 1)(n + 2) + (p^2 - 2)(n + 1))p^{-4}x^n W_{-n}^2 + p^{-4}(p^2 - 4)W_0^2 - (p^2 - 4)(2x - p^{-2})W_1^2 + 2p^{2n-2}(p^2 - 4)(x^n + nx^n - p^{-2n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0).$$

(e): $(m = -2, j = 0)$

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, i.e.,

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k W_{-2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1}W_{-2n+2}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1}W_{-2n}^2 + (x - p^{-4})(p^2 - 4)p^{-8}W_0^2 + (p^{-4} - x)(p^2 - 4)x(-W_0 + p^2W_1)^2p^{-4} + 2p^{4n-4}(p^2 - 4)(x^n - p^{-4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, and $p^2 \neq 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(and $p^2 \neq 2$, i.e., $p \neq -\sqrt{2}, p \neq \sqrt{2}$) then

$$\sum_{k=0}^n x^k W_{-2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-4}(n + 1))x^n W_{-2n+2}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + (x - p^{-4})((n + 2)x - (p^4 - 4p^2 + 2)p^{-4}(n + 1))x^n)W_{-2n}^2 + p^{-8}(p^2 - 4)W_0^2 - (p^2 - 4)(2x - p^{-4})(-W_0 + p^2W_1)^2p^{-4} + 2p^{4n-4}(p^2 - 4)(x^n + nx^n - p^{-4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0),$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0, x \neq p^{-4} = \frac{1}{4}$ and $p^2 = 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k W_{-2k}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4 \left(-\frac{1}{4}\right)^n (n+1)^2 W_{-2n+2}^2 + \frac{1}{4} \left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) W_{-2n}^2 + (W_0 - 2W_1)^2 - n(n+1)2^{2n-1} \left(-\frac{1}{4}\right)^{n-1} (2W_1^2 + W_0^2 - 2W_1W_0).$$

(f): ($m = -2, j = 1$)

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, i.e.,

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k W_{-2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1}W_{-2n+3}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1}W_{-2n+1}^2 + (x - p^{-4})(p^2 - 4)p^{-8}W_1^2 + (p^{-4} - x)(p^2 - 4)p^{-4}(-W_0 - W_1 + p^2W_1)^2x + 2p^{4n-6}(p^2 - 4)(x^n - p^{-4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0)x,$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, and $p^2 \neq 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(and $p^2 \neq 2$, i.e., $p \neq -\sqrt{2}, p \neq \sqrt{2}$) then

$$\sum_{k=0}^n x^k W_{-2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$$\Lambda = (p^2 - 4)((n+2)x - p^{-4}(n+1))x^n W_{-2n+3}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + (x - p^{-4})((n+2)x - (p^4 - 4p^2 + 2)p^{-4}(n+1))x^n)W_{-2n+1}^2 + p^{-8}(p^2 - 4)W_1^2 - (p^2 - 4)(2x - p^{-4})p^{-4}(-W_0 - W_1 + p^2W_1)^2 + 2p^{4n-6}(p^2 - 4)(x^n + nx^n - p^{-4n})(W_1^2 + \frac{1}{p^2}W_0^2 - W_1W_0),$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0, x \neq p^{-4} = \frac{1}{4}$ and $p^2 = 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k W_{-2k+1}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4 \left(-\frac{1}{4}\right)^n (n+1)^2 W_{-2n+3}^2 + \frac{1}{4} \left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) W_{-2n+1}^2 + (W_0 - W_1)^2 - n(n+1)2^{2n-2}(2W_1^2 + W_0^2 - 2W_1W_0) \left(-\frac{1}{4}\right)^{n-1}.$$

Note that in the last proposition, the case $p^2 \neq 4$ so that $p^2 - 4 \neq 0$ is considered. The sum formulas for the case $p^2 = 4$ so that $p^2 - 4 = 0$ is given in Soykan [8, Proposition 2.1].

From the above proposition, we have the following corollary which gives sum formulas of modified p-Oresme numbers (take $W_n = G_n$ with $G_0 = 0, G_1 = 1$).

COROLLARY 3. *Suppose that $p^2 \neq 4$ so that $p^2 - 4 \neq 0$. For $n \geq 0$, modified p-Oresme numbers have the following properties:*

(a): ($m = 1, j = 0$)

If $(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4) \neq 0$, i.e.,

$$x \neq \frac{p^2(p^2 - 2) + \sqrt{p^6(p^2 - 4)}}{2}, x \neq \frac{p^2(p^2 - 2) - \sqrt{p^6(p^2 - 4)}}{2}, x \neq p^2,$$

then

$$\sum_{k=0}^n x^k G_k^2 = \frac{\Phi}{(p^2 - 4)(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4)}$$

where

$$\Phi = (x - p^2(p^2 - 2))(x - p^2)(p^2 - 4)x^{n+1}G_n^2 + (x - p^2)(p^2 - 4)x^{n+1}G_{n-1}^2 - p^{-2n+4}(p^2 - 4)(p^{2n}(x + p^2) - 2p^2x^n)x,$$

and

if $(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4) = 0$, i.e.,

$$x = \frac{p^2(p^2 - 2) + \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = \frac{p^2(p^2 - 2) - \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = p^2,$$

then

$$\sum_{k=0}^n x^k G_k^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^2(p^2 - 1)x + p^4(p^2 - 1))}$$

where

$$\Lambda = (p^2 - 4)((x - p^2(p^2 - 2))x^{n+1} + (x - p^2)((n+2)x - (n+1)p^2(p^2 - 2))x^n)G_n^2 + (p^2 - 4)((n+2)x - p^2(n+1))x^nG_{n-1}^2 + p^{-2n+4}(p^2 - 4)(-p^{2n}(2x + p^2) + 2p^2(n+1)x^n).$$

(b): ($m = 2, j = 0$)

If $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) \neq 0$, i.e.,

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k G_{2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)}$$

where

$$\Phi = (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}G_{2n}^2 + (x - p^4)(p^2 - 4)x^{n+1}G_{2n-2}^2 - p^{-4n+8}(p^2 - 4)(p^{4n}(x + p^4) - 2p^4x^n)x,$$

$$\text{and if } (x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = 0 \text{ and } p^2 \neq 2 \text{ i.e.,}$$

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k G_{2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))}$$

where

$$\Lambda = (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n + 2)x - p^4(p^4 - 4p^2 + 2)(n + 1))x^n)G_{2n}^2 + (p^2 - 4)((n + 2)x - p^4(n + 1))x^n G_{2n-2}^2 + p^{-4n+8}(p^2 - 4)(-p^{4n}(2x + p^4) + 2p^4(n + 1)x^n),$$

$$\text{and if } (x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = (x - 4)(x + 4)^2 = 0, x \neq p^4 = 4 \text{ and } p^2 = 2$$

i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k G_{2k}^2 = \frac{\Theta}{32}$$

where

$$\Theta = 64(-4)^{n-1}(n^2 - 2)G_{2n}^2 + (-4)^{n+1}(n + 1)^2 G_{2n-2}^2 + 16(G_0 - 2G_1)^2 - n(n + 1)(-4)^{n-1}2^{8-2n}$$

(c): ($m = 2, j = 1$)

$$\text{If } (x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) \neq 0, \text{ i.e.,}$$

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k G_{2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)},$$

where

$$\Phi = (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}G_{2n+1}^2 + (x - p^4)(p^2 - 4)x^{n+1}G_{2n-1}^2 - p^{-4n+4}(p^2 - 4)(p^{4n}(x^2 - 2p^4x + 2p^6x + p^8) - 2p^6x^{n+1}),$$

$$\text{and if } (x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = 0 \text{ and } p^2 \neq 2 \text{ i.e.,}$$

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k G_{2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))},$$

where

$$\Lambda = (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n + 2)x - p^4(p^4 - 4p^2 + 2)(n + 1))x^n)G_{2n+1}^2 +$$

$$(p^2 - 4)((n + 2)x - p^4(n + 1))x^n G_{2n-1}^2 + 2p^{-4n+4}(p^2 - 4)(p^{4n}(-x - p^6 + p^4) + p^6(n + 1)x^n),$$

$$\text{and if } (x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = (x - 4)(x + 4)^2 = 0, x \neq p^4 = 4 \text{ and } p^2 = 2$$

i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k G_{2k+1}^2 = \frac{\Theta}{32},$$

where

$$\Theta = 64(-4)^{n-1}(n^2 - 2)G_{2n+1}^2 + (-4)^{n+1}(n + 1)^2 G_{2n-1}^2 + 16(G_0 - G_1)^2 - n(n + 1)(-4)^{n-1} 2^{7-2n}$$

(d): ($m = -1, j = 0$)

If $(p^2 x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) \neq 0$, i.e.,

$$x \neq \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq p^{-2},$$

then

$$\sum_{k=0}^n x^k G_{-k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-2})(x^2 - (p^2 - 2)p^{-2}x + p^{-4})},$$

where

$$\Phi = (x - p^{-2})(p^2 - 4)x^{n+1}G_{-n+1}^2 + (x - (p^2 - 2)p^{-2})(x - p^{-2})(p^2 - 4)x^{n+1}G_{-n}^2 + (p^2 - 4)p^{-2}(2p^{2n}x^n - p^2x - 1)x$$

and if $(p^2 x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) = 0$, i.e.,

$$x = \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = p^{-2},$$

then

$$\sum_{k=0}^n x^k G_{-k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)p^{-2}x + (p^2 - 1)p^{-4})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-2}(n + 1))x^n G_{-n+1}^2 + (p^2 - 4)(p^4(n + 3)x^2 - p^2x(p^2 - 1)(n + 2) +$$

$$(p^2 - 2)(n + 1))p^{-4}x^n G_{-n}^2 + (p^2 - 4)p^{-2}(2p^{2n}(n + 1)x^n - 2p^2x - 1)$$

(e): ($m = -2, j = 0$)

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, i.e.,

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k G_{-2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1}G_{-2n+2}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1}G_{-2n}^2 + (p^2 - 4)p^{-4}(2p^{4n}x^n - p^4x - 1)x$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, and $p^2 \neq 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(and $p^2 \neq 2$, i.e., $p \neq -\sqrt{2}, p \neq \sqrt{2}$) then

$$\sum_{k=0}^n x^k G_{-2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-4}(n + 1))x^n G_{-2n+2}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + (x - p^{-4})((n + 2)x - (p^4 - 4p^2 + 2)p^{-4}(n + 1))x^n)G_{-2n}^2 + (p^2 - 4)p^{-4}(2p^{4n}(n + 1)x^n - 2p^4x - 1)$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0$, $x \neq p^{-4} = \frac{1}{4}$ and $p^2 = 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k G_{-2k}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4\left(-\frac{1}{4}\right)^n (n + 1)^2 G_{-2n+2}^2 + \frac{1}{4}\left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) G_{-2n}^2 + 4(n(n + 1)2^{2n} \left(-\frac{1}{4}\right)^n + 1)$$

(f): ($m = -2, j = 1$)

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, i.e.,

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k G_{-2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1}G_{-2n+3}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1}G_{-2n+1}^2 + (p^2 - 4)p^{-12}(-p^8(p - 1)^2(p + 1)^2x^2 + p^4(p^4 - 4p^2 + 2)x + 2p^{4n+6}x^{n+1} - 1)$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, and $p^2 \neq 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(and $p^2 \neq 2$, i.e., $p \neq -\sqrt{2}, p \neq \sqrt{2}$) then

$$\sum_{k=0}^n x^k G_{-2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$$\begin{aligned} \Lambda = & (p^2 - 4)((n + 2)x - p^{-4}(n + 1))x^n G_{-2n+3}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + \\ & (x - p^{-4})((n + 2)x - (p^4 - 4p^2 + 2)p^{-4}(n + 1))x^n)G_{-2n+1}^2 + (p^2 - 4)p^{-8}(-2p^4x(p - 1)^2(p + 1)^2 + \\ & 2p^{4n+2}(n + 1)x^n + p^4 - 4p^2 + 2) \end{aligned}$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0$, $x \neq p^{-4} = \frac{1}{4}$ and $p^2 = 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k G_{-2k+1}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4\left(-\frac{1}{4}\right)^n (n + 1)^2 G_{-2n+3}^2 + \frac{1}{4}\left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) G_{-2n+1}^2 + 2n(n + 1)2^{2n}\left(-\frac{1}{4}\right)^n + 1.$$

Note that in the last corollary, the case $p^2 \neq 4$ so that $p^2 - 4 \neq 0$ is considered. The sum formulas for the case $p^2 = 4$ so that $p^2 - 4 = 0$ is given in Soykan [8, Corollary 2.2].

Taking $W_n = H_n$ with $H_0 = 2, H_1 = 1$ in the last proposition, we have the following corollary which presents sum formulas of p-Oresme-Lucas numbers.

COROLLARY 4. Suppose that $p^2 \neq 4$ so that $p^2 - 4 \neq 0$. For $n \geq 0$, p-Oresme-Lucas numbers have the following properties:

(a): ($m = 1, j = 0$)

If $(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4) \neq 0$, i.e.,

$$x \neq \frac{p^2(p^2 - 2) + \sqrt{p^6(p^2 - 4)}}{2}, x \neq \frac{p^2(p^2 - 2) - \sqrt{p^6(p^2 - 4)}}{2}, x \neq p^2,$$

then

$$\sum_{k=0}^n x^k H_k^2 = \frac{\Phi}{(p^2 - 4)(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4)}$$

where

$$\begin{aligned} \Phi = & (x - p^2(p^2 - 2))(x - p^2)(p^2 - 4)x^{n+1}H_n^2 + (x - p^2)(p^2 - 4)x^{n+1}H_{n-1}^2 + p^{-2n+4}(p^2 - \\ & 4)(p^{2n}(p^2(3x - 4) - x(x + 4)) - 2(p^2 - 4)x^{n+1}), \end{aligned}$$

and

if $(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4) = 0$, i.e.,

$$x = \frac{p^2(p^2 - 2) + \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = \frac{p^2(p^2 - 2) - \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = p^2,$$

then

$$\sum_{k=0}^n x^k H_k^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^2(p^2 - 1)x + p^4(p^2 - 1))}$$

where

$$\Lambda = (p^2 - 4)((x - p^2(p^2 - 2))x^{n+1} + (x - p^2)((n + 2)x - (n + 1)p^2(p^2 - 2))x^n)H_n^2 + (p^2 - 4)((n + 2)x - p^2(n + 1))x^n H_{n-1}^2 + p^{-2n+4}(p^2 - 4)(p^{2n}(-2x + 3p^2 - 4) - 2(n + 1)(p^2 - 4)x^n).$$

(b): ($m = 2, j = 0$)

If $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) \neq 0$, i.e.,

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k H_{2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)}$$

where

$$\Phi = (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}H_{2n}^2 + (x - p^4)(p^2 - 4)x^{n+1}H_{2n-2}^2 + p^{-4n+4}(p^2 - 4)(p^{4n}(4p^2x^2 - p^4x^2 - 4x^2 + 8p^4x - 12p^6x + 3p^8x - 4p^8) - 2p^6(p^2 - 4)x^{n+1})$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = 0$ and $p^2 \neq 2$ i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k H_{2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))}$$

where

$$\Lambda = (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n + 2)x - p^4(p^4 - 4p^2 + 2)(n + 1))x^n)H_{2n}^2 + (p^2 - 4)((n + 2)x - p^4(n + 1))x^n H_{2n-2}^2 + p^{-4n+4}(p^2 - 4)(p^{4n}(-8x + 8p^2x - 2p^4x + 8p^4 - 12p^6 + 3p^8) - 2p^6(p^2 - 4)(n + 1)x^n)$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = (x - 4)(x + 4)^2 = 0$, $x \neq p^4 = 4$ and $p^2 = 2$

i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k H_{2k}^2 = \frac{\Theta}{32}$$

where

$$\Theta = 64(-4)^{n-1}(n^2 - 2)H_{2n}^2 + (-4)^{n+1}(n + 1)^2H_{2n-2}^2 - n(n + 1)2^{-2n+8}(-4)^{n-1}$$

(c): ($m = 2, j = 1$)

If $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) \neq 0$, i.e.,

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k H_{2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)},$$

where

$$\Phi = (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}H_{2n+1}^2 + (x - p^4)(p^2 - 4)x^{n+1}H_{2n-1}^2 - p^{-4n+4}(p^2 - 4)(p^{4n}(x^2 + 6p^4x - 2p^6x + p^8) + 2p^4(p^2 - 4)x^{n+1})$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = 0$ and $p^2 \neq 2$ i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k H_{2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))},$$

where

$$\Lambda = (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n + 2)x - p^4(p^4 - 4p^2 + 2)(n + 1))x^n)H_{2n+1}^2 + (p^2 - 4)((n + 2)x - p^4(n + 1))x^n H_{2n-1}^2 - 2p^{-4n+4}(p^2 - 4)(p^{4n}(x + 3p^4 - p^6) + p^4(p^2 - 4)(n + 1)x^n)$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = (x - 4)(x + 4)^2 = 0$, $x \neq p^4 = 4$ and $p^2 = 2$

i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k H_{2k+1}^2 = \frac{\Theta}{32},$$

where

$$\Theta = 64(-4)^{n-1}(n^2 - 2)H_{2n+1}^2 + (-4)^{n+1}(n + 1)^2 H_{2n-1}^2 + 2^{-2n+4}(2^{2n} + 2n(n + 1)(-4)^n).$$

(d): ($m = -1, j = 0$)

If $(p^2x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) \neq 0$, i.e.,

$$x \neq \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq p^{-2},$$

then

$$\sum_{k=0}^n x^k H_{-k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-2})(x^2 - (p^2 - 2)p^{-2}x + p^{-4})},$$

where

$$\Phi = (x - p^{-2})(p^2 - 4)x^{n+1}H_{-n+1}^2 + (x - (p^2 - 2)p^{-2})(x - p^{-2})(p^2 - 4)x^{n+1}H_{-n}^2 - (p^2 - 4)p^{-6}(p^6x^2 + 4p^2x - 3p^4x + 2p^{2n+2}(p^2 - 4)x^{n+1} + 4)$$

and if $(p^2x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) = 0$, i.e.,

$$x = \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = p^{-2},$$

then

$$\sum_{k=0}^n x^k H_{-k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)p^{-2}x + (p^2 - 1)p^{-4})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-2}(n + 1))x^n H_{-n+1}^2 + (p^2 - 4)(p^4(n + 3)x^2 - p^2x(p^2 - 1)(n + 2) + (p^2 - 2)(n + 1)p^{-4}x^n H_{-n}^2 + (p^2 - 4)p^{-4}(-2p^{2n}(p^2 - 4)(n + 1)x^n - 2p^4x + 3p^2 - 4)$$

(e): $(m = -2, j = 0)$

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, i.e.,

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k H_{-2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1}H_{-2n+2}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1}H_{-2n}^2 + (p^2 - 4)p^{-12}(-p^8(p^2 - 2)^2x^2 + p^4(3p^4 - 12p^2 + 8)x - 2p^{4n+6}(p^2 - 4)x^{n+1} - 4)$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, and $p^2 \neq 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(and $p^2 \neq 2$, i.e., $p \neq -\sqrt{2}, p \neq \sqrt{2}$) then

$$\sum_{k=0}^n x^k H_{-2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-4}(n + 1))x^n H_{-2n+2}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + (x - p^{-4})((n + 2)x - (p^4 - 4p^2 + 2)p^{-4}(n + 1))x^n)H_{-2n}^2 + (p^2 - 4)p^{-8}(-2p^2x^n p^{4n}(p^2 - 4)(n + 1) - 2p^4(p^2 - 2)^2x + 3p^4 - 12p^2 + 8)$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0, x \neq p^{-4} = \frac{1}{4}$ and $p^2 = 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k H_{-2k}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4\left(-\frac{1}{4}\right)^n (n + 1)^2 H_{-2n+2}^2 + \frac{1}{4}\left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) H_{-2n}^2 - n(n + 1) 2^{2n} \left(-\frac{1}{4}\right)^{n-1}$$

(f): ($m = -2, j = 1$)

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, i.e.,

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k H_{-2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1}H_{-2n+3}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1}H_{-2n+1}^2 + (p^2 - 4)p^{-12}(-p^8(p^2 - 3)^2x^2 + p^4(p^4 - 4p^2 + 2)x - 2p^{4n+4}(p^2 - 4)x^{n+1} - 1)$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, and $p^2 \neq 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(and $p^2 \neq 2$, i.e., $p \neq -\sqrt{2}, p \neq \sqrt{2}$) then

$$\sum_{k=0}^n x^k H_{-2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-4}(n + 1))x^n H_{-2n+3}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + (x - p^{-4})((n + 2)x - (p^4 - 4p^2 + 2)p^{-4}(n + 1))x^n)H_{-2n+1}^2 + (p^2 - 4)p^{-8}(-2p^{4n}(p^2 - 4)(n + 1)x^n - 2p^4(p^2 - 3)^2x + p^4 - 4p^2 + 2)$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0$, $x \neq p^{-4} = \frac{1}{4}$ and $p^2 = 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k H_{-2k+1}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4\left(-\frac{1}{4}\right)^n (n + 1)^2 H_{-2n+3}^2 + \frac{1}{4}\left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) H_{-2n+1}^2 + n(n + 1)2^{2n+1}\left(-\frac{1}{4}\right)^n + 1.$$

Note that in the last corollary, the case $p^2 \neq 4$ so that $p^2 - 4 \neq 0$ is considered. The sum formulas for the case $p^2 = 4$ so that $p^2 - 4 = 0$ is given in Soykan [8, Corollary 2.3].

From the above proposition, we have the following corollary which gives sum formulas of p-Oresme numbers (take $W_n = O_n$ with $O_0 = 0, O_1 = \frac{1}{p}$).

COROLLARY 5. Suppose that $p^2 \neq 4$ so that $p^2 - 4 \neq 0$. For $n \geq 0$, p-Oresme numbers have the following properties:

(a): ($m = 1, j = 0$)

If $(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4) \neq 0$, i.e.,

$$x \neq \frac{p^2(p^2 - 2) + \sqrt{p^6(p^2 - 4)}}{2}, x \neq \frac{p^2(p^2 - 2) - \sqrt{p^6(p^2 - 4)}}{2}, x \neq p^2,$$

then

$$\sum_{k=0}^n x^k O_k^2 = \frac{\Phi}{(p^2 - 4)(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4)}$$

where

$$\Phi = (x - p^2(p^2 - 2))(x - p^2)(p^2 - 4)x^{n+1}O_n^2 + (x - p^2)(p^2 - 4)x^{n+1}O_{n-1}^2 - p^{-2n+2}(p^2 - 4)(p^{2n}(x + p^2) - 2p^2x^n)x$$

and

if $(x - p^2)(x^2 - p^2(p^2 - 2)x + p^4) = 0$, i.e.,

$$x = \frac{p^2(p^2 - 2) + \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = \frac{p^2(p^2 - 2) - \sqrt{p^6(p^2 - 4)}}{2} \text{ or } x = p^2,$$

then

$$\sum_{k=0}^n x^k O_k^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^2(p^2 - 1)x + p^4(p^2 - 1))}$$

where

$$\Lambda = (p^2 - 4)((x - p^2(p^2 - 2))x^{n+1} + (x - p^2)((n + 2)x - (n + 1)p^2(p^2 - 2))x^n)O_n^2 + (p^2 - 4)((n + 2)x - p^2(n + 1))x^n O_{n-1}^2 + p^{-2n+2}(p^2 - 4)(-p^{2n}(2x + p^2) + 2p^2(n + 1)x^n)$$

(b): ($m = 2, j = 0$)

If $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) \neq 0$, i.e.,

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k O_{2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)}$$

where

$$\Phi = (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}O_{2n}^2 + (x - p^4)(p^2 - 4)x^{n+1}O_{2n-2}^2 - p^{-4n+6}(p^2 - 4)(p^{4n}(x + p^4) - 2p^4x^n)x$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = 0$ and $p^2 \neq 2$ i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k O_{2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))}$$

where

$$\Lambda = (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n + 2)x - p^4(p^4 - 4p^2 + 2)(n + 1))x^n)O_{2n}^2 + (p^2 - 4)((n + 2)x - p^4(n + 1))x^n O_{2n-2}^2 + p^{-4n+6}(p^2 - 4)(-p^{4n}(2x + p^4) + 2p^4(n + 1)x^n)$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = (x - 4)(x + 4)^2 = 0$, $x \neq p^4 = 4$ and $p^2 = 2$ i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k O_{2k}^2 = \frac{\Theta}{32}$$

where

$$\Theta = 64(-4)^{n-1}(n^2 - 2)O_{2n}^2 + (-4)^{n+1}(n + 1)^2 O_{2n-2}^2 + p^{-2}2^{-2n+6}(2^{2n} + n(n + 1)(-4)^n)$$

(c): ($m = 2, j = 1$)

If $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) \neq 0$, i.e.,

$$x \neq \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2}, x \neq p^4,$$

then

$$\sum_{k=0}^n x^k O_{2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8)},$$

where

$$\Phi = (x - p^4(p^4 - 4p^2 + 2))(x - p^4)(p^2 - 4)x^{n+1}O_{2n+1}^2 + (x - p^4)(p^2 - 4)x^{n+1}O_{2n-1}^2 - p^{-4n+2}(p^2 - 4)(p^{4n}(p^8 + 2p^6x - 2p^4x + x^2) - 2p^6x^{n+1}),$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = 0$ and $p^2 \neq 2$ i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} \text{ or } x = p^4,$$

(and $p^2 \neq 2$) then

$$\sum_{k=0}^n x^k O_{2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2p^4(p^2 - 1)(p^2 - 3)x + p^8(p^2 - 1)(p^2 - 3))},$$

where

$$\Lambda = (p^2 - 4)((x - p^4(p^4 - 4p^2 + 2))x^{n+1} + (x - p^4)((n + 2)x - p^4(p^4 - 4p^2 + 2)(n + 1))x^n)O_{2n+1}^2 + (p^2 - 4)((n + 2)x - p^4(n + 1))x^n O_{2n-1}^2 + 2p^{-4n+2}(p^2 - 4)(p^{4n}(-x - p^6 + p^4) + p^6(n + 1)x^n),$$

and if $(x - p^4)(x^2 - p^4(p^4 - 4p^2 + 2)x + p^8) = (x - 4)(x + 4)^2 = 0$, $x \neq p^4 = 4$ and $p^2 = 2$ i.e.,

$$x = \frac{p^4(p^4 - 4p^2 + 2) + \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = \frac{p^4(p^4 - 4p^2 + 2) - \sqrt{p^{10}(p^2 - 4)(p^2 - 2)^2}}{2} = -4,$$

and $x \neq p^4 = 4$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n (-4)^k O_{2k+1}^2 = \frac{\Theta}{32},$$

where

$$\Theta = 64(-4)^{n-1} (n^2 - 2) O_{2n+1}^2 + (-4)^{n+1} (n+1)^2 O_{2n-1}^2 + 2^{-2n+4} p^{-2} (2^{2n} + 2n(n+1) (-4)^n).$$

(d): ($m = -1, j = 0$)

If $(p^2 x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) \neq 0$, i.e.,

$$x \neq \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2}, x \neq p^{-2},$$

then

$$\sum_{k=0}^n x^k O_{-k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-2})(x^2 - (p^2 - 2)p^{-2}x + p^{-4})},$$

where

$$\Phi = (x - p^{-2})(p^2 - 4)x^{n+1}O_{-n+1}^2 + (x - (p^2 - 2)p^{-2})(x - p^{-2})(p^2 - 4)x^{n+1}O_{-n}^2 + (p^2 - 4)p^{-4}(2p^{2n}x^n - p^2x - 1)x$$

and if $(p^2 x - 1)p^{-2}(x^2 - (p^2 - 2)p^{-2}x + p^{-4}) = 0$, i.e.,

$$x = \frac{(p^2 - 2)p^{-2} + \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = \frac{(p^2 - 2)p^{-2} - \sqrt{(p^2 - 4)p^{-2}}}{2} \text{ or } x = p^{-2},$$

then

$$\sum_{k=0}^n x^k O_{-k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)p^{-2}x + (p^2 - 1)p^{-4})},$$

where

$$\Lambda = (p^2 - 4)((n+2)x - p^{-2}(n+1))x^n O_{-n+1}^2 + (p^2 - 4)(p^4(n+3)x^2 - p^2x(p^2 - 1)(n+2) + (p^2 - 2)(n+1))p^{-4}x^n O_{-n}^2 + (p^2 - 4)p^{-4}(2p^{2n}(n+1)x^n - 2p^2x - 1)$$

(e): ($m = -2, j = 0$)

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, i.e.,

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k O_{-2k}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1}O_{-2n+2}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1}O_{-2n}^2 + (p^2 - 4)p^{-6}(2p^{4n}x^n - p^4x - 1)x$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, and $p^2 \neq 2$ i.e.,

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(and $p^2 \neq 2$, i.e., $p \neq -\sqrt{2}, p \neq \sqrt{2}$) then

$$\sum_{k=0}^n x^k O_{-2k}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$\Lambda = (p^2 - 4)((n + 2)x - p^{-4}(n + 1))x^n O_{-2n+2}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + (x - p^{-4})((n + 2)x - (p^4 - 4p^2 + 2)p^{-4}(n + 1))x^n) O_{-2n}^2 + (p^2 - 4)p^{-6}(2p^{4n}(n + 1)x^n - 2p^4x - 1),$
and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0, x \neq p^{-4} = \frac{1}{4}$ *and*
 $p^2 = 2$ *i.e.,*

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (*and* $p^2 = 2$, *i.e.,* $p = -\sqrt{2}, p = \sqrt{2}$) *then*

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k O_{-2k}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4\left(-\frac{1}{4}\right)^n (n + 1)^2 O_{-2n+2}^2 + \frac{1}{4}\left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) O_{-2n}^2 + 4p^{-2}(2^{2n}n(n + 1)\left(-\frac{1}{4}\right)^n + 1).$$

(f): ($m = -2, j = 1$)

If $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) \neq 0$, *i.e.,*

$$x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2}, x \neq p^{-4},$$

then

$$\sum_{k=0}^n x^k O_{-2k+1}^2 = \frac{\Phi}{(p^2 - 4)(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8})},$$

where

$$\Phi = (x - p^{-4})(p^2 - 4)x^{n+1} O_{-2n+3}^2 + (x - (p^4 - 4p^2 + 2)p^{-4})(x - p^{-4})(p^2 - 4)x^{n+1} O_{-2n+1}^2 + (p^2 - 4)p^{-14}(-p^8(p - 1)^2(p + 1)^2x^2 + p^4(p^4 - 4p^2 + 2)x + 2p^{4n+6}x^{n+1} - 1),$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = 0$, *and* $p^2 \neq 2$ *i.e.,*

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} \text{ or } x = p^{-4},$$

(*and* $p^2 \neq 2$, *i.e.,* $p \neq -\sqrt{2}, p \neq \sqrt{2}$) *then*

$$\sum_{k=0}^n x^k O_{-2k+1}^2 = \frac{\Lambda}{(p^2 - 4)(3x^2 - 2(p^2 - 1)(p^2 - 3)p^{-4}x + (p^2 - 1)(p^2 - 3)p^{-8})},$$

where

$$\Lambda = (p^2 - 4)((n + 2)x - p^{-4}(n + 1))x^n O_{-2n+3}^2 + (p^2 - 4)((x - (p^4 - 4p^2 + 2)p^{-4})x^{n+1} + (x - p^{-4})((n + 2)x - (p^4 - 4p^2 + 2)p^{-4}(n + 1))x^n) O_{-2n+1}^2 + (p^2 - 4)p^{-10}(2p^{4n+2}(n + 1)x^n - 2p^4(p - 1)^2(p + 1)^2x + p^4 - 4p^2 + 2),$$

and if $(x - p^{-4})(x^2 - (p^4 - 4p^2 + 2)p^{-4}x + p^{-8}) = \frac{1}{64}(4x - 1)(4x + 1)^2 = 0, x \neq p^{-4} = \frac{1}{4}$ *and*
 $p^2 = 2$ *i.e.,*

$$x = \frac{(p^4 - 4p^2 + 2)p^{-4} + \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = \frac{(p^4 - 4p^2 + 2)p^{-4} - \sqrt{(p^2 - 4)((p^2 - 2)^2 p^{-6}}}{2} = -\frac{1}{4},$$

and $x \neq p^{-4} = \frac{1}{4}$ (and $p^2 = 2$, i.e., $p = -\sqrt{2}, p = \sqrt{2}$) then

$$\sum_{k=0}^n \left(-\frac{1}{4}\right)^k O_{-2k+1}^2 = \frac{\Theta}{2},$$

where

$$\Theta = -4 \left(-\frac{1}{4}\right)^n (n+1)^2 O_{-2n+3}^2 + \frac{1}{4} \left(-\frac{1}{4}\right)^{n-1} (n^2 - 2) O_{-2n+1}^2 + p^{-2} (n(n+1) 2^{2n+1} \left(-\frac{1}{4}\right)^n + 1).$$

Note that in the last corollary, the case $p^2 \neq 4$ so that $p^2 - 4 \neq 0$ is considered. The sum formulas for the case $p^2 = 4$ so that $p^2 - 4 = 0$ is given in Soykan [8, Corollary 2.4].

Taking $x = 1$ in the last three corollaries we get the following corollary.

COROLLARY 6. *Suppose that $p^2 \neq 4$ so that $p^2 - 4 \neq 0$. For $n \geq 0$, modified p -Oresme numbers and p -Oresme-Lucas numbers and p -Oresme numbers have the following properties:*

(1) *modified p -Oresme numbers:*

(a): *If $p^2 \neq 1$ then*

$$\sum_{k=0}^n G_k^2 = \frac{\Phi}{(p^2 - 4)(1 - p^2)(2p^2 + 1)}$$

where

$$\Phi = (-p^4 + 2p^2 + 1)(1 - p^2)(p^2 - 4)G_n^2 + (1 - p^2)(p^2 - 4)G_{n-1}^2 - p^{-2n+4}(p^2 - 4)(p^{2n}(1 + p^2) - 2p^2)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n G_k^2 = \frac{1}{3}(2G_n^2 + G_{n-1}^2 + 2n - 1).$$

(b): *If $p^2 \neq 1$ then*

$$\sum_{k=0}^n G_{2k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^4)(2p^2 + 1)(2p^4 - 2p^2 + 1)}$$

where

$$\Phi = (-p^8 + 4p^6 - 2p^4 + 1)(1 - p^4)(p^2 - 4)G_{2n}^2 + (1 - p^4)(p^2 - 4)G_{2n-2}^2 - p^{-4n+8}(p^2 - 4)(p^{4n}(1 + p^4) - 2p^4)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n G_{2k}^2 = \frac{1}{3}(2G_{2n}^2 + G_{2n-2}^2 + 2n - 1).$$

(c): *If $p^2 \neq 1$ then*

$$\sum_{k=0}^n G_{2k+1}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^4)(2p^2 + 1)(2p^4 - 2p^2 + 1)}$$

where

$$\Phi = (-p^8 + 4p^6 - 2p^4 + 1)(1 - p^4)(p^2 - 4)G_{2n+1}^2 + (1 - p^4)(p^2 - 4)G_{2n-1}^2 - p^{-4n+4}(p^2 - 4)(p^{4n}(1 - 2p^4 + 2p^6 + p^8) - 2p^6)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n G_{2k+1}^2 = \frac{1}{3}(2G_{2n+1}^2 + G_{2n-1}^2 + 2n).$$

(d): If $p^2 \neq 1$ then

$$\sum_{k=0}^n G_{-k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-2})(2p^{-2} + p^{-4})}$$

where

$$\Phi = (1 - p^{-2})(p^2 - 4)G_{-n+1}^2 + 2p^{-2}(1 - p^{-2})(p^2 - 4)G_{-n}^2 + (p^2 - 4)p^{-2}(2p^{2n} - p^2 - 1)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n G_{-k}^2 = \frac{1}{3}(G_{-n+1}^2 + 2G_{-n}^2 + 2n - 1).$$

(e): If $p^2 \neq 1$ then

$$\sum_{k=0}^n G_{-2k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-4})(1 - (p^4 - 4p^2 + 2)p^{-4} + p^{-8})}$$

where

$$\Phi = (1 - p^{-4})(p^2 - 4)G_{-2n+2}^2 + (1 - (p^4 - 4p^2 + 2)p^{-4})(1 - p^{-4})(p^2 - 4)G_{-2n}^2 + (p^2 - 4)p^{-4}(2p^{4n} - p^4 - 1)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n G_{-2k}^2 = \frac{1}{3}(G_{-2n+2}^2 + 2G_{-2n}^2 + 2n - 1).$$

(f): If $p^2 \neq 1$ then

$$\sum_{k=0}^n G_{-2k+1}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-4})(1 - (p^4 - 4p^2 + 2)p^{-4} + p^{-8})}$$

where

$$\Phi = (1 - p^{-4})(p^2 - 4)G_{-2n+3}^2 + (1 - (p^4 - 4p^2 + 2)p^{-4})(1 - p^{-4})(p^2 - 4)G_{-2n+1}^2 + (p^2 - 4)p^{-12}(-p^8(p - 1)^2(p + 1)^2 + p^4(p^4 - 4p^2 + 2) + 2p^{4n+6} - 1)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n G_{-2k+1}^2 = \frac{1}{3}(G_{-2n+3}^2 + 2G_{-2n+1}^2 + 2n + 1).$$

(2) p -Oresme-Lucas numbers:

(a): If $p^2 \neq 1$ then

$$\sum_{k=0}^n H_k^2 = \frac{\Phi}{(p^2 - 4)(1 - p^2)(2p^2 + 1)}$$

where

$$\Phi = (p^2 - 1)(p^2 - 4)(p^4 - 2p^2 - 1)H_n^2 + (1 - p^2)(p^2 - 4)H_{n-1}^2 + p^{-2n+4}(p^2 - 4)(8 - 2p^2 - p^{2n}(p^2 + 5))$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n H_k^2 = \frac{1}{3}(2H_n^2 + H_{n-1}^2 + 6n + 3).$$

(b): If $p^2 \neq 1$ then

$$\sum_{k=0}^n H_{2k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^4)(1 - p^4(p^4 - 4p^2 + 2) + p^8)}$$

where

$$\Phi = (1 - p^4(p^4 - 4p^2 + 2))(1 - p^4)(p^2 - 4)H_{2n}^2 + (1 - p^4)(p^2 - 4)H_{2n-2}^2 + p^{-4n+4}(p^2 - 4)(p^{4n}(-p^8 - 12p^6 + 7p^4 + 4p^2 - 4) - 2p^6(p^2 - 4))$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n H_{2k}^2 = \frac{1}{3}(2H_{2n}^2 + H_{2n-2}^2 + 6n + 3).$$

(c): If $p^2 \neq 1$ then

$$\sum_{k=0}^n H_{2k+1}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^4)(1 - p^4(p^4 - 4p^2 + 2) + p^8)}$$

where

$$\Phi = (1 - p^4(p^4 - 4p^2 + 2))(1 - p^4)(p^2 - 4)H_{2n+1}^2 + (1 - p^4)(p^2 - 4)H_{2n-1}^2 - p^{-4n+4}(p^2 - 4)(p^{4n}(1 + 6p^4 - 2p^6 + p^8) + 2p^4(p^2 - 4))$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n H_{2k+1}^2 = \frac{1}{3}(2H_{2n+1}^2 + H_{2n-1}^2 + 6n).$$

(d): If $p^2 \neq 1$ then

$$\sum_{k=0}^n H_{-k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-2})(2p^{-2} + p^{-4})}$$

where

$$\Phi = (1 - p^{-2})(p^2 - 4)H_{-n+1}^2 + 2p^{-2}(1 - p^{-2})(p^2 - 4)H_{-n}^2 - (p^2 - 4)p^{-6}(p^6 + 4p^2 - 3p^4 + 2p^{2n+2}(p^2 - 4) + 4)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n H_{-k}^2 = \frac{1}{3}(H_{-n+1}^2 + 2H_{-n}^2 + 6n + 3).$$

(e): If $p^2 \neq 1$ then

$$\sum_{k=0}^n H_{-2k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-4})(1 - (p^4 - 4p^2 + 2)p^{-4} + p^{-8})}$$

where

$$\Phi = (1 - p^{-4})(p^2 - 4)H_{-2n+2}^2 + (1 - (p^4 - 4p^2 + 2)p^{-4})(1 - p^{-4})(p^2 - 4)H_{-2n}^2 + (p^2 - 4)p^{-12}(-p^8(p^2 - 2)^2 + p^4(3p^4 - 12p^2 + 8) - 2p^{4n+6}(p^2 - 4) - 4)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n H_{-2k}^2 = \frac{1}{3}(H_{-2n+2}^2 + 2H_{-2n}^2 + 6n + 3).$$

(f): If $p^2 \neq 1$ then

$$\sum_{k=0}^n H_{-2k+1}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-4})(1 - (p^4 - 4p^2 + 2)p^{-4} + p^{-8})}$$

where

$$\Phi = (1 - p^{-4})(p^2 - 4)H_{-2n+3}^2 + (1 - (p^4 - 4p^2 + 2)p^{-4})(1 - p^{-4})(p^2 - 4)H_{-2n+1}^2 + (p^2 - 4)p^{-12}(-p^8(p^2 - 3)^2 + p^4(p^4 - 4p^2 + 2) - 2p^{4n+4}(p^2 - 4) - 1)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n H_{-2k+1}^2 = \frac{1}{3}(H_{-2n+3}^2 + 2H_{-2n+1}^2 + 6n - 3).$$

(3) p -Oresme numbers:

(a): If $p^2 \neq 1$ then

$$\sum_{k=0}^n O_k^2 = \frac{\Phi}{(p^2 - 4)(1 - p^2)(2p^2 + 1)}$$

where

$$\Phi = (-p^4 + 2p^2 + 1)(1 - p^2)(p^2 - 4)O_n^2 + (1 - p^2)(p^2 - 4)O_{n-1}^2 - p^{-2n+2}(p^2 - 4)(p^{2n}(1 + p^2) - 2p^2)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n O_k^2 = \frac{1}{3}(2O_n^2 + O_{n-1}^2 + 2n - 1).$$

(b): If $p^2 \neq 1$ then

$$\sum_{k=0}^n O_{2k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^4)(1 - p^4(p^4 - 4p^2 + 2) + p^8)}$$

where

$$\Phi = (1 - p^4(p^4 - 4p^2 + 2))(1 - p^4)(p^2 - 4)O_{2n}^2 + (1 - p^4)(p^2 - 4)O_{2n-2}^2 - p^{-4n+6}(p^2 - 4)(p^{4n}(1 + p^4) - 2p^4)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n O_{2k}^2 = \frac{1}{3}(2O_{2n}^2 + O_{2n-2}^2 + 2n - 1).$$

(c): If $p^2 \neq 1$ then

$$\sum_{k=0}^n O_{2k+1}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^4)(1 - p^4(p^4 - 4p^2 + 2) + p^8)}$$

where

$$\Phi = (1 - p^4(p^4 - 4p^2 + 2))(1 - p^4)(p^2 - 4)O_{2n+1}^2 + (1 - p^4)(p^2 - 4)O_{2n-1}^2 + p^2(p^2 - 4)(2p^4 p^{4n} - 2p^6 p^{4n} - p^8 p^{4n} - p^{4n} + 2p^6)p^{-4n}$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n O_{2k+1}^2 = \frac{1}{3}(2O_{2n+1}^2 + O_{2n-1}^2 + 2n).$$

(d): If $p^2 \neq 1$ then

$$\sum_{k=0}^n O_{-k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-2})(2p^{-2} + p^{-4})}$$

where

$$\Phi = (1 - p^{-2})(p^2 - 4)O_{-n+1}^2 + 2p^{-2}(1 - p^{-2})(p^2 - 4)O_{-n}^2 + (p^2 - 4)p^{-4}(2p^{2n} - p^2 - 1)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n O_{-k}^2 = \frac{1}{3}(O_{-n+1}^2 + 2O_{-n}^2 + 2n - 1).$$

(e): If $p^2 \neq 1$ then

$$\sum_{k=0}^n O_{-2k}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-4})(1 - (p^4 - 4p^2 + 2)p^{-4} + p^{-8})}$$

where

$$\Phi = (1 - p^{-4})(p^2 - 4)O_{-2n+2}^2 + (1 - (p^4 - 4p^2 + 2)p^{-4})(1 - p^{-4})(p^2 - 4)O_{-2n}^2 + (p^2 - 4)p^{-6}(2p^{4n} - p^4 - 1)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n O_{-2k}^2 = \frac{1}{3}(O_{-2n+2}^2 + 2O_{-2n}^2 + 2n - 1).$$

(f): If $p^2 \neq 1$ then

$$\sum_{k=0}^n O_{-2k+1}^2 = \frac{\Phi}{(p^2 - 4)(1 - p^{-4})(1 - (p^4 - 4p^2 + 2)p^{-4} + p^{-8})}$$

where

$$\Phi = (1 - p^{-4})(p^2 - 4)O_{-2n+3}^2 + (1 - (p^4 - 4p^2 + 2)p^{-4})(1 - p^{-4})(p^2 - 4)O_{-2n+1}^2 + (p^2 - 4)p^{-14}(-p^8(p - 1)^2(p + 1)^2 + p^4(p^4 - 4p^2 + 2) + 2p^{4n+6} - 1)$$

and if $p^2 = 1$ then

$$\sum_{k=0}^n O_{-2k+1}^2 = \frac{1}{3}(O_{-2n+3}^2 + 2O_{-2n+1}^2 + 2n + 1).$$

Note that in the last corollary, the case $p^2 \neq 4$ so that $p^2 - 4 \neq 0$ is considered. The sum formulas for the case $p^2 = 4$ so that $p^2 - 4 = 0$ is given in Soykan [8, Corollary 2.5].

3. Conclusion

Recently, there have been so many studies of the sequences of numbers in the literature and the sequences of numbers were widely used in many research areas, such as architecture, nature, art, physics and engineering. In this work, sum identities were proved. The method used in this paper can be used for the other linear recurrence sequences, too. We have written sum identities in terms of the generalized p-Oresme sequence, and then we have presented the formulas as special cases the corresponding identity for the modified p-Oresme, p-Oresme-Lucas and p-Oresme numbers. All the listed identities in the corollaries may be proved by induction, but that method of proof gives no clue about their discovery. We give the proofs to indicate how these identities, in general, were discovered.

We can summarize the sections as follows:

- In section 1, we present some background about generalized p-Oresme numbers.
- In section 2, summation formulas have been presented for the generalized p-Oresme numbers. As special cases, summation formulas of modified p-Oresme, p-Oresme-Lucas and p-Oresme numbers have been given.

Competing Interests

Author have declared that no competing interests exist.

References

- [1] Clagett, M., Nicole Oresme and the Medieval Geometry of Qualities and Motions: A Treatise on the Uniformity and Difformity of Intensities Known as Tractatus de confurationibus qualitatum et motuum, The University of Wisconsin Press, Wisconsin, 1968.
- [2] Clagett, M., "Oresme, Nicole", C.C. Gillespie (ed.), Dictionary of Scientific Biography, Vol. 9, Charles Scribner's Sons, New York, 223–230, 1981.
- [3] Cook, C.K., Some sums related to sums of Oresme numbers, In: Howard F. T. (eds) Applications of Fibonacci Numbers, volume 9, Proceedings of the Tenth International Research Conference on Fibonacci Numbers and their Applications, Kluwer Academic Publishers, 87–99, 2004.
- [4] Horadam, A.F., Oresme Numbers, Fibonacci Quarterly, 12(3), 267-271, 1974.
- [5] Manguiera, M.C.S., Vieira, R.P.M., Alves, F.R.V., Catarino, P.M.C., The Oresme Sequence: The Generalization of its Matrix Form and its Hybridization Process, Notes on Number Theory and Discrete Mathematics, 27(1), 101-111, 2021. DOI: 10.7546/nntdm.2021.27.1.101-111.
- [6] Soykan, Y., Generalized Oresme Numbers, Earthline Journal of Mathematical Sciences, 7(2), 333-367, 2021. <https://doi.org/10.34198/ejms.7221.333367>
- [7] Soykan, Y., A Study On Generalized p-Oresme Numbers, Asian Journal of Advanced Research and Reports, 15(7), 1-25, 2021. ISSN: 2582-3248. DOI: 10.9734/AJARR/2021/v15i630410
- [8] Soykan, Y., A Study on the Sum of the Squares of Generalized Oresme Numbers: The Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$, Asian Journal of Pure and Applied Mathematics, 4(1), 16-27, 2022; Article no.AJPAM.679
- [9] Soykan, Y., On the Sum of the Squares of Generalized Mersenne Numbers: the Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$, International Journal of Advances in Applied Mathematics and Mechanics, 9(2), 28-37, 2021.

- [10] Soykan, Y., Taşdemir, E., Dikmen, C.M., A Study on the Sum of the Squares of Generalized Balancing Numbers: the Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$, Journal of Innovative Applied Mathematics and Computational Sciences, 1(1), 16–30, 2021.
- [11] Soykan, Y., Generalized Fibonacci Numbers with Indices in Arithmetic Progression and Sum of Their Squares: the Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$, Journal of Advances in Mathematics and Computer Science, 36(6), 30-62, 2021; ISSN: 2456-9968. DOI: 10.9734/JAMCS/2021/v36i630371
- [12] Soykan, Y., On the Sums of Squares of Generalized Tribonacci Numbers: Closed Formulas of $\sum_{k=0}^n x^k W_k^2$, Archives of Current Research International, 20(4), 22-47, 2020. DOI: 10.9734/ACRI/2020/v20i430187
- [13] Soykan, Y., Formulae For The Sums of Squares of Generalized Tribonacci Numbers: Closed Form Formulas of $\sum_{k=0}^n k W_k^2$, IOSR Journal of Mathematics, 16(4), 1-18, 2020. DOI: 10.9790/5728-1604010118
- [14] Soykan, Y., A Study On the Sums of Squares of Generalized Tribonacci Numbers: Closed Form Formulas of $\sum_{k=0}^n k x^k W_k^2$, Journal of Scientific Perspectives, 5(1), 1-23, 2021. DOI: <https://doi.org/10.26900/jsp.5.1.02>
- [15] Soykan, Y., On Generalized Tetranacci Numbers: Closed Form Formulas of the Sum $\sum_{k=0}^n W_k^2$ of the Squares of Terms, International Journal of Advances in Applied Mathematics and Mechanics, 8(1), 15-26, 2020.
- [16] Soykan, Y., A Study on Generalized Tetranacci Numbers: Closed Form Formulas $\sum_{k=0}^n x^k W_k^2$ of Sums of the Squares of Terms, Asian Research Journal of Mathematics, 16(10), 109-136, 2020. DOI: 10.9734/ARJOM/2020/v16i1030234