

Current Lipid Management Guidelines in Atherosclerotic Cardiovascular Disease

ABSTRACT

The use of coronary artery calcium (CAC) score to reclassify risk individuals at either borderline or intermediate risk, for whom the danger of statin therapy is unknown, is another significant addition to the 2018 recommendations. Increased risk of recurrent cardiovascular (CV) events is linked to having established atherosclerotic cardiovascular disease (ASCVD). Many modifiable risk factors for cardiovascular disease have been connected in the literature, including blood pressure, low-density lipoprotein (LDL), high-density lipoprotein (HDL), glucose intolerance, and smoking. High dose statins and strict adherence to dietary and lifestyle changes have been shown to reduce the risk of coronary events, with the advent of newer lipid-lowering medications, statins may one day become obsolete.

Keywords: High-density lipoprotein; cardiovascular disease; coronary artery calcium; Atherosclerosis.

1. INTRODUCTION

“Cardiovascular disease is the main cause of death, an estimated 17.9 million people died from CVDs in 2019, representing 32% of all global deaths” [1,2]. Cardiovascular disease kills 5 individuals every hour in Saudi Arabia, accounting for 37% of all fatalities in the kingdom [3]. Due to an unstable or ruptured plaque obstructing blood flow and raising the risk of CV events, early and vigorous treatment of Atherosclerosis may lower the risk of CV events [4-7]. The underlying cause of CVD that develops early in childhood is atherosclerosis caused by elevated LDL-C.


Increased risk of recurrent cardiovascular (CV) events is linked to having established atherosclerotic cardiovascular disease (ASCVD). “Many modifiable risk factors for cardiovascular disease have been connected in the literature, including blood pressure, low-density lipoprotein (LDL), high-density lipoprotein (HDL), glucose intolerance, and smoking” [8]. “Total cholesterol is more than or equivalent to 240 mg/dL, which includes LDL, HDL, and triglycerides. Low HDL levels are defined as serum HDL concentrations of less than 40 mg/dL. Low HDL levels were seen in 18.4% of people in the United States” [9].

“71 million (33.5%) of U.S. adults had an increased LDL level, with just 23 million (33%) of these people regulating their LDL levels, despite having a target LDL level of less than 100 mg/dL” [10]. Low-density lipoproteins (LDL) induce ASCVD, according to evidence from genetic, epidemiologic, and clinical investigations [11]. “Lowering LDL cholesterol (LDL-C) results in significant reductions in new ASCVD events, according to randomized clinical studies using cholesterol-lowering medications” [12,13].

Elevated lipids have the greatest impact on MI Risk, according to the INTERHEART Study [14]. Low-density lipoprotein (LDL) is strongly linked to coronary heart disease; every 39 mg/dL (1 mmol/L) reduction in LDL-C reduces the relative risk of CHD by 20-25 percent [15,16,17]. “However, despite awareness that lowering LDL-C levels in patients with acute coronary syndrome (ACS) provides benefits, few of these high-risk patients attain their LDL-C target goals” [18]. Lipoprotein(a), often known as Lp(a), is a modified version of LDL-C that has long been recognized as a CVD risk factor. It's worth testing in individuals who have a familial history of early ASCVD or a personal history of ASCVD that can't be explained by other risk factors [19].

2019 European Society of Cardiology/European Atherosclerosis Society (ESC/EAS) Guidelines and 2018 American Heart Association/American College of Cardiology (AHA/ACC) Guidelines (Tables 1,3)

Table 1. Recommended treatment goals for LDL-lower therapy: main changes from 2016-2019

EAS  **ESC**
European Society of Cardiology

**Recommended treatment goals for LDL-lowering therapy:
main changes from 2016 to 2019**

Risk category	LDL goals (starting with untreated LDL-C)	
	2016	2019
Very-high-risk	<1.8 mmol/L (70 mg/dL) or >50% ↓ if LDL-C 1.8-3.5 (70 - 135 mg/dL)	<1.4 mmol/L (55 mg/dL) and >50% ↓
High-risk	<2.6 mmol/L (100mg/dL) or >50% ↓ if LDL-C 2.6-5.2 (100 - 200 mg/dL)	<1.8 mmol/L (70 mg/dL) and >50% ↓
Moderate-risk	<3.0 mmol/L (115 mg/dL)	< 2.6 mmol/L (100 mg/dL)
Low-risk	<3.0 mmol/L (115 mg/dL)	<3.0 mmol/L (115 mg/dL)

Treatment goals for low-density lipoprotein cholesterol (LDL-C) across categories of total cardiovascular disease risk




Fig. 1. Treatment goals for low-density lipoprotein cholesterol (LDL-C) across categories of total cardiovascular disease risk

Table 3. Recommendations for treatment goals for low-density lipoprotein cholesterol

Recommendations for treatment goals for low-density lipoprotein cholesterol (1)

Recommendations	Class	Level
In secondary prevention patients at very-high risk ^c , an LDL-C reduction of at least 50% from baseline ^d and an LDL-C goal of <1.4 mmol/L (<55 mg/dL) are recommended.	I	A
In primary prevention, for individuals at very-high risk but without FH ^c , an LDL-C reduction of at least 50% from baseline ^d and an LDL-C goal of <1.4 mmol/L (<55 mg/dL) are recommended.	I	C
In primary prevention, for individuals with FH at very-high risk, an LDL-C reduction of at least 50% from baseline and an LDL-C goal of <1.4 mmol/L (<55 mg/dL) should be considered.	IIa	C

“Any patient with confirmed ASCVD (e.g., past ACS, stable angina, coronary revascularization, stroke, transient ischemic attack, or peripheral artery disease) is deemed extremely high risk, according to the recommendations. In very high-risk individuals, the ultimate treatment objective for secondary prevention is a 50% decrease in LDL-C from baseline and a target LDL-C of 55 mg/dL. Patients who do not meet their LDL-C objectives on maximally tolerated statin medication should be given a statin plus ezetimibe combination therapy. A proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor is advised in individuals who have not met their LDL-C target while using a maximally tolerated statin in conjunction with ezetimibe” [20].

“A history of multiple major ASCVD events or one major ASCVD event plus multiple high-risk conditions (age 65 years, heterozygous familial hypercholesterolemia, prior coronary artery bypass surgery or percutaneous coronary intervention outside of the major ASCVD event(s), diabetes mellitus, hypertension, chronic kidney disease, current smoker, persistently elevated LDL-C > or equal to 100 mg/dL on a maximally tolerated statin and ezetimibe, and a history of congestive heart failure). High-intensity or maximally tolerated statin medication should be given to patients with ASCVD who are at very high risk. It is appropriate to add ezetimibe to maximally tolerated statin medication if LDL-C is not below 70 mg/dL. It is appropriate to add a PCSK9 inhibitor to people on a maximally tolerated statin and ezetimibe medication who have not attained an LDL-C 70 mg/dL” [12,13].

2. PATHOPHYSIOLOGY

“Dyslipidemias are characterized by clinically elevated cholesterol and/or triglycerides, as well

as lower HDL levels. Genetic factors, high carbohydrate diets, excessive alcohol use, obesity, insulin resistance, and nephrotic syndrome can all contribute to increased hepatic VLDL production” [21]. “The absolute amount and cumulative length of LDL-C exposure have been directly linked to the risk of atherosclerotic cardiovascular disease (ASCVD)” [22]. “There is currently no evidence that HDL has a preventive effect in the prevention of atherosclerosis” [23]. Increased cholesterol levels in the body can cause plaque to develop and build up in the vasculature, resulting in atherosclerotic cardiovascular disease [24].

3. CLINICAL PRESENTATION

Hypercholesterolemia is a substantial risk factor for cardiovascular disease despite the lack of obvious symptoms. Xanthomas and corneal arcus are symptoms of high cholesterol levels, which are most commonly found in familial hypercholesterolemia. “Carotid artery disease, stroke, peripheral vascular disease, high blood pressure, and type 2 diabetes mellitus are all complications of poorly controlled hypercholesterolemia (T2DM). Psoriasis, Crohn’s disease, inflammatory bowel disease, chronic obstructive lung disease, depression, chronic pain, and chronic kidney disease are examples of systemic disorders that affect dyslipidemia” [25].

“According to 2017 observational research of 7,641 Europeans over 50, 1,591 (20.8 percent) of the participants had high triglyceride or low HDL values. These individuals were also more likely to be obese, have type 2 diabetes, and consume more alcohol than the recommended weekly limit. Patients with high T.G. and low HDL received no lipid treatment in 55% of cases” [26]. “When the ACC/AHA released the 2018

Guideline for the Management of Blood Cholesterol, it made more revisions to the 2013 lipid recommendations. It acknowledges that, while there is no optimum LDL-C blood level, it is critical to maintain those levels low, since people with LDL-C levels of 100mg/dL or less are less likely to suffer from heart disease and stroke. The 2018 guidelines recommend looking into risk-enhancing factors in older patients aged 40 to 75, such as family history and other factors of health conditions, in addition to risk factors that are commonly associated with cardiovascular diseases such as high cholesterol, high blood pressure, and smoking” [27].

“The use of coronary artery calcium (CAC) score (Table 4) to reclassify risk individuals at either

borderline or intermediate risk, for whom the danger of statin therapy is unknown, is another significant addition to the 2018 recommendations” [28]. Unless the patient has a very high risk of ASCVD, lifestyle changes such as reducing saturated fat consumption and exercising for at least 40 minutes three to four times per week are usually the initial step in therapy for high cholesterol [29]. With the emergence of non-statin medications like ezetimibe and bile acid sequestrants, the 2018 recommendations provide for more tailored care and treatment alternatives, but they also incorporate new risk-assessment tools to reduce inappropriate statin prescriptions.

Table 4. Agatston coronary artery calcium scoring [30]

CAC score(Agatston method)	Risk analysis	Clinical correlation
0	Absent/no risk	Low risk of future cardiovascular events
1-10	Minimal	Minimal atherosclerosis may be present with a low risk of future cardiovascular events
11-100	Mild	There is likely mild to minimum coronary artery stenosis. A mild risk of coronary artery disease exists
101-400	Moderate	Reasonable amount of plaque can be confirmed. Has a moderately increased risk of future cardiovascular events
>400	High	A high coronary calcium score correlated with a significant risk of having a cardiovascular event (such as myocardial ischemia) in near future

Statins and Ezetimibe: “Statins and healthy lifestyle changes are the first-line medication and healthy lifestyle modifications for lowering LDL-C to reduce CV risk, according to a large body of data from CV outcomes studies. In a review of many clinical studies, an ACC/AHA Expert Panel discovered that initiating moderate-intensity statin medication (to drop LDL-C by 30% to 50%) or high-intensity statin therapy (to lower LDL-C by 50%) is a crucial factor in reducing ASCVD occurrences. Ezetimibe is a selective cholesterol absorption inhibitor that reduces LDL-C levels by preventing cholesterol absorption in the gut. When this medicine is added to a statin regimen, the amount of LDL-C that is reduced increases by 20% to 25%” [12].

PCSK9 inhibitors: Monoclonal antibodies that attach to the PCSK9 protein are used in these treatments. PCSK9 normally blocks LDL receptor recycling by targeting the LDL receptor for destruction when it attaches to it, preventing the LDL receptor from returning to the hepatocyte's surface to bind new LDL particles [19].

On top of all types and dosages of statin medication and ezetimibe decrease in LDL-C by treatment and background therapy, evolocumab provides up to 75% additional LDL-C lowering [29,31,32]. A consistent decrease of LDL-C from a baseline of 60% was seen throughout all evolocumab clinical studies [32]. LDL-C Lowering has been shown To reduce atherosclerotic burden, and percent atheroma volume [33]. Patients who had a recent MI were more likely to have serious unfavourable CV events than those who had a remote MI. Evolocumab decreased the risk of the main endpoint by 19% in recent MI patients, and the risk of CV mortality, MI, or stroke was lowered by 25% [34].

“Inclisiran might be the first and only siRNA therapy to diminish LDL-C levels. It should be injected subcutaneously by a healthcare expert with an initial dosage, followed by a second dose three months later, and then every six months after that. Inclisiran promotes LDL-C absorption by hepatocytes and decreases LDL-C levels in the circulation by enhancing LDL-C receptor recycling and expression on the hepatocyte cell surface” [35]. Inclisiran is a cholesterol-lowering

small interfering RNA-based medication that got European Medicines Agency clearance in December 2020 for use as an addition to diet in individuals with primary hypercholesterolemia or mixed dyslipidemia. US food and drug administration (FDA) officially approved Inclisiran add on therapy to lower cholesterol on December 22, 2021 [36,37]. Evinacumab inhibits angiopoietin-like 3, a lipoprotein and endothelial lipase inhibitor that increases triglyceride and other lipid levels [38]. Vupanorsen inhibits the formation of the ANGPTL3 protein, lowering LDL-C levels in the process. Vupanorsen has also been demonstrated to lower total cholesterol, triglycerides and non-HDL-C [39]. Through its liver-directed downregulation of hepatic apolipoprotein C-III, Gemcabene can improve VLDL clearance in plasma and limit cholesterol and triglycerides synthesis in the liver [40]. "Bempedoic acid inhibits ATP-citrate lyase, a component of the cholesterol synthesis pathway" [41]. "The United States Food and Drug Administration (FDA) has authorized bempedoic acid as an adjuvant to diet and maximally tolerated statin medication for treating persons with heterozygous familial hypercholesterolemia or established ASCVD who require further LDL-C reduction" [42]. "In adults with elevated triglycerides (150 mg/dL) who have either established CVD or type 2 diabetes (T2D) and 2 CVD risk factors, icosapent ethyl is approved as an adjunct to maximally tolerated statin therapy to reduce the risk of MI, stroke, coronary revascularization, and unstable angina requiring hospitalization" [43]. Fibrates aid to reduce triglyceride levels by reducing VLDL synthesis in the liver and assisting in the removal of triglycerides from the blood. Fibrates can marginally enhance HDL-C levels and have varying effects on LDL-C levels, depending on the existence of other lipid disorders [44]. "Niacin is a B-complex vitamin that decreases triglycerides by 20% to 50%, lowers LDL-C by 10% to 20%, and improves HDL-C by 15% to 35%" [45]. Bile acid sequestrants bind to bile and prevent its absorption in the gut. Because bile is made up of cholesterol, these drugs lower total cholesterol and LDL-C levels in the body. Cholestyramine has been proven to decrease CV events in hypercholesterolemic males when given as monotherapy [17].

Lifestyle changes for ASCVD patients are an important aspect of secondary prevention. Smoking cessation, weight loss, and engaging in moderate-to-vigorous physical exercise 3–4 times per week for around 30-40 minutes per session are strongly advocated, in addition to

participating in cardiac rehabilitation programs whenever possible[12,13].

4. CONCLUSION

Low-density lipoprotein is a good indicator of a patient's lipid profile. LDL levels should be less than 100 mg/dL in healthy people; however, in those with heart disease or diabetes, they should be less than 70 mg/dL. Patients with a larger than 7.5 percent ASCVD risk, those with a history of familial hypercholesterolemia, patients with a documented history of atherosclerotic disease, and patients with diabetes mellitus all benefit from LDL reduction. Statins with high dose and strict adherence to dietary and lifestyle changes have been shown to reduce the risk of coronary events, with the advent of newer lipid lowering medications, statins may one day become obsolete.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO world health organization, cardiovascular key facts; 2019.
2. Murphy SL, Xu J, Kochanek KD, Arias E, Tejada-Vera B. Deaths: Final data for 2018. National Vital Statistics Reports. Available: <https://www.cdc.gov/nchs/data/>
3. WHO world health organization non-communicable diseases (NCD), country profile; 2015.
4. Pepine CJ. The effects of angiotensin-converting enzyme inhibition on endothelial dysfunction: potential role in myocardial ischemia. *Am J Cardiol.* 1998;82(10A): 23S-27S. DOI:10.1016/s0002-9149(98)00805-4
5. Ross R. Atherosclerosis--an inflammatory disease. *N Engl J Med.* 1999;340(2):115-126. DOI:10.1056/NEJM199901143400207
6. Stary HC, Chandler AB, Dinsmore RE, et al. A definition of advanced types of atherosclerotic lesions and a histological

- classification of atherosclerosis. A report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation*. 1995;92(5):1355-1374.
DOI:101161/01.cir.92.5.1355
7. Okazaki S, Yokoyama T, Miyauchi K, et al. Early statin treatment in patients with acute coronary syndrome: demonstration of the beneficial effect on atherosclerotic lesions by serial volumetric intravascular ultrasound analysis during half a year after coronary event: The ESTABLISH Study. *Circulation*. 2004;110(9):1061-1068.
DOI:101161/01.CIR.0000140261.58966.A4
8. Vasan RS, Sullivan LM, Wilson PW, et al. Relative importance of borderline and elevated levels of coronary heart disease risk factors, *Annals of Internal Medicine*. 2005;142(6):393–402.
9. Carroll MD, Fryar CD, Nguyen DT, National Center for Health Statistics (U.S.), "Total and high-density lipoprotein cholesterol in adults: the United States, 2015-2016," in U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics; 2017.
10. Centers for Disease Control and Prevention (CDC. (2011), "Disparities in diagnoses of HIV infection between blacks/A- African Americans and other racial/ethnic populations—37 states, 2005-2008," *MMWR. Morbidity and mortality weekly report*. 2011;60(4):93–98.
11. Ference BA, Ginsberg HN, Graham I, et al. Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel. *Eur Heart J*. 2017;38(32):2459-2472.
12. Grundy SM, Stone NJ, Bailey AL, et al; 2018AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;139:e1082-e1143. Erratum in: *Circulation*. 2019;139:e1182-e1186.
13. Mach F, Baigent C, Catapano AL, et al. 2019 ESC/EAS guidelines for managing dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J*. 2020;41:111-188. Erratum in: *Eur Heart J*. 2020;41:4255.
14. Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet*. 2004;364(9438):937-952.
DOI:101016/S0140-6736(04)17018-9
15. Kolodgie FD, Virmani R, Burke AP, et al. Pathologic assessment of the vulnerable human coronary plaque. *Heart*. 2004;90(12):1385-1391.
DOI:10.1136/hrt.2004.041798
16. Grundy SM, Cleeman JI, Merz CN, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines [published correction appears in *Circulation*. 2004;110(6):763]. *Circulation*. 2004;110(2):227-239.
doi:10.1161/01.CIR.0000133317.49796.0E
17. Cannon CP, Steinberg BA, Murphy SA, Mega JL, Braunwald E. Meta-analysis of cardiovascular outcomes trials comparing intensive versus moderate statin therapy. *J Am Coll Cardiol*. 2006;48(3):438-445.
DOI:101016/j.jacc.2006.04.070
18. European Society of Cardiology/ European Atherosclerosis Society (ESC/EAS) Guidelines; 2019.
19. Muneera Al Taweel, Sarah AlMukhaylid, Nada Alsultan. Latest in High-Risk Atherosclerotic Cardiovascular Disease Lipid GuidelinesMini Review. *Open J Cardiol Heart Dis*. 2022;3(5). JCHD.000572.
20. American Heart Association/ American College of Cardiology (AHA/ACC) Guidelines; 2018.
21. Raviglione MC, O'Brien RJ, Longo D, Fauci A, Kasper D, Hauser S. Harrison's principles of internal medicine, McGraw-Hill Education. 2012;1:2888–2892.
22. Ference BA, Ginsberg HN, Graham I, et al. Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel, *European Heart Journal*. 2017;38(32):2459–2472.
23. Libby P, Buring JE, Badimon L, et al. "Atherosclerosis," *Nature Reviews Disease Primers*. 2019;5:1.
24. Karr S. Epidemiology and management of hyperlipidemia. *The American Journal of*

- Managed Care. 2017;23;9 Supplement: S139–S148, 2017.
25. Hill MF, Bordoni B. Hyperlipidemia, StatPearls Publishing; 2020.
 26. Halcox JP, Banegas JR, Roy C, et al. Prevalence and treatment of atherogenic dyslipidemia in the primary prevention of cardiovascular disease in Europe: EUREKA, a cross-sectional observational study, BMC Cardiovascular Disorders. 2017;17:1.
 27. Grundy SM, Stone NJ, Bailey AL, et al. Guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines, Journal of the American College of Cardiology. 2019;73(24):e285–e350.
 28. Rosenson RS, Hegele RA, Fazio S, Cannon CP. The Evolving Future of PCSK9 Inhibitors. J Am Coll Cardiol. 2018;72:314-329.
 29. Robinson JG, et al. JAMA. 2014;311:1870-1882. Raal FJ, et al. Lancet. 2015;385:331-34
 30. Shreya D, Zamora DI, Patel GS, et al. Coronary Artery Calcium Score - A Reliable Indicator of Coronary Artery Disease?; 2021.
 31. Nissen SE et al. Presented at American Heart Association Annual Scientific Sessions; 2016, New Orleans, LA.
 32. Sabatine MS, Giugliano RP, Keech AC, et al. Evolocumab and Clinical Outcomes in Patients with Cardiovascular Disease. N Engl J Med. 2017;376(18):1713-1722. DOI:101056/NEJMoa1615664
 33. Puri R, Nissen SE, Ballantyne CM, et al. Factors underlying regression of coronary atheroma with potent statin therapy. Eur Heart J. 2013;34(24):1818-1825. DOI:101093/eurheartj/eh084
 34. Gencer B, et al. Poster presented at American Heart Association Scientific Sessions; 2019. Philadelphia, PA
 35. Ray KK, Wright RS, Kallend D, et al. Two phases 3 trials of inclisiran in patients with elevated LDL cholesterol, New England Journal of Medicine. 2020;382(16):1507–1519.
 36. Lamb YN. Inclisiran: First Approval [published correction appears in Drugs. 2021;81(9):1129]. Drugs. 2021;81(3):389-395. DOI:101007/s40265-021-01473-6
 37. FDA, US food and drug administration; 2022.
 38. Real FJ, Rosenson RS, Reeskamp LF, et al. Evinacumab for homozygous familial hypercholesterolemia, New England Journal of Medicine. 2020;383(8):711–720.
 39. Jia X, AlRifai M, Hussain A, Martin S, Agarwala A, Virani SS. Highlights from studies in cardiovascular disease prevention presented at the Digital 2020 European Society of Cardiology Congress: prevention is alive and well, Current Atherosclerosis Reports. 2020;22:12.
 40. Stein E, Bays H, Koren M, Bakker-Arkema R, Bisgaier C. Efficacy and safety of gemcabene as add-on to stable statin therapy in hypercholesterolemic patients. Journal of Clinical Lipidology. 2016;10(5):1212–1222.
 41. Laufs U, Banach M, Mancini GJ. "Efficacy and safety of bempedoic acid in patients with hypercholesterolemia and statin intolerance," Journal of the American Heart Association. 2019;8:7, article e011662.
 42. NEXLETOL (bempedoic acid) [Prescribing Information]. Ann Arbor, MI: Esperion Therapeutics
 43. VASCEPA (icosapent ethyl) [Prescribing Information]. Bridgewater, NJ: Amarin Pharma, Inc; 2019.
 44. Rubins H, Robins S, Collins D, et al. Gemfibrozil for the secondary prevention of coronary heart disease in men with low high-density lipoprotein cholesterol levels. Veterans Affairs High-density Lipoprotein Cholesterol Intervention Trial Study Group. N Engl J Med. 1999;341:410-418.
 45. Robinson J. Management of complex lipid abnormalities with a fixed-dose combination of simvastatin and extended-release niacin. Vasc Health Risk Man. 2009;5:31-43.

Peer-review history:
The peer review history for this paper can be accessed here: