

Healthy Lifestyles during Pregnancy is Associated with Better Maternal and Fetal Health

Abstract

Evidence indicates that healthy lifestyles can significantly improve maternal and fetal outcomes during pregnancy. Several major physiological changes take place during a normal pregnancy, all aimed at keeping the mother and the growing fetus healthy. Five major lifestyles: tobacco smoking, alcohol consumption, obesity, sedentary lifestyle, and inadequate nourishment can deleteriously alter these processes. Several myths, addictions, societal stigmas, and other factors often lie behind continuance of unhealthy lifestyles. Besides directly impacting the course of a normal pregnancy, fetal development, delivery and lactation with breastfeeding, these unhealthy lifestyles also impact several chronic diseases that invariably have a negative effect on pregnancy. This manuscript briefly reviews the effects of these lifestyles.

Keywords: pregnancy, smoking, exercise, alcohol, diet, obesity, maternal health, fetal health

Introduction

Pregnancy is associated with several physiological changes, including an increase in the production of estrogens, progesterone, prolactin, and the placental human chorionic gonadotropin¹⁻³. The gestational bodyweight increases, mainly due to the additional weight of the fetus, the placenta, enlarging uterus, amniotic fluid, mammary glands, and adipose tissue⁴. Several cardiac and hematological changes also occur. Plasma volume and cardiac output may increase by about 40% - resulting in peripheral vasodilatation and a decrease in systemic vascular resistance. There is a reduction in hemoglobin concentration, hematocrit, red blood cell and platelet count⁵. Oxygen demand increases by 20 to 30%. There is an increase in respiratory rate and an increase in ventilation. Several other systems are also affected, primarily to adapt to the growing needs of pregnancy⁶. These changes continue to evolve as the pregnancy progresses and start resolving with the birth of the baby and expulsion of the placenta⁶. Lactation and breastfeeding provide the offspring with hydration and nutrients for the first 4–6 months of life⁷. Pregnancy is a complex physiological process. Lifestyles play a significant role during the entire perinatal period. This is briefly discussed in this manuscript.

Discussion

The five major lifestyles that have a significant impact on human health are: smoking, obesity, exercise, alcohol intake and diet. They play a major role in the pathogenesis of most major chronic diseases. Li et al. estimated that avoidance of these five unhealthy lifestyles can significantly increase longevity⁸. In their estimate, adherence to healthy lifestyles at age 50, provide women 14.0 years, and men 12.2 years of additional life. These five lifestyles also have a major impact on the health of both the mother and the fetus before conception, during pregnancy and after childbirth.

Smoking/Tobacco Use

Most pregnant smokers are aware that smoking in pregnancy is harmful, although many lack a detailed knowledge of the associated risks⁹. However, despite this general awareness, approximately 12% of pregnant women in the United Kingdom and United States smoke throughout pregnancy¹⁰. Tobacco is the most common substance abused during pregnancy¹¹. Its use is associated increased maternal, fetal, and infant morbidity and mortality¹². Serious outcomes from tobacco use include ectopic pregnancy, increased risk of miscarriage, placenta previa, premature rupture of membranes¹³, preterm birth^{14,15}, antepartum and intrapartum stillbirth¹⁶, low birthweight^{17,18}, decreased head circumference, perinatal death¹⁹ and sudden unexpected infant death^{20,21}. Smoking women also tend not to breastfeed, wean offspring earlier, and produce less milk than non-smokers²². Waterpipe smoking may also increase the risk of delivering a low birthweight infant as well as other pregnancy complications²³. Snuff or chewing tobacco is also associated with an increased risk of stillbirth, low birthweight, prematurity, and infant death²⁴. Secondhand smoke (SHS) exposure to infants causes increased risk of SIDS and lower respiratory illness²⁵. Almost every developing organ system in the fetus, including the lungs, brain, heart, and ears, appears to be affected by prenatal exposure to tobacco^{26,27}. The damage is usually long-term, and results in future lower respiratory tract infections, asthma, otitis media, dental caries, hearing loss, and the metabolic syndrome²⁸. Exposure to secondhand smoke is also dangerous for pregnant women – studies have shown that they have a 20% greater chance of giving birth to a low birthweight baby than women who are not exposed to secondhand smoke (SHS) during pregnancy²⁹. Nicotine and cotinine are known to accumulate in concentrations two- to three-fold higher in breast milk than in plasma, posing additional risk in breastfeeding infants³⁰. Prenatal third hand smoke is also dangerous to the lungs in utero of the unborn child³¹. After birth, the newly born child may continue to be exposed to environmental dust with thirdhand smoke particles in a home where smoking is or was present³². Studies show that most women who smoked during pregnancy also continue to do so after the birth of the child³³. Among children exposed to tobacco smoke either prenatally or postnatally, increased rates of behavioral problems are seen very early in life³⁴. Newborns exposed in utero have heightened startle responses, tremors, hypertonicity, have more severe reactions to bowel movements and diapering, and tend to be fussier and cry more³⁴. By 18 months, children exposed to SHS in utero manifest increased externalizing behaviors, including defiance, not responding to punishment, inattentiveness, and hyperactivity³⁵. During early school years, children exposed prenatally to tobacco smoke also demonstrate increased rates of behavior problems. These include hyperactivity, oppositional defiant disorder, delinquency, and both internalizing and externalizing behaviors³⁶. An association with ADHD has also been noted³⁷. During pre-school years, children prenatally or postnatally exposed (also with SHS) show more behavioral problems than unexposed children^{38,39}. These include demanding attention, changes in mood, emotional instability, arguing, aggression, and destructive behavior^{39,40}. These behaviors are often persistent into adolescence^{41,42}. E-cigarettes, unlike cigarettes⁴³, exposes non-pregnant adults to lower levels of carcinogens and toxins⁴⁴. Exposure to second-hand e-cigarette vapor may pose less risk than exposure to second-hand cigarette smoke⁴⁵. However, although the data on any harm to the pregnant woman or her baby from exposure to vaping is limited⁴⁶⁻⁴⁸, precautions should be taken to avoid such an exposure. And finally, children of women who

smoke cigarettes are more likely to take up smoking themselves⁴⁹. There is obviously no safe level of exposure to tobacco smoke during pregnancy.

Obesity

Obesity (BMI>30) is increasing in pregnant women. It has recently become one of the most important health issues in pregnancy⁵⁰. Current Centers for Disease Control and Prevention (USA) data suggest excessive weight gain is reported in roughly 59% of overweight women and 56% of obese women⁵¹. Excessive weight gain and obesity impacts pregnancy even during the preconception period. Obesity in females is associated with subfertility and with a longer time to achieve pregnancy^{52,53}. Observational studies indicate that bariatric surgery improves fertility in women with obesity⁵⁴. Following conception, the complication rates in abnormally overweight pregnant women go up dramatically⁵⁵. They are more likely to experience miscarriage⁵⁶ – one meta-analysis showed that in women with a BMI>25, the increased risks had an odds ratio (OR) of 1.67⁵⁷. Preeclampsia and gestational hypertension, usually occurring in the second half of pregnancy or soon after childbirth in obese women, can lead to dangerous cardiovascular complications⁵⁸. A Swedish cohort study of 805,275 pregnancies found that 2.8% of women with a BMI of 29.1–35.0 had preeclampsia compared to 1.4% of women with a BMI of 19.8–26.0 (adjusted OR 2.62)⁵⁹. Duckitt et al. found that there was a 50% increase in the risk of preeclampsia in pregnant women who had an increased BMI at the outset and this risk doubled if the BMI was >35⁶⁰. Gestational diabetes is also dangerous. It leads to more cesarean sections, offspring born with high body weight, and a higher risk of T2DM in both the mother and the child in the future. One study found that almost 42% of mothers developed T2DM over a 10 year follow up following gestational diabetes⁶¹. Several studies have documented the high risk of gestational DM in overweight/obese pregnant women. A retrospective UK study of 287,213 pregnancies between 1989 and 1997 showed that after adjusting for confounding factors, women with a BMI ≥ 30 were more likely to develop gestational diabetes than women with a BMI of 20.0–24.9 (odds ratio 3.6)⁶². A subsequent Australian study of 14,230 pregnancies found that the risk of developing gestational diabetes was 2.95 times higher in obese women (BMI 30.01–40.00) compared with normal-weight (BMI 20.01–25.00) women⁵⁸. Obesity also increases the risk of death from VTE. There are several factors contributing to an increased risk of VTE in obese pregnant patients. These women often have reduced mobility, suffer from co-morbid conditions, increased frequency of operative delivery, and possibly higher levels of coagulation factors VIII and IX⁶³. In a United Kingdom study, pregnant women with a BMI ≥ 30 had an adjusted OR of 2.65 for antenatal pulmonary thromboembolism⁶⁴. The offspring is exposed to an increased risk of being still born or preterm^{65,66}. Ultrasound examination of the fetus may be difficult in obese pregnant women. In a systematic review and meta-analyses, Aune et al. determined that the relative risk for each five unit increase in maternal BMI in overweight and obese women was 1.21 for fetal death, 1.24 for stillbirth, 1.16 for perinatal death, 1.15 for neonatal death, and 1.18 for infant death⁶⁷. During labor, they experience problems with difficulty with anesthesia, and cesarean delivery. The BMI inversely proportional to the length of labor in nulliparous women⁶⁸. Cesarean delivery is 1.46 (unadjusted OR) in overweight and 2.05 (unadjusted OR) is more common in obese women, compared with normal weight women⁶⁹. Anesthesia is more problematic, with obese women – epidural failure is more common⁷⁰ and

endotracheal intubation may be difficult if general anesthesia is used⁷¹. Cesarean section is more likely to be complicated by surgical site infections in the overweight (odds ratio 1.6), obese class I (2.4), and obese classes II and III (3.7)⁷². Macrosomia (birth weight of $\geq 4,000$ grams regardless of gestational age) is more common and may complicate childbirth (especially birth trauma e.g., shoulder dystocia, fracture of clavicle, damage to the brachial plexus^{59,73}). The risk of congenital abnormalities in the offspring is also increased in overweight/obese pregnant women⁷⁴. A systematic review and meta-analysis found the increased risk as follows: spina bifida (Odds Ratio or OR: 2.24), neural tube defects (OR: 1.87), limb reduction anomalies (OR: 1.34), cardiovascular anomalies (OR: 1.30), and cleft lip and palate (OR: 1.20) – in obese when compared with non-obese women⁷⁵. The risk increases with increasing levels of obesity⁷⁶. Children born with excess bodyweight increase their risk of later life obesity⁷⁷. After delivery, obese women also demonstrate decreased rates of initiating breastfeeding and breastfeed for shorter durations compared to normal weight women⁷⁸. Obese women face several hurdles, such as delayed lactation, embarrassment with body size, larger breasts and nipples, and comorbid medical conditions, that may interfere with breast feeding. It also results in postpartum weight retention and obesity^{79,80}. They are also prone to long-term obesity and future co-morbidities such as diabetes and cardiovascular disease^{65,77}. According to the USA Institute of Medicine optimal gestational weight gain (GWG) in pregnancy of 11 to 20 lbs. (5 to 9 kg) for women who are obese (BMI >30.0 kg/m²) and 15 to 25 lbs. (6.8 to 11.3 kg) for women who are in the overweight (BMI of 25.0-29.9 kg/m²) category. They recommend a weight gain of 25 to 42 lbs. (about 11 to 19 kilograms) for obese women carrying twins or multiples. For underweight (BMI <18.5 kg/m²) women a weight gain of 28-40 lbs. (12.5-18 kg) and for normal weight women (BMI of 18.5-24.9 kg/m²), a weight gain of 25-35 lbs. (11.5-16 kg) is appropriate⁸¹. The optimal weight gain recommended may differ for Asian women.

Alcohol

It is estimated that an estimated 10% of women globally, consume alcohol during pregnancy⁸². Data suggests that in Europe about a quarter of women drink alcohol during pregnancy. However, this use is much higher in some countries such as Russia (36.5%), the United Kingdom (41.3%), Denmark (45.8%), Belarus (46.6%), and Ireland (60.4%)⁸². Alcohol use and binge drinking among women of childbearing age has been increasing in most countries globally⁸³. Denny et al found that (in a report published in 2019), almost one in nine (11.5%) of pregnant women drink alcohol during pregnancy and 3.9% report binge drinking in the past 30 days⁸⁴. Prenatal alcohol exposure (PAE) can affect multiple aspects of infant health and development⁸⁵. The harm from alcohol starts preconception, as it adversely affects the quality and quantity of nutrient intake⁸⁶. Since almost one half of all pregnancies in the United States are unintended, and often unrecognized early on, alcohol intake may harm a fetus during embryologic development (3–8 wk. of pregnancy)⁸⁷. The teratogenic effects of alcohol are most significant during this stage⁸⁸. Alcohol use in women who are sexually active and in the reproductive-aged, should therefore be, (current recommendation from the American College of Obstetricians and Gynecologists, Center for Disease Control (CDC), Surgeon General, and medical societies from other countries including the Society of Obstetricians and Gynecologists of Canada) completely avoided⁸⁹⁻⁹¹. Alcohol exposure during the second and third trimester is associated with neuronal

loss⁹²⁻⁹⁴. Overall, alcohol use during pregnancy may lead to several detrimental effects which include miscarriage, preterm labor, intrauterine growth restriction, and stillbirth^{95,96}. Alcohol consumption also may lead to the fetal alcohol spectrum disorder (FASD). This includes growth retardation, craniofacial dysmorphisms, central nervous system dysfunction, or neurobehavioral disabilities in the child⁹⁷⁻⁹⁹. Although alcohol use specifically is harmful to the fetus, pregnant and parenting women who use alcohol often shy away from necessary health care¹⁰⁰. The related fear of a punitive response may result in avoidance of the much-needed health and social services. To summarize, there is no safe threshold for alcohol consumption during pregnancy¹⁰¹ and it should be totally avoided during pregnancy and breastfeeding¹⁰². Alcohol consumption and its harmful effects should be discussed with all pregnant women as early in the pregnancy as possible and at every prenatal visit¹⁰³. WHO also recommends detoxification if the pregnant woman is addicted to alcohol¹⁰⁴. Withdrawal symptoms (visual, auditory, and tactile disturbances, tremors, sweating, agitation, anxiety), if severe may require inpatient monitoring during detoxification¹⁰⁵.

Physical Activity

Exercise tends to decrease in many women with pregnancy^{106,107}. It is estimated that almost 60% of women are inactive in pregnancy¹⁰⁸, only 15%–38% of pregnant women follow physical activity guidelines¹⁰⁹, and only 8% meet the recommendations in the 3rd trimester¹¹⁰. Moreover, the decrease in levels of physical activity during pregnancy tends to persist up for to six months after delivery¹¹¹. Several barriers contribute towards this increase in sedentary behavior, especially worries that physical activity might contribute to the mother suffering a miscarriage. Other justifications include the nausea experienced during pregnancy, together with increased body weight and size¹¹². Physical activity and physical exercise during pregnancy is not only safe but helps improve maternal and fetal health and prevents several pregnancy-related complications^{113,114}. It helps prevent weight gain¹¹⁵. Exercise before and during pregnancy helps reduce the risk of maternal diabetes¹¹⁶ and if present, makes it easier to control¹¹⁷. It may also help prevent preeclampsia¹¹⁸. Regular exercise may shorten the duration of labor and reduce the risk of Cesarean section and operative-assisted vaginal delivery¹¹⁹. Exercise also helps reduce peri-natal depression¹²⁰. PA does not lead to adverse effects on birthweight and does not increase the risk of stillbirth. Fetal benefits include reduced preterm birth and decreased fat mass¹²¹. There is no detrimental effect on lactation or infant growth¹²². Even non-traditional exercise like yoga help reduce several pregnancies related complications^{123,124}. Exercise also helps relieve stress, lowers blood pressure and cholesterol, decreases depression, and enhances self-confidence^{125,126}. In general, exercise reduces the morbidity and mortality associated with cardiovascular disease, hypertension, and type 2 diabetes mellitus and several other chronic diseases¹²⁷. Several professional associations, including the American College of Obstetrics and Gynecology (ACOG) recommends that pregnant women should perform at least 30 minutes of moderate exercise, five days per week^{128,129}. Aerobic exercise can be safely initiated in the first trimester, and continued until delivery^{130,131}. Resistance exercises, properly designed, may also help complement aerobic exercises^{132,133}. Besides improving cardiorespiratory fitness, they also help decrease both weight gain, decrease lower limb edema, and mitigate depression^{134,135}. Exercises may be modified as the pregnancy progresses. Postpartum exercises also help urinary urge

incontinence decrease lumbopelvic pain, and help strengthen the recti abdominus muscles^{136,137}. **In general, exercises that harm should be avoided.** These include exercise above 5250 feet, those increasing the risk of contact, falling, or abdominal trauma, and scuba diving¹³⁸. Hot yoga and hot Pilates may also cause harm. Women may not be able to exercise if they have severe preeclampsia, severe respiratory/heart/endocrine/hematological or neurological diseases. They may also be unable to exercise if they suffer from placenta previa, second or third trimester bleeding, vasa previa, intrauterine growth restriction, and preterm labor^{139,140}. These patients can minimize the risk of VTE by ambulating regularly rather than exercising¹⁴¹. Pregnant women should stop exercising and seek immediate medical attention if there are unexplained symptoms – such as dizziness, shortness of breath, chest pain or calf pain. Many symptoms related to pregnancy are also indications for seeking medical advice and include painful contractions, vaginal bleeding, a leak of the amniotic fluid or rupture of membranes^{142,143}.

Nutrition

Adherence to the nutritional recommendations is low in pregnant women^{144,145}. Most women start avoiding alcohol and caffeine when they find out that they are pregnant¹⁴⁶. However, diet modification during pregnancy should go beyond that¹⁴⁷. Non-pregnancy requirement for protein is generally 0.8g/kg/day. However, in the pregnant women, this goes up to 1.1g/kg/day¹⁴⁸. Pregnant women should consume 45-64% of daily calories from carbohydrates and 20-35% of daily calories from fats¹⁴⁷. The micronutrient intake also goes up – both during pregnancy and during lactation. Three micronutrients are specifically important – folic acid, iron and B12¹⁴⁹. The requirements for folic acid increase 10-20-fold, for iron two- to three-fold and for B12, two-fold¹⁴⁹. Folic acid is necessary to support cell growth and nucleotide synthesis for fetal and placental development. Several studies have shown that folic acid supplementation helps reduce neural tube defects in the offspring¹⁵⁰. It is recommended that folate consumption should increase from 400 µg/d to 600 µg/d during pregnancy and should be around 500 µg/d during lactation¹⁵⁰. Another micro-nutrient – iron, needs to nearly double in intake during pregnancy – the requirements go up from 18 mg/d to 27 mg/d. Iron deficiency anemia (ferritin <15 µg/L,) can cause harm and needs to be aggressively treated. Although other micronutrient requirements also go up, these can usually be provided by intake of a daily prenatal multivitamin before conception and during pregnancy. Additional 500 additional kcal/day intake is needed by breast feeding women beyond what is recommended for **non-pregnant women to ensure an appropriate amount of weight gain by the newborn**¹⁵¹. Requirements of many micronutrients also continue to be high during this postnatal lactating phase. These can be usually remedied by taking a prenatal vitamin daily. Vegetarian diets during pregnancy have been found to have no protective associations with adverse maternal outcomes. The fluid requirements during pregnancy also increase by about 300 ml per day during the last few months of pregnancy¹⁵².

A proper diet during pregnancy (and the peri-pregnancy) period is a crucial in preventing deleterious pregnancy outcomes¹⁵³⁻¹⁶⁷. These include maternal ailments such as unwanted obesity, preeclampsia, gestational hypertension, and postpartum depression¹⁵³⁻¹⁵⁴. The impact on the fetus (especially from an inadequate intake of folate and other B vitamins by the pregnant mother) may cause preterm birth, neural defects, cleft palate, and congenital heart disease¹⁵⁵⁻¹⁶⁰.

The neonatal may experience eczema, cough and wheezing during the first year of life^{161,162}. The growing infant may develop obesity, hypertension and adverse cardiovascular outcomes¹⁶³. Inadequate intake of vitamin B12 may also result in long term damage to the child's nervous system^{158,159,164}. A vitamin D and/or iodine deficient diet may affect bone mineralization in the offspring¹⁶⁵⁻¹⁶⁷. Adequate and balanced intake of macronutrients and a multivitamin supplement before, during and often after pregnancy is therefore extremely crucial for good maternal and fetal health¹⁶⁸.

Conclusion

Lifestyles are extremely important in modifying the results of pregnancy. Obesity has a harmful relationship with pregnancy, and this is often unrecognized, overlooked, or ignored. Obesity impacts pregnancy even before conception as it reduces fertility. During pregnancy, excess body weight significantly increases the risk of maternal and fetal complications. The repercussion persist into future life of both the mother and the child. The quality of diet also makes a significant impact on maternal and fetal health. The increased protein requirement and the critical role played by folic acid and iron in the health of the growing fetus is well recognized. The Institute of Medicine dietary guidelines are helpful. The deleterious role of smoking and alcohol for both the pregnant mother and the offspring is well recognized. Even a single cigarette or alcoholic drink are unsafe in pregnancy. And finally, the myth of exercise not being safe during pregnancy has now long been proven false. Most healthy nulliparas are advised about 30 minutes of moderate intensity exercise about 5 days a week, leading to better maternal and offspring health. The adoption of healthy lifestyles should begin before pregnancy and continue through the postpartum period.

Acknowledgement: None

Funding: None

Conflict of Interest: None

References

1. <https://americanpregnancy.org/>.
2. ACOG Committee on Obstetric Practice and Society for Maternal-Fetal Medicine. (2013; Reaffirmed 2015). Committee Opinion No. 579. Definition of term pregnancy.
3. Soma-Pillay P., Nelson-Piercy C., Tolppanen H., Mebazaa A. Physiological changes in pregnancy. *Cardiovasc. J. Afr.* 2016;27:89–94. doi: 10.5830/CVJA-2016-021.
4. Institute of Medicine . Weight Gain during Pregnancy: Reexamining the Guidelines. National Academies Press; Washington, DC, USA: 2009.
5. Tkachenko O., Shchekochikhin D., Schrier R.W. Hormones and Hemodynamics in Pregnancy. *Int. J. Endocrinol. Metab.* 2014;12:e14098. doi: 10.5812/ijem.14098.

6. Soma-Pillay P., Nelson-Piercy C., Tolppanen H., Mebazaa A. Physiological changes in pregnancy. *Cardiovasc. J. Afr.* 2016;27:89–94. doi: 10.5830/CVJA-2016-021.
7. Truchet S., Honvo-Houéto E. Physiology of milk secretion. *Pract. Res. Clin. Endocrinol. Metab.* 2017;31:367–384. doi: 10.1016/j.beem.2017.10.008.
8. Li Yanping, An Pan, Dong D. et al. Impact of Healthy Lifestyle Factors on Life Expectancies in the US Population. *Circulation.* 2018;138:345–355. <https://doi.org/10.1161/CIRCULATIONAHA.117.032047>.
9. Flemming K, McCaughan, Angus K, et al. Qualitative systematic review: barriers and facilitators to smoking cessation experienced by women in pregnancy and following childbirth. *J Adv Nurs.* 2014;71(6):1210–1226. doi:10.1111/jan.12580.
10. NHS Information Centre. Infant Feeding Survey 2010: Early Results. The NHS Information Centre; 2011. ;Centers for Disease Control and Prevention. Pregnancy Risk Assessment Monitoring System 2009–2010. National Center for Chronic Disease Prevention and Health Promotion; 2009.
11. Oh S, Reingle Gonzalez JM, Salas-Wright CP, Vaughn MG, DiNitto DM. Prevalence and correlates of alcohol and tobacco use among pregnant women in the United States: Evidence from the NSDUH 2005-2014. *Prev Med.* 2017 Apr;97:93-99.
12. Committee opinion no. 721 summary: smoking cessation during pregnancy. *Obstetrics and Gynecology.* 2017; 130(4): 1. DOI: 10.1097/AOG.0000000000002348.
13. Centers for Disease Control and Prevention. The health consequences of smoking: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services; 2004.
14. Kelkay B, Omer A, Teferi Y, Moges Y. Factors associated with singleton preterm birth in Shire Suhul General Hospital, Northern Ethiopia, 2018. *Journal of Pregnancy.* 2019; 4629101. DOI: 10.1155/2019/4629101.
15. Liu B, Xu G, Sun Y, et al. Maternal cigarette smoking before and during pregnancy and the risk of preterm birth: a dose-response analysis of 25 million mother-infant pairs. *PLoS Medicine.* 2020; 17(8): e1003158. DOI: 10.1371/journal.pmed.1003158.
16. Bjørnholt SM, Leite M, Albieri V, Kjaer SK, Jensen A. Maternal smoking during pregnancy and risk of stillbirth: results from a nationwide Danish register-based cohort study. *Acta Obstetrica et Gynecologica Scandinavica.* 2016; 95(11): 1305–1312. DOI: 10.1111/aogs.13011.
17. Philips, EM, Santos S, Trasande L, et al. Changes in parental smoking during pregnancy and risks of adverse birth outcomes and childhood overweight in Europe and North America: an individual participant data meta-analysis of 229,000 singleton births. *PLoS Medicine.* 2020; 17(8): e1003182. DOI: 10.1371/journal.pmed.1003182.
18. Tong VT, England LJ, Rockhill KM, D'Angelo DV. Risks of preterm delivery and small for gestational age infants: effects of nondaily and low-intensity daily smoking during pregnancy. *Paediatric and Perinatal Epidemiology.* 2017; 31(2): 144–148. DOI: 10.1111/ppe.12343.
19. CInerney C, Ibiebele I, Ford JB, et al. Benefits of not smoking during pregnancy for Australian Aboriginal and Torres Strait Islander women and their babies: a retrospective

- cohort study using linked data. *BMJ Open*. 2019; 9(11): e032763. DOI: 10.1136/bmjopen-2019-032763.
20. Anderson TM, Lavista Ferres JM, Ren SY, et al. Maternal smoking before and during pregnancy and the risk of sudden unexpected infant death. *Pediatrics*. 2019; 143(4): e20183325. DOI: 10.1542/peds.2018-3325.
 21. Prabhu N, Smith N, Campbell D, et al. First trimester maternal tobacco smoking habits and fetal growth. *Thorax* 2010;65(3):235–40.
 22. Amir LH. Maternal smoking and reduced duration of breastfeeding: a review of possible mechanisms. *Early Hum Dev*. 2001 Aug;64(1):45-67. doi: 10.1016/s0378-3782(01)00170-0
 23. Maziak W, Ward KD, Afifi Soweid RA, Eissenberg T. Tobacco smoking using a waterpipe: a re-emerging strain in a global epidemic. *Tob Control*. 2004;13(4):327–33. ; Mirahmadizadeh A, Nakhaee N. Prevalence of waterpipe smoking among rural pregnant women in Southern Iran. *Med Princ Pract*. 2008;17(6):435–9.
 24. Gupta PC, Sreevidya S. Smokeless tobacco use, birthweight, and gestational age: population based, prospective cohort study of 1217 women in Mumbai, India. *BMJ*. 2004;328(7455):1538. ; Gupta PC, Subramoney S. Smokeless tobacco use and risk of stillbirth: a cohort study in Mumbai, India. *Epidemiology*. 2006;17(1):47–51.
 25. Centers for Disease Control and Prevention. The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services; 2006.
 26. Joad JP. Smoking and pediatric respiratory health. *Clin Chest Med* 2000;21(1):37–46 [vii].
 27. Cook DG, Strachan DP. Health effects of passive smoking. 3. Parental smoking and prevalence of respiratory symptoms and asthma in school age children. *Thorax* 1997;52(12): 1081–94.
 28. Gergen PJ, Fowler JA, Maurer KR, Davis WW, Overpeck MD. The burden of environmental tobacco smoke exposure on the respiratory health of children 2 months through 5 years of age in the United States: third National Health and Nutrition Examination Survey, 1988 to 1994. *Pediatrics* 1998;101(2):[E8].
 29. ;;
 30. Luck W, Nau H. Nicotine and cotinine concentrations in serum and urine of infants exposed via passive smoking or milk from smoking mothers. *J Pediatr* 1985;107(5):816–20.
 31. Rehan VK, Sakurai R, Torday JS. Thirdhand smoke: a new dimension to the effects of cigarette smoke on the developing lung. *Am J Physiol Lung Cell Mol Physiol* 2011;301(1):L1–8.
 32. Dreyfuss JH. Thirdhand smoke identified as potent, enduring carcinogen. *CA Cancer J Clin* 2010;60(4):203–4.
 33. The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. Atlanta, GA; 2006. A study by Weitzman et al. Weitzman M, Gortmaker S, Sobol A. Maternal smoking and behavior problems of children. *Pediatrics* 1992;90(3):342–9.

34. Reijneveld SA, Brugman E, Hirasing RA. Excessive infant crying: definitions determine risk groups. *Arch Dis Child* 2002;87(1):43–4.
35. Stene-Larsen K, Borge AIH, Vollrath M. Maternal smoking in pregnancy and externalizing behavior in 18-month-old children: results from a population-based prospective study. *J Am Acad Child Adolesc Psychiatry* 2009;48(3):283–9.
36. Twardella D, Bolte G, Fromme H, Wildner M, von Kries R. Exposure to secondhand tobacco smoke and child behaviour —results from a cross-sectional study among preschool children in Bavaria. *Acta Paediatr* 2010;99(1):106–11.
37. Gatzke Kopp L, Beauchaine T. Direct and passive prenatal nicotine exposure and the development of externalizing psychopathology. *Child Psychiatry Hum Dev* 2007;38(4):255–69.
38. Kahn R, Zuckerman B, Bauchner H, Homer C, Wise P. Women’s health after pregnancy and child outcomes at age 3 years: a prospective cohort study. *Am J Public Health* 2002;92(8):1312–8.
39. Day NL, Richardson GA, Goldschmidt L, Cornelius MD. Effects of prenatal tobacco exposure on preschoolers’ behavior. *J Dev Behav Pediatr* 2000;21(3):180–8.
40. Williams GM, O’Callaghan M, Najman JM, et al. Maternal cigarette smoking and child psychiatric morbidity: a longitudinal study. *Pediatrics* 1998;102(1):[E11].
41. Wakschlag LS, Lahey BB, Loeber R, Green SM, Gordon RA, Leventhal BL. Maternal smoking during pregnancy and the risk of conduct disorder in boys. *Arch Gen Psychiatry* 1997; 54(7):670–6.
42. Indredavik M, Brubakk A-M, Romundstad P, Vik T. Prenatal smoking exposure and psychiatric symptoms in adolescence. *Acta Paediatr* 2007;96(3):377–82.
43. McNeill, A., Brose, L. S., Calder, R., Hitchman, S. C., Hajek, P., & McRobbie, H. (2015). E-cigarettes: an evidence update. A report commissioned by Public Health England (PHE publications gateway number: 2015260).
44. Goniewicz M.L., Smith D.M., Edwards K.C., Blount B.C., Caldwell K.L., Feng J...Hyland A.J. Comparison of nicotine and toxicant exposure in users of electronic cigarettes and combustible cigarettes. *JAMA Network Open*. 2018;1(8):e185937. doi: 10.1001/jamanetworkopen.2018.5937.
45. Hess, I. M., Lachireddy, K., & Capon, A. (2016). A systematic review of the health risks from passive exposure to electronic cigarette vapour. *Public Health Research & Practice*, 26(2). 10.17061/phrp2621617.6.
46. Cardenas V., Cen R., Clemens M., Moody H., Ekanem U., Policherla A....Boysen G. Use of Electronic Nicotine Delivery Systems (ENDS) by pregnant women I: Risk of small-for-gestational-age birth. *Tobacco Induced Diseases*. 2019;17(May) doi: 10.18332/tid/106089.
47. Froggatt S., Covey J., Reissland N. Infant neurobehavioural consequences of prenatal cigarette exposure: A systematic review and meta-analysis. *Acta Paediatrica*. 2020;109(6):1112–1124. doi: 10.1111/apa.v109.610.1111/apa.15132.
48. McDonnell B.P., Dicker P., Regan C.L. Electronic cigarettes and obstetric outcomes: A prospective observational study. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2020;127(6):750–756. doi: 10.1111/bjo.v127.610.1111/1471-0528.16110.

49. Leonardi-Bee J., Jere M.L., Britton J. Exposure to parental and sibling smoking and the risk of smoking uptake in childhood and adolescence: A systematic review and meta-analysis. *Thorax*. 2011;66(10):847–855. doi: 10.1136/thx.2010.153379.
50. <https://www.acog.org/womens-health/faqs/obesity-and-pregnancy>.
51. Flegal K, Carroll M, Kit B, et al. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA*. 2012;307:491–497.
52. Gesink Law DC, Maclehose RF, Longnecker MP. Obesity and time to pregnancy. *Hum Reprod*. 2007;22(2):414–420.
53. Nohr EA, Timpson NJ, Andersen CS, Davey Smith G, Olsen J, Sorensen TI. Severe obesity in young women and reproductive health: the Danish National Birth Cohort. *PLoS One*. 2009;4(12):e8444.
54. Milone M, De Placido G, Musella M, et al. Incidence of Successful Pregnancy After Weight Loss Interventions in Infertile Women: a Systematic Review and Meta-Analysis of the Literature. *Obes Surg*. 2016;26(2):443–451.
55. Fitzsimons KJ, Modder J, Greer IA. Obesity in pregnancy: risks and management. *Obstet Med*. 2009 Jun;2(2):52-62.
56. Lu GC, Rouse DJ, DuBard M, Cliver S, Kimberlin D, Hauth JC. The effect of the increasing prevalence of maternal obesity on perinatal morbidity. *Am J Obstet Gynecol*. 2001 Oct;185(4):845–849.
57. Metwally M, Ong KJ, Ledger WL, Li TC. Does high body mass index increase the risk of miscarriage after spontaneous and assisted conception? A meta-analysis of the evidence. *Fertil Steril* 2008;356:714-26. 10.1016/j.fertnstert.2007.07.1290.
58. Callaway LK, Prins JB, Chang AM, McIntyre HD. The prevalence and impact of overweight and obesity in an Australian obstetric population. *Med J Aust* 2006;184:56–9,
59. Cedergren MI. Maternal morbid obesity and the risk of adverse pregnancy outcome. *Obstet Gynecol* 2004;103:219–24.
60. Duckitt K, Harrington D. Risk factors for pre-eclampsia at antenatal booking: systematic review of controlled studies. *BMJ*. 2005 Mar 12;330(7491):565. doi: 10.1136/bmj.38380.674340.E0/
61. Lauenborg J, Hansen T, Jensen DM, et al. Increasing incidence of diabetes after gestational diabetes: A long-term follow-up in a Danish population. *Diab Care* 2004;27:1194–9.
62. Sebire NJ, Jolly M, Harris JP, Wadsworth J, Joffe M, Beard RW, Regan L, Robinson S. Maternal obesity and pregnancy outcome: a study of 287,213 pregnancies in London. *Int J Obes Relat Metab Disord*. 2001 Aug;25(8):1175-82. doi: 10.1038/sj.ijo.0801670.
63. Abdollahi M, Cushman M, Rosendaal FR. Obesity: risk of venous thrombosis and the interaction with coagulation factor levels and oral contraceptive use. *Thromb Haemost* 2003;89:493–8.
64. Knight M; UKOSS. Antenatal pulmonary embolism: risk factors, management and outcomes. *BJOG*. 2008 Mar;115(4):453-61. doi: 10.1111/j.1471-0528.2007.01622.x.
65. Poston, L. , Caleyachetty, R. , Cnattingius, S. , Corvalán, C. , Uauy, R. , Herring, S. , & Gillman, M. W. (2016). Preconceptional and maternal obesity: epidemiology and health

- consequences. *The Lancet Diabetes and Endocrinology*, 4(12), 1025–1036. 10.1016/S2213-8587(16)30217-0.
66. Vézina-Im, L.-A. , Nicklas, T. A. , & Baranowski, T. (2018). Intergenerational effects of health issues among women of childbearing age: A review of the recent literature. *Current Nutrition Reports*, 7(4), 274–285. 10.1007/s13668-018-0246-x).
 67. Aune D, Saugstad OD, Henriksen T, Tonstad S. Maternal body mass index and the risk of fetal death, stillbirth, and infant death: a systematic review and meta-analysis. *JAMA* 2014;356:1536-46. 10.1001/jama.2014.2269 pmid:24737366.
 68. Nuthalapaty FS, Rouse DJ, Owen J. The association of maternal weight with cesarean risk, labor duration, and cervical dilation rate during labor induction. *Obstet Gynecol* 2004;356:452-6.
 69. Chu SY, Kim SY, Schmid CH, et al. Maternal obesity and risk of cesarean delivery: a meta-analysis. *Obes Rev* 2007;356:385-94. 10.1111/j.1467-789X.2007.00397.x.
 70. Mhyre JM. Anesthetic management for the morbidly obese pregnant woman. *Int Anesthesiol Clin* 2007;356:51-70. 10.1097/AIA.0b013e31802b8a90.
 71. Dresner M, Brocklesby J, Bamber J. Audit of the influence of body mass index on the performance of epidural analgesia in labour and the subsequent mode of delivery. *BJOG* 2006;356:1178-81. 10.1111/j.1471-0528.2006.01048.x.
 72. Wloch C, Wilson J, Lamagni T, Harrington P, Charlett A, Sheridan E. Risk factors for surgical site infection following caesarean section in England: results from a multicentre cohort study. *BJOG* 2012;356:1324-33. 10.1111/j.1471-0528.2012.03452.x,
 73. Catalano PM, Shankar K. Obesity and pregnancy: mechanisms of short term and long term adverse consequences for mother and child. *BMJ*. 2017 Feb 08;356:j1.
 74. Anderson JL, Waller DK, Canfield MA, Shaw GM, Watkins ML, Werler MM. Maternal obesity, gestational diabetes, and central nervous system birth defects. *Epidemiology*. 2005 Jan;16(1):87-92.
 75. Stothard KJ, Tennant PW, Bell R, Rankin J. Maternal overweight and obesity and the risk of congenital anomalies: a systematic review and meta-analysis. *JAMA* 2009;356:636-50. 10.1001/jama.2009.113.
 76. Brite J, Laughon SK, Troendle J, Mills J. Maternal overweight and obesity and risk of congenital heart defects in offspring. *Int J Obes (Lond)* 2014;356:878-82. 10.1038/ijo.2013.244.
 77. Vézina-Im, L.-A. , Nicklas, T. A. , & Baranowski, T. (2018). Intergenerational effects of health issues among women of childbearing age: A review of the recent literature. *Current Nutrition Reports*, 7(4), 274–285. 10.1007/s13668-018-0246-x.
 78. Wojcicki JM. Maternal prepregnancy body mass index and initiation and duration of breastfeeding: a review of the literature. *J Womens Health (Larchmt)* 2011;20(3):341–347.
 79. Pirkola J, Pouta A, Bloigu A, et al. Prepregnancy overweight and gestational diabetes as determinants of subsequent diabetes and hypertension after 20-year follow-up. *J Clin Endocrinol Metab*. 2010;95:772–778.
 80. Walter JR, Perng W, Kleinman KP, Rifas-Shiman SL, Rich-Edwards JW, Oken E. Associations of trimester-specific gestational weight gain with maternal adiposity and

systolic blood pressure at 3 and 7 years postpartum. *Am J Obstet Gynecol*. 2015 Apr;212(4):499.e1-12.

81. Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Rasmussen KM, Yaktine AL, editors. National Academies Press (US); Washington (DC): 2009.
82. Popova S, Lange S, Probst C, Gmel G, Rehm J. Estimation of national, regional, and global prevalence of alcohol use during pregnancy and fetal alcohol syndrome: a systematic review and meta-analysis. *The Lancet Global Health*. 2017;5(3):e290–e9. doi: 10.1016/S2214-109X(17)30021-9.
83. World Health Organization . *Global Status Report on Alcohol and Health*. Geneva: World Health Organization; 2018.
84. Denny CH, et al. Consumption of alcohol beverages and binge drinking among pregnant women aged 18–44 years—United States, 2015–2017. *MMWR Morb Mortal Wkly Rep* 2019;68(16):365–8.
85. Abel EL. Prenatal effects of alcohol. *Drug Alcohol Depend*. 1984;14(1):1–10. doi: 10.1016/0376-8716(84)90012-7.
86. Breslow R.A., Guenther P.M., Smothers B.A. Alcohol drinking patterns and diet quality: The 1999–2000 national health and nutrition examination survey. *Am. J. Epidemiol*. 2006;163:359–366. doi: 10.1093/aje/kwj050.
87. Finer LB, Zolna MR. Unintended pregnancy in the United States: incidence and disparities, 2006. *Contraception*. 2011;84:478–485.
88. West JR, Chen WJ, Pantazis NJ. Fetal alcohol syndrome: the vulnerability of the developing brain and possible mechanisms of damage. *Metab Brain Dis* 1994;9:291–322.
89. American College of Obstetricians and Gynecologists. Committee on Health Care for Underserved Women. Committee opinion no. 496: at-risk drinking and alcohol dependence obstetric and gynecologic implications. *Obstet Gynecol*. 2011; 118(pt 1):383–388.
90. Carson G, Cox LV, Crane J, et al. No. 245-Alcohol use and pregnancy consensus clinical guidelines. *J Obstet Gynaecol Can*. 2017;39:e220–e254.
91. Williams JF, Smith VC. Committee on substance abuse. Fetal alcohol spectrum disorders. *Pediatrics*. 2015;136:e1395–e1406.; Centers for disease control and prevention: fetal alcohol spectrum disorders (FASDs). 2018. Available at: www.cdc.gov/ncbddd/fasd/alcohol-use.html Accessed June 18, 2018.
92. West JR, Chen WJ, Pantazis NJ. Fetal alcohol syndrome: the vulnerability of the developing brain and possible mechanisms of damage. *Metab Brain Dis* 1994;9:291–322.
93. West JR, Chen WJ, Pantazis NJ. Fetal alcohol syndrome: the vulnerability of the developing brain and possible mechanisms of damage. *Metab Brain Dis* 1994;9:291–322.
94. Jeffrey BG, Weisinger HS, Neuringer M, Mitchell DC. The role of docosahexaenoic acid in retinal function. *Lipids* 2001;36:859–71.
95. Henderson J, Gray R, Brocklehurst P. Systematic review of effects of low-moderate prenatal alcohol exposure on pregnancy outcome. *BJOG*. 2007;114:243–252.

96. Muggli E, Matthews H, Penington A, et al. Association between prenatal alcohol exposure and craniofacial shape of children at 12 months of age. *JAMA Pediatr.* 2017;171:771–780.
97. Bertrand J, Floyd LL, Weber MK; Fetal Alcohol Syndrome Prevention Team, Division of Birth Defects and Developmental Disabilities, National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention (CDC). Guidelines for identifying and referring persons with fetal alcohol syndrome [published correction appears in *MMWR Morb Mortal Wkly Rep.* 2006;55(20):568].
98. Young JK, Giesbrecht HE, Eskin MN, Aliani M, Suh M. Nutrition implications for fetal alcohol spectrum disorder. *Adv Nutr.* 2014 Nov 14;5(6):675-92. doi: 10.3945/an.113.004846.
99. Williams JF, Smith VC, Committee on substance abuse. Fetal Alcohol Spectrum Disorders. *Pediatrics.* 2015; 136(5):e1395-406.
100. Hubberstey C., Rutman D. HerWay Home Program for Pregnant and Parenting Women Using Substances: A Brief Social Return on Investment Analysis. *Can. J. Addict.* 2019;11:6–14. doi: 10.1097/CXA.0000000000000086.
101. Ceccanti M, Iannitelli A, Fiore M. Italian Guidelines for the treatment of alcohol dependence. *Riv Psichiatr* 2018;53:105–6. doi:10.1708/2925.29410.
102. Gibson L, Porter M. Drinking or Smoking While Breastfeeding and Later Cognition in Children. *Pediatrics* 2018;142:e20174266. doi:10.1542/peds.2017-4266.
103. WHO, 2014b.
104. WHO. Guidelines for the Identification and Management of Substance Use and Substance Use Disorders in Pregnancy. World Health Organization; Geneva, Switzerland: 2014.
105. Bacon O, Robert S, VandenBerg A. Evaluating nursing satisfaction and utilization of the Clinical Institute Withdrawal Assessment for Alcohol, revised version (CIWA-Ar). *Ment Health Clin.* 2016 Jun;6(3):114-119.
106. Amezcua-Prieto C, Olmedo-Requena R, Jiménez-Mejías E, Hurtado-Sánchez F, Mozas-Moreno J, Lardelli-Claret P, Jiménez-Moleón JJ. Changes in leisure time physical activity during pregnancy compared to the prior year. *Matern Child Health J.* 2013 May;17(4):632-8. doi: 10.1007/s10995-012-1038-3.
107. Aguilar-Cordero M.J., Sánchez-López A.M., Rodríguez-Blancque R., Noack-Segovia J.P., Pozo-Cano M.D., López-Contreras G., Villar N.M. Physical activity by pregnant women and its influence on maternal and foetal parameters; a systematic review. *Nutr. Hosp.* 2014;30:719–726.
108. Poudevigne, M. S. , & O'Connor, P. J. (2006). A review of physical activity patterns in pregnant women and their relationship to psychological health. *Sports Medicine*, 36(1), 19–38. 10.2165/00007256-200636010-00003.
109. Broberg, L. , Ersbøll, A. S. , Backhausen, M. G. , Damm, P. , Tabor, A. , & Hegaard, H. K. (2015). Compliance with national recommendations for exercise during early pregnancy in a Danish cohort. *BMC Pregnancy and Childbirth*, 15(1), 317 10.1186/s12884-015-0756-0.

110. Evenson KR, Wen F. Prevalence and correlates of objectively measured physical activity and sedentary behavior among US pregnant women. *Prev Med* 2011;53:39–43.
111. Pereira M.A., Rifas-Shiman S.L., Kleinman K.P., Rich-Edwards J.W., Peterson K.E., Gillman M.W. Predictors of change in physical activity during and after pregnancy: Project Viva. *Am. J. Prev. Med.* 2007;32:312–319. doi: 10.1016/j.amepre.2006.12.017.
112. Hausenblas H., Giacobbi P., Cook B., Rhodes R., Cruz A. Prospective examination of pregnant and nonpregnant women's physical activity beliefs and behaviours. *J. Reprod. Infant Psychol.* 2011;29:308–319. doi: 10.1080/02646838.2011.629993.
113. Haakstad LA, Bo K. Exercise in pregnant women and birth weight: a randomized controlled trial. *BMC Pregnancy Childbirth.* 2011;11:66 ;Lee I., Djoussé L., Sesso H.D., Wang L., Buring J.E. Physical activity and weight gain prevention. *JAMA.* 2010;303(12):1173–1179.
114. Davenport, M. , Meah, V. , Ruchat, S. , Davies, G. , Skow, R. , Barrowman, N. , ... Mottola, M. F. (2018). Impact of prenatal exercise on neonatal and childhood outcomes: A systematic review and meta-analysis. *British Journal of Sports Medicine*, 52(21), 1386–1396. 10.1136/bjsports-2018-099836.
115. Ruchat, S.-M. , Mottola, M. F. , Skow, R. J. , Nagpal, T. S. , Meah, V. L. , James, M. , ... Davenport, M. H. (2018). Effectiveness of exercise interventions in the prevention of excessive gestational weight gain and postpartum weight retention: A systematic review and meta-analysis. *British Journal of Sports Medicine*, 52(21), 1347–1356. 10.1136/bjsports-2018-099399.
116. Ramos-Leví A.M., Pérez-Ferre N., Fernández M.D. Risk factors for gestational diabetes mellitus in a large population of women living in Spain: implications for preventative strategies. *Int J Endocrinol.* 2012:312529.
117. Deierlein AL, Siega-Riz AM, Evenson KR. Physical activity during pregnancy and the risk of hyperglycemia. *J Womens Health (Larchmt).* 2012;21:769-775.
118. Kasawara KT, do Nascimento SL, Costa ML, Surita FG, e Silva JL. Exercise and physical activity in the prevention of pre-eclampsia: systematic review. *Acta Obstet Gynecol Scand.* 2012;91:1147-1157.
119. Kardel KR, Johanse B, Voldner N, Iversen PO, Henriksen T. Association between aerobic fitness in late pregnancy and duration of labor in nulliparous women. *Acta Obstet Gynecol Scand.* 2009;88:948-952.
120. Sánchez-Polán M, Franco E, Silva-José C, Gil-Ares J, Pérez-Tejero J, Barakat R, Refoyo I. Exercise During Pregnancy and Prenatal Depression: A Systematic Review and Meta-Analysis. *Front Physiol.* 2021 Jun 28;12:640024. doi: 10.3389/fphys.2021.640024.
121. Magro-Malosso ER, Saccone G, Di Mascio D, Di Tommaso M, Berghella V. Exercise during pregnancy and risk of preterm birth in overweight and obese women: a systematic review and meta-analysis of randomized controlled trials. *Acta Obstet Gynecol Scand.* 2017 Mar;96(3):263-273. doi: 10.1111/aogs.13087.
122. Cary GB, Quinn TJ. Exercise and lactation: are they compatible? *Can J Appl Physiol* 2001;26:55–75. 10.1139/h01-004.

123. Rakhshani A, Nagarathna R, Mhaskar R, Mhaskar A, Thomas A, Gunasheela S. The effects of yoga in prevention of pregnancy complications in high-risk pregnancies: A randomized controlled trial. *Prev Med* 2012;55:333–340.
124. Bouya S, Rezaie Keikhaie L, Hosseini S, Rezaie Keikhaie K. The effect of yoga on uterine artery Doppler indices, maternal and fetal complications in pregnant women: A quasi-experimental study. *J Ayurveda Integr Med* 2021;12:70–74.
125. Dhana K, Haines J, Liu G, Zhang C, Wang X, Field AE, Chavarro JE, Sun Q. Association between maternal adherence to healthy lifestyle practices and risk of obesity in offspring: results from two prospective cohort studies of mother-child pairs in the United States. *BMJ*. 2018 Jul 04;362:k2486.
126. Szumilewicz A. Who and how should prescribe and conduct exercise programs for pregnant women? Recommendation based on the European educational standards for pregnancy and postnatal exercise specialists. *Dev Period Med*. 2018;22(2):107-112.
127. Melzer K, Keyser B, Pichard C. Physical activity: the health benefits outweigh the risks. *Curr Opin Clin Nutr Metab Care*. 2004;7:641-647.
128. Mottola MF, Davenport MH, Ruchat SM, et al. Canadian Guideline for Physical Activity throughout Pregnancy. *J Obstet Gynaecol Can*. 2018 Nov;40(11):1528-1537. doi: 10.1016/j.jogc.2018.07.001.
129. American College of Obstetricians and Gynecologists Physical activity and exercise during pregnancy and the postpartum period. Committee Opinion No. 650. *Obstet Gynecol*. 2015;126:e135–e142. doi: 10.1097/AOG.0000000000001214.
130. Di Mascio D, Magro-Malosso ER, Saccone G, et al. . Exercise during pregnancy in normal-weight women and risk of preterm birth: a systematic review and meta-analysis of randomized controlled trials. *Am J Obstet Gynecol* 2016;215:561–71. 10.1016/j.ajog.2016.06.014.
131. Magro-Malosso ER, Saccone G, Di Mascio D, et al. . Exercise during pregnancy and risk of preterm birth in overweight and obese women: a systematic review and meta-analysis of randomized controlled trials. *Acta Obstet Gynecol Scand* 2017;96:263–73. 10.1111/aogs.13087.
132. Perales M, Santos-Lozano A, Ruiz JR, Lucia A, Barakat R. Benefits of aerobic or resistance training during pregnancy on maternal health and perinatal outcomes: A systematic review. *Early Hum Dev*. 2016 Mar;94:43-8. doi: 10.1016/j.earlhumdev.2016.01.004.
133. Barakat R, Perales M. Resistance Exercise in Pregnancy and Outcome. *Clin Obstet Gynecol*. 2016 Sep;59(3):591-9. doi: 10.1097/GRF.0000000000000213.
134. O'Toole ML, Sawicki MA, Artal R. Structured diet and physical activity prevent postpartum weight retention. *J Womens Health* 2003;12:991–8. doi:10.1089/154099903322643910.
135. Davenport MH, McCurdy AP, Mottola MF, et al. Impact of prenatal exercise on both prenatal and postnatal anxiety and depressive symptoms: a systematic review and meta-analysis. *Br J Sports Med* 2018;52:1376–85. doi:10.1136/bjsports-2018-099697.
136. Nygaard I, Girts T, Fultz NH, et al. . Is urinary incontinence a barrier to exercise in women? *Obstet Gynecol* 2005;106:307–14. 10.1097/01.AOG.0000168455.39156.0f.

137. Sancho MF, Pascoal AG, Mota P, et al. Abdominal exercises affect inter-rectus distance in postpartum women: a two-dimensional ultrasound study. *Physiotherapy* 2015;101:286–91. 10.1016/j.physio.2015.04.004.
138. American Academy of Pediatrics, American College of Obstetricians and Gynecologists . Guidelines for perinatal care. 8th ed. Elk Grove Village, IL and Washington, DC: AAP, American College of Obstetricians and Gynecologists, 2017.
139. Mottola MF. Components of exercise prescription and pregnancy. *Clin Obstet Gynecol* 2016;59:552–8. 10.1097/GRF.0000000000000207.
140. Meah VL, Davies GA, Davenport MH. Why can't I exercise during pregnancy? Time to revisit medical 'absolute' and 'relative' contraindications: systematic review of evidence of harm and a call to action. *Br J Sports Med* 2020;54:1395–404. 10.1136/bjsports-2020-102042.
141. Ferrari N, Graf C. [Recommendations for Physical Activity During and After Pregnancy]. *Gesundheitswesen*. 2017 Mar;79(S 01):S36-S39.
142. American College of Obstetricians and Gynecologists. ACOG Committee opinion no. 650: physical activity and exercise during pregnancy and the postpartum period. *Obstet Gynecol* 2015;126:e135–42. 10.1097/AOG.0000000000001214.
143. Mottola MF, Davenport MH, Ruchat S-M, et al. No. 367-2019 Canadian guideline for physical activity throughout pregnancy. *J Obstet Gynaecol Can* 2018;40:1528–37. 10.1016/j.jogc.2018.07.001.
144. Bookari, K., Yeatman, H., & Williamson, M. (2016). Falling short of dietary guidelines—What do Australian pregnant women really know? A cross sectional study. *Women and Birth*, 30(1), 9–17. 10.1016/j.wombi.2016.05.010).
145. Jarman, M., Bell, R. C., Nerenberg, K., & Robson, P. J. (2017). Adherence to Canada's food guide recommendations during pregnancy. *Current Developments in Nutrition*, 1(7), e000356 10.3945/cdn.116.000356.
146. Bookari, K., Yeatman, H., & Williamson, M. (2016). Falling short of dietary guidelines—What do Australian pregnant women really know? A cross sectional study. *Women and Birth*, 30(1), 9–17. 10.1016/j.wombi.2016.05.010.
147. Kominiarek MA, Rajan P. Nutrition Recommendations in Pregnancy and Lactation. *Med Clin North Am*. 2016;100(6):1199-1215. doi:10.1016/j.mcna.2016.06.004.
148. Trumbo P, Yates AA, Poos M. Food and Nutrition Board, Institute of Medicine: Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. National Academies Press; Washington, DC: 2002. SS.
149. Adam, Ishag, Ali, Abdelaziem. "Anemia During Pregnancy". *Nutritional Deficiency*, edited by Pinar Erkekoglu, Belma Kocer-Gumusel, IntechOpen, 2016. 10.5772/63211.
150. Pitkin RM. Folate and neural tube defects. *Am J Clin Nutr*. 2007;85(1):285S–288S.
151. Guidelines for perinatal care. 7th American Academy of Pediatrics and the American College of Obstetricians and Gynecologists; Washington, DC: 2012.

152. EFSA Panel on Dietetic Products, Nutrition and Allergies: Scientific opinion on dietary reference values for water. *EFSA J* 2010;8:1459-1507.
153. Oken E, Ning Y, Rifas-Shiman SL, et al. Diet during pregnancy and risk of preeclampsia or gestational hypertension. *Ann Epidemiol.* 2007;17:663–668. doi: 10.1016/j.annepidem.2007.03.003.
154. Renault KM, Carlsen EM, Hædersdal S et al (2017) Impact of lifestyle intervention for obese women during pregnancy on maternal metabolic and inflammatory markers. *Int J Obes.* 10.1038/ijo.2017.9.
155. van Beynum IM, Kapusta L, Bakker MK, den Heijer M, Blom HJ, de Walle HE: Protective effect of periconceptional folic acid supplements on the risk of congenital heart defects: a registry-based case-control study in the northern Netherlands. *Eur Heart J* 2010;31:464-471.
156. Czeizel AE: Reduction of urinary tract and cardiovascular defects by periconceptional multivitamin supplementation. *Am J Med Genet* 1996;62:179-183.
157. Pereira-da-Silva L, Cabo C, Moreira A, et al. The adjusted effect of maternal body mass index, energy and macronutrient intakes during pregnancy, and gestational weight gain on body composition of full-term neonates. *Am J Perinatol.* 2013;31:875–882. doi: 10.1055/s-0033-1363502.
158. Honzik T, Adamovicova M, Smolka V, Magner M, Hrubá E, Zeman J: Clinical presentation and metabolic consequences in 40 breastfed infants with nutritional vitamin B12 deficiency - what have we learned? *Eur J Paediatr Neurol* 2010;14:488-495.
159. von Schenck U, Bender-Gotze C, Koletzko B: Persistence of neurological damage induced by dietary vitamin B-12 deficiency in infancy. *Arch Dis Child* 1997;77:137-139.
160. Fekete K, Berti C, Trovato M, Lohner S, Dullemeijer C, Souverein OW, et al: Effect of folate intake on health outcomes in pregnancy: a systematic review and meta-analysis on birth weight, placental weight and length of gestation. *Nutr J* 2012;11:75.
161. Grieger JA, Clifton VL, Tuck AR, et al. In utero programming of allergic susceptibility. *Int Arch Allergy Immunol.* 2016;169:80–92. doi: 10.1159/000443961.
162. Thorburn AN, McKenzie CI, Shen S, et al. Evidence that asthma is a developmental origin disease influenced by maternal diet and bacterial metabolites. *Nat Commun.* 2015;6:7320. doi: 10.1038/ncomms8320.
163. Petry N, Olofin I, Boy E et al (2016) The effect of low dose iron and zinc intake on child micronutrient status and development during the first 1000 days of life: a systematic review and meta-analysis. *Nutrients.* 10.3390/nu8120773.
164. Dror DK, Allen LH: Effect of vitamin B12 deficiency on neurodevelopment in infants: current knowledge and possible mechanisms. *Nutr Rev* 2008;66:250-255.
165. Bischoff-Ferrari HA: Vitamin D: role in pregnancy and early childhood. *Ann Nutr Metab* 2011;59:17-21.
166. Dawodu A, Wagner CL: Prevention of vitamin D deficiency in mothers and infants worldwide - a paradigm shift. *Paediatr Int Child Health* 2012;32:3-13.
167. Hyppönen E: Preventing vitamin D deficiency in pregnancy: importance for the mother and child. *Ann Nutr Metab* 2011;59:28-31.

168. Guidelines for perinatal care. 7th American Academy of Pediatrics and the American College of Obstetricians and Gynecologists; Washington, DC: 2012.