

Ethanollic extract of whole unripe plantain *Musa paradisiaca* ameliorates carbon tetrachloride-induced hepatotoxicity and nephrotoxicity in wistar rat

ABSTRACT

Aim: Globally, burden of liver and kidney diseases has been on the increase in recent times. The present study therefore investigates the hepatoprotective and nephroprotective potentials of unripe plantain *Musa paradisiaca* on CCl₄-induced oxidative damage in albino rat. This was with the aim of providing a locally available and potent therapeutic alternative to the conventional drugs used in the management of liver and kidney diseases.

Place and Duration of study: The study was conducted at the Department of Medical Biochemistry, College of Medicine, Ekiti State University, Ado Ekiti between July 2018 and January, 2019.

Methodology: Twenty-five adult male albino rats were placed into seven groups of 5 animals each. Group I animals received distilled water throughout the duration of the experiment, while group II were exposed to CCl₄ only. Groups III, IV, V and VI received 3 ml/kg b.w of CCl₄ intraperitoneally but were post treated with 50 mg/kg and 100 mg/kg of unripe plantain extract respectively while group seven were post-treated with silymarin by oral gavage. Animals were sacrificed for the excision of the liver and kidney. Activities of creatinine kinase (CK), aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), as well as levels of urea, uric acid, bilirubin and lipid profile were assessed. Tissue antioxidant level of reduced glutathione (GSH) and activities of antioxidant enzymes such as superoxide dismutase (SOD) and catalase (CAT) were also determined.

Results: Exposure to CCl₄ caused a significant derangement in lipid profile, resulting in the increase in serum triglyceride, total cholesterol and low density lipoprotein (LDL) while high

density lipoprotein (HDL) level was diminished. Liver and kidney biomarkers (ALT, AST, ALP, CK, urea, uric acid and bilirubin) were also significantly elevated in the serum relative to the control animals following exposure to CCl₄. Activities of antioxidant enzymes in the serum were markedly inhibited by CCl₄ exposure. Treatment with *Musa paradisiaca* extract caused a dose-dependent restoration of all biochemical parameters determined, while histopathological observation was in agreement with biochemical results.

Conclusion: These findings showed that *Musa paradisiaca* extract exhibited positive modulatory effects on the liver and kidney subjected to oxidative attack, hence, its potential usefulness in the management of diseases associated with these organs.

Keywords: biomarkers, carbon tetrachloride, unripe plantain, liver, kidney.

1.0 INTRODUCTION

The liver and kidney are critical to several biochemical processes including metabolism of drugs, hormones, toxins as well as maintenance of cellular homeostasis [1]. The liver is the main metabolic organ in the body and is considered a viable defense system against both environmental and metabolic toxicants [2]. Conversely, the kidneys are mainly involved in urinary excretion of metabolites and osmoregulation. The biological roles of these organs make them vulnerable to oxidative attack by xenobiotics [3]. Several synthetic drugs have been used in the management of liver and kidney diseases. Although, their potency cannot be overemphasized, they are often very expensive, hence they are not affordable by a large percentage of people in developing nations. Moreover, these therapies have been suggested to partially compensate for metabolic derangements seen in diseases, they do not necessarily correct the fundamental biochemical lesions [4]. Considering these limitations, the need for plant-based medicines as alternative therapies becomes critical. Plants are natural and are biologically friendly. They synthesize certain compounds that are primarily designed to protect

them against invasion by both micro and macro-predators. Co-incidentally, these compounds such as polyphenolic substances, melatonin, carotenoids, quercetin, resveratrol, vitamin E, vitamin C, L-carnitine and coenzyme Q10 [5-8] have been found to possess medicinal properties that can be exploited by man in the management of diseases. These phytochemicals, which act as antioxidants protect critical organs and macromolecules from oxidative attack of endogenous and exogenous free radicals [9,10].

Plantain (*Musa paradisiaca*) is cultivated in several tropical countries of the world. It is rich in fiber, iron, serotonin, minerals and vitamins [11,12]. According to an estimate by WHO, about 80% of the population in developing nations of the world recognized the use of plantain for enhancing wound healing [13-18] In fact, unripe plantain pulp has been used as poultice [18, 16] in the management of skin inflammation due to its ability to stimulate angiogenesis by virtue of collagen fibres synthesis and remodeling [19-21]. Recently, reports have recommended plantain meal as dietary approach for the management of diabetes and other diseases related to dysfunctional lipid profile [22,23]. Anticancer and anti-inflammatory potentials of plantain in cell lines and animal models respectively have been documented [24, 15,16]. Folkloric reports have indicated that unripe plantain is helpful in the management of hepatic and renal disorders including anemia [11,12]. Considering the vast medicinal relevance of plantain in traditional parlance, coupled with an ever-increasing burden of liver and kidney diseases, there is a dire need to investigate its curative potential in animal models of hepatotoxicity and nephrotoxicity. Hence, this study.

2.0 MATERIALS AND METHODS

2.1 Collection, preparation and extraction of unripe plantain

Unripe plantain fruit were obtained from a local farm in Ado Ekiti and authenticated at the Department of Plant Science, Ekiti State University, Ado Ekiti, Nigeria.

2.2 Reagents and chemicals

All experimental parameters were determined using diagnostic kits from Rando Chemicals Ltd, England.

2.3 Preparation of extract

Fresh unripe plantains (300 g) were washed with distilled water, weighed, chopped into pieces without peeling and extracted in 80% ethanol for 72 hours to allow for extraction. The supernatant was filtered using Whatman filter paper. The filtrate was then freeze dried, labelled as crude extract and kept in an airtight container in the refrigerator.

2.4 Experimental animals

Twenty-five (25) male wistar rats of average weight 200g were purchased from the animal house of the College of Medicine, Ekiti State University, Ado-Ekiti. All experimental animals were acclimatized for one week and housed in neat metallic cages at standard temperature, relative humidity as well as 12/12-h light and dark cycle. The animals were granted unrestricted access to food and water ad libitum on a daily basis. Rat bedding were routinely changed and replaced for the period of the experiment. The 25 rats were randomly placed into five groups and treated as follows:

2.5 List 1: Animal Treatment

Group	Treatment
I	Animal feed and distilled water only
II	3 ml/kg bw CCl ₄ only
III	3 ml/kg b.w CCl ₄ + 50 mg/kg bw. <i>M. paradisiaca</i>
IV	3 ml/kg b.w CCl ₄ + 100 mg/kg bw. <i>M. paradisiaca</i>
V	3 ml/kg b.w CCl ₄ + 100 mg/kg bw. silymarin

Animals were fasted 24 h before sacrifice on the 15th day of commencement of the work.

2.6 Preparation of organs homogenate

Animals were decapitated under very mild, cold-ether anesthesia and quickly dissected to excise the liver, kidney as well as serum. Ten percent (10%) homogenate each of the liver and kidney were prepared separately in 6.8mM potassium phosphate buffer, (pH 7.4) using chilled pestle

and mortar. The resulting homogenate was centrifuged at 10,000 rpm for 10 min at 4 °C to obtain a supernatant which was stored in a refrigerator and used for the determination of biochemical parameters.

2.7 Preparation of serum

Heart of experimental animals were punctured immediately after decapitation while the animals were still breathing. Blood was collected into plain sample bottles and allow to stand for coagulation. Whole blood (coagulated) was then centrifuged at 3000 rpm for 15 min to obtain the serum which was gently decanted and kept in the refrigerator.

2.8 Determination of biochemical parameters

Creatine kinase was measured following the method of Mattenheimer [25]. Aspartate aminotransferase (AST) and alanine amino transferase (ALT) activity were assayed as described by Reitman and Frankel [26], while and alkaline phosphatase (ALP) was determined according to Englehardt *et al.* [27]. Lipid profile: triglycerides [28], total cholesterol [29], LDL [30] and HDL [31] were assayed according to established protocols. Catalase (CAT) and superoxide dismutase (SOD) were assayed following the methods of Sinha [32] and Misra [33] respectively. Reduced glutathione (GSH) level was determined by the method of Beutler *et al.* [34] while the modified Biuret method described by Weichselbaum [35] was followed for the determination of total protein.

2.9 Statistical analysis

All values are expressed as mean \pm SD. Statistical evaluation was done using One Way Analysis of Variance (ANOVA) followed by Duncan's Multiple Range Test (DMRT) by using SPSS 11.09 for windows. The significance level was set at $p = 0.05$.

3.0 RESULTS

Exposure to CCl₄ resulted in marked (P=0.05) derangement in lipid profile in all tissues of experimental animals analyzed (Table 1). Treatment with *M. paradisiaca* extract caused a dose-dependent restoration of deranged lipid profile in a manner comparable to animals treated with silymarin (Table 1). Serum activities of liver enzyme biomarkers such as alkaline phosphatase (ALP), alanine aminotransferase (ALT) and aspartate aminotransferase (AST) were significantly elevated following exposure to CCl₄ (Table 2) relative to the negative control animals. Post administration of *M. paradisiaca* extract, however, reversed the toxic trend at a level comparable to animals treated with standard drug (Table 2). Serum and tissue levels of kidney biomarkers such as urea, uric acid and bilirubin were significantly (P=0.05) elevated after administering CCl₄, while oral administration of *M. paradisiaca* extract to exposed rats brought about a marked reversal of the toxic trend in a dose-dependent manner (Table 2). Activities of antioxidant enzymes such as superoxide dismutase and catalase as well as creatine kinase were markedly (P=0.05) depleted in animals exposed to CCl₄ toxicity (Table 3), while treatment with *M. paradisiaca* extract caused a dose-dependent restoration of activities in a manner comparable with animals treated with standard drug. On the other hand, serum level of GSH and total protein (Table 3) respectively was significantly decreased following exposure to CCl₄ (Table 3). However, treatment with plantain extract restored the parameters to a level comparable to the negative control.

Table 1. Effect of unripe plantain extract on lipid profile of animals exposed to CCl₄ toxicity

Parameters (mg/dL)	Negative control	Positive control CCl ₄ (3ml/kg bw) only	CCl ₄ + Unripe Plantain (50 ml/kgbw)	CCl ₄ + Unripe Plantain (100 ml/kgbw)	CCl ₄ + Silymarin (100mg/kgbw)
SERUM					
CHOL	73.28±1.84 ^a	140.21±0.00 ^b	105.24±0.56 ^a	88.43±1.25 ^a	77.76±0.96 ^a
TRIG	23.05±1.30 ^a	46.22±1.18 ^b	39.87±0.61	31.46±0.75 ^a	25.35±0.81 ^a
HDL	15.91±0.68 ^a	10.10±0.04 ^b	11.33±0.59	12.88±0.43 ^a	13.35±0.28 ^a
LDL	54.10±0.53 ^a	120.87±0.46 ^b	102.15±1.53 ^a	71.38±1.31 ^a	59.34±1.22 ^a
KIDNEY					
CHOL	23.96±1.80 ^a	38.49±0.65 ^b	35.22±0.59 ^b	27.83±0.43 ^a	24.16±5.89 ^a
TRIG	7.75±0.65 ^a	11.6±0.50 ^b	10.42±0.51 ^b	9.63±0.89 ^a	9.51±0.74 ^a
HDL	6.40±0.07 ^a	3.57±0.11 ^b	4.32±0.42 ^a	4.14±0.61 ^a	5.61±0.25 ^a
LDL	16.01±1.71 ^a	32.6±0.68 ^b	28.56±0.10 ^b	21.12±0.04 ^a	16.64±0.16 ^a
LIVER					
CHOL	72.96±1.01 ^a	119.75±1.52 ^b	103.31±1.23 ^a	97.39±1.20 ^a	65.36±1.19 ^a
TRIG	64.40±0.16 ^a	96.58±0.27 ^b	85.63±0.47 ^a	71.06±0.36 ^a	74.11±0.25 ^a
HDL	53.62±0.11 ^a	34.19±0.10 ^b	40.05±0.83 ^a	49.73±0.43 ^a	54.15±0.19 ^a
LDL	6.53±1.06 ^a	67.44±3.41 ^b	54.64±2.15 ^b	37.62±2.23 ^a	13.10±1.77 ^a

Data represents mean ± SEM values animal experiments performed in triplicate 'a' indicates significant difference (p = 0.05) from the control, (n= 5).

Table 2. Effect of unripe plantain extract on selected biomarkers of liver and kidney injury

Parameters (IU/L)	Positive Control	Negative control CCl ₄ (3ml/kg bw)	CCl ₄ + Unripe Plantain (50 ml/kg bw)	CCl ₄ + Unripe Plantain (100 ml/kg bw)	CCl ₄ + Silymarin (100 mg/kg bw)
SERUM					
ALP	85.37±0.00 ^a	145.96±0.00 ^b	121.78±4.03	98.83±2.45 ^a	96.39±0.00 ^a
ALT	66.09±0.88 ^a	106.69±1.7 ^b	76.92±0.34 ^a	64.36±0.34 ^a	66.92±0.46 ^a
AST	75.19±1.06 ^a	113.01±1.08 ^b	83.30±1.46 ^a	78.84±0.98 ^a	73.36±0.49 ^a
KIDNEY					
ALP	80.17±0.00 ^a	125.96±0.00 ^b	120.37±4.32 ^a	80.63±2.43 ^a	96.39±0.00 ^a
ALT	56.14±0.94 ^a	158.71±7.56 ^b	73.50±0.39 ^a	63.88±0.56 ^a	64.06±0.97 ^a
AST	69.58±1.28 ^a	99.74±1.08 ^b	74.90±0.64 ^a	68.90±1.24 ^a	75.27±1.47 ^a
UREA	52.69±0.67 ^a	93.08±0.00 ^b	57.38±2.18 ^a	54.65±0.63 ^a	58.85±0.00 ^a
URIC	24.36±0.29 ^a	50.48±0.33 ^b	47.00±3.59 ^b	33.45±3.58 ^a	25.55±0.55 ^a
LIVER					
ALP	55.08±0.00 ^a	112.91±0.00 ^b	94.88±5.01 ^a	69.56±2.86 ^a	65.41±1.38 ^a
ALT	44.99±3.23 ^a	116.55±3.18 ^b	84.53±0.36 ^a	73.08±0.45 ^a	66.03±0.96 ^a
AST	69.07±1.55 ^a	104.59±4.32 ^b	84.34±1.05 ^a	73.50±1.44 ^a	68.64±1.74 ^a
T. BIL	24.85±1.28 ^a	46.48±0.18 ^b	41.03±0.47 ^b	32.11±1.04 ^a	26.65±1.15 ^a

Data represents mean ± SEM values animal experiments performed in triplicate 'a' indicates significant difference (p = 0.05) from the control, (n= 5).

Table 3. Effect of unripe plantain extract on selected antioxidant enzymes

Parameters (IU/L)	Positive Control	Negative control CCl ₄ (3ml/kg bw)	CCl ₄ + Unripe Plantain (50 mg/kg bw)	CCl ₄ + Unripe Plantain (100 mg/kg bw)	CCl ₄ + Silymarin (100 mg/kg bw)
SERUM					
SOD	8.47±0.49 ^a	5.02±0.35 ^b	5.73±0.53 ^b	6.91±0.88 ^a	7.80±1.04 ^a
CAT	4.36±0.18 ^a	1.84±0.05 ^b	2.16±0.72 ^a	2.99±0.81 ^a	3.77±0.60 ^a
GSH	6.81±1.10 ^a	4.53±1.22 ^b	5.03±0.95 ^b	6.17±0.66 ^a	6.04±0.87 ^a
TP	3.75±0.20 ^a	1.67±0.60 ^b	2.14±0.57 ^a	2.79±0.27 ^a	2.59±0.18 ^a
KIDNEY					
SOD	5.26±0.02 ^a	2.28±0.12 ^b	3.63±0.27 ^a	3.84±0.36 ^a	3.62±0.21 ^a
CAT	1.36±0.11 ^a	0.64±0.07 ^b	0.95±0.04 ^a	1.39±0.14 ^a	1.47±0.60 ^a
GSH	2.52±0.17 ^a	0.93±0.02 ^b	1.64±0.08 ^a	2.27±0.28 ^a	1.95±0.04 ^a
TP	2.97±0.02 ^a	2.14±0.01 ^b	2.48±0.12 ^a	2.93±0.16 ^a	2.74±0.11 ^a
LIVER					
SOD	2.04±0.10 ^a	1.42±0.42 ^b	1.83±0.32 ^a	1.97±0.19 ^a	2.11±0.32 ^a
CAT	1.14±0.13 ^a	0.26±0.21 ^b	0.74±0.16 ^a	1.04±0.43 ^a	0.97±0.22 ^a
GSH	1.93±0.03 ^a	0.39±0.01 ^b	0.65±0.08 ^a	1.41±0.12 ^a	1.79±0.03 ^a
TP	1.88±0.13 ^a	1.07±0.09 ^b	1.39±0.05 ^a	1.69±0.45 ^a	1.58±0.43 ^a

Data represents mean ± SEM values animal experiments performed in triplicate 'a' indicates significant difference (p = 0.05) from the control, (n= 5).

4.0 DISCUSSION

The therapeutic potential of unripe plantain pulp in the management of hepatic and renal diseases was investigated. Established liver and kidney markers have been used routinely in monitoring the health status of both organs. Specifically, aspartate aminotransferase (AST), alanine aminotransaminase (ALT) and alkaline phosphatase are reliable markers of hepatic disorders [36-38]. Exposure of experimental rats to carbon tetrachloride toxicity resulted in significant elevation in the serum level of these biomarkers relative to animals that were not exposed. Notably, the surge in serum level of these biomarkers implies free-radical induced oxidative injury to the hepatocytes. Increased ALT level indicates hepatocellular damage followed by cardiac tissue damage and is usually accompanied by a rise in AST activity. ALP is a marker of biliary function and cholestasis. However, functional alterations of liver can lead to different pathologies in other organs. Serum level of these enzymes become elevated in cases of liver diseases or damage because they are normally contained inside hepatocyte. They only leak into the blood stream when the liver cells are damaged. The spill-over of the enzyme into the blood is routinely measured as a marker of abnormal cell damage; hence, a markedly raised serum activity indicates severe liver disease [39]. These biochemical parameters were on the increase as a result of tetrachloromethane (CCl₄)-induced oxidative attack on the liver cells. This might be due to deterioration and necrosis of the liver cells by the oxidative attack of trichloromethyl radicals resulting in the release of transaminases into the blood stream [40-41]. This observation is in line with the earlier reports of [42-45] where liver biomarkers of experimental animals were increased following exposure to pesticides. Treatment of exposed animals with unripe plantain extract caused a significant, dose- dependent restoration of AST, ALT and ALP activities in the serum and tissue homogenates relative to exposed animals that were not treated with the extract. These findings are in agreement with Chinedum and Polycarp [46], and suggests the

hepatoprotective potentials of unripe plantain extract which can be exploited in the management of liver diseases.

Lipid profile (low density lipoprotein cholesterol (LDL-C), high density lipoprotein cholesterol (HDL-C), total cholesterol (TC) and triacylglycerol (TG) could provide useful information on the predisposition of the heart to atherosclerosis and its associated coronary heart disease (Yakubu *et al.*, 2008). The significant reduction in triacylglycerol of exposed but untreated animals may be associated with impaired lipolysis while reduction in HDL-C at all doses investigated may not be clinically beneficial to the animals since the rate at which plasma cholesterol are carried to the liver were also decreased. Furthermore, the enhanced level of cholesterol and LDL-C may suggest cardiovascular risk in the animals. This is supported in the present study by the increase in the computed atherogenic index, a useful indicator of cardiovascular diseases [47]. The unripe plantain extract was able to restore the level of triacylglycerol and carp up the level of HDL-C compared to the control (Table1). **This observation is in agreement with earlier reports [48-50]** and suggests the potential usefulness of plantain in the management of cardiovascular diseases.

Superoxide dismutase (SOD) and catalase (CAT) are the two major radical scavenging enzymes. Superoxide dismutase is the main enzymatic defense against the superoxide anion. It detoxifies the superoxide anion, thus converting it into hydrogen peroxide and water. Although SOD is an antioxidant enzyme, some studies have suggested that it's over expression is actually harmful to cells [51]. The toxic effect of reactive oxygen species observed in many cells with over expressed SOD has been linked to elevated levels of hydrogen peroxide (H₂O₂) and accompanying oxidative damage following hydroxyl radical formation [52]. In the present study, treatment of CCl₄-exposed animals with *M. parasidiaca* extract reactivated the activity of SOD in the serum. This is in agreement with Friday and Chinedu [53].

Catalase is a heme protein that catalyzes the reduction of hydrogen peroxides and protects tissues from hydroxyl radicals. Consequently, the decrease in activities of SOD and CAT as well as GSH level in both liver and kidney during disease condition may be due to over-production of reactive oxygen species in animals [54]. The overproduction of free radicals in turn, causes oxidative damages to membrane's lipid and protein, and ultimately leads to a decrease in the content of GSH and activity of its dependent enzyme. However, treatment of exposed animals with unripe plantain extract ameliorated the activities of these antioxidant enzymes as well as GSH (Table 3). **This is in agreement with Ji *et al.* [55].**

In the present study, integrity of the kidney was assessed through serum urea and uric acid levels. All animals exposed to CCl₄ showed significantly increased level of serum urea and uric acid relative to unexposed animals. This suggests the nephrotoxic potential of CCl₄ (Table 2). However, administration of unripe plantain extract restored the serum urea and uric acid levels to values comparable with unexposed animals. This observation is in line with the report of Friday and Chinedu [53]. This is an indication that unripe plantain is a potential nephroprotective agent that can be exploited in the management of kidney diseases.

In agreement with previous research [48-50], unripe plantain restored deranged lipid profile, remedied distorted liver and kidney biomarkers as well as reactivated the activities of antioxidant enzymes. Hence, it is a potential plant with hepatoprotective and nephroprotective properties that can be exploited in the management of liver and kidney diseases.

Ethical Approval

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

Acknowledgement

Authors acknowledge the support of Mr Oyelade of the Federal Polytechnic Ado Ekiti, Nigeria.

Authors Contribution

Author 1 designed the work, analyzed the data and wrote the final draft. Authors 2 did literature search and wrote the initial draft of the manuscript.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

Hilly M, Adams ML, Nelson SC. A study of digit fusion in the mouse embryo. *Clin Exp Allergy*. 2002;32(4):489-98.

[1] Thuawaini MM, Mawahib BG, Abbas KF. Hepatoprotective and Nephroprotective Effects of the Aqueous Extract of Turmeric (*Curcuma Longa*) in Rifampicin and Isoniazid-Induced Hepatotoxicity and Nephrotoxicity in Rats. *Asian Journal of Pharmaceutical and clinical Research*, 2019;12 (3)2455-3891.

[2] Manfo FPT, Nantia EA, Kuete V. Hepatotoxicity and hepatoprotective effects of African medicinal plants. *Toxicological survey of African medicinal plants*. V. Kuete, Ed. 2014; 323–355.

[3] Brzówska MM, Moniuszko-Jakoniuk J, Piłat-Marcinkiewicz B, Sawicki B. Liver and kidney function and histology in rats exposed to cadmium and ethanol. *Alcohol and Alcoholism*. 2003;38 (1):2–10.

[4]. Chatila R, West AB. Hepatomegaly and abnormal liver tests due to glycogenesis in adults with diabetes. *Med Balt* 1996;75: 327–333.

[5] Wang Y, Wu Y, Luo K, Liu Y, Zhou M, Yans S, Shi H, Cai Y. The protective effects of selenium on cadmium-induced oxidative stress and apoptosis via mitochondria pathway in mice Sidney. *Food Chem. Toxicol*. 2013;58:61–67.

[6] Ibrahim TA, Tarboush HA, Alada A, Mohanna MA. The effect of selenium and lycopene on oxidative stress in bone tissue in rats exposed to cadmium. *Food Nutr. Sci*.2014; 5:1420–1429.

[7] El-Boshy ME, Risha EF, Abdelhamid FM, Mubarak MS, Hadda, TB. Protective effects of selenium against cadmium induced hematological disturbances, immunosuppressive, oxidative stress and hepatorenal damage in rats. *Trace Elem. Med. Biol*. 2015;29:104–110.

[8] El-Sayed YS, El-Gazzar AM, El-Nahas AF, Ashry KM. Vitamin C modulates cadmium-induced hepatic antioxidants gene transcripts and toxicopathic changes in Nile tilapia, *Oreochromis niloticus*. *Environ. Sci. Pollut. Res*. 2016;23:1664–1670.

[9] Lakshmi BVS, Sudhakar M, Aparna M. Protective effect of black grapes on cadmium induced hepatotoxicity in rats. *World Pharm. Sci*. 2014; 2: 276–282.

[10] Baiomy AA, Mansour AA. Genetic and histopathological responses to cadmium toxicity in rabbit's kidney and liver: Protection by ginger (*Zingiberocinale*). *Biol. Trace Elem. Res*.2016; 170: 320–329.

[11] Iweala EJ, Obichi IC, Omotosho IE. Biochemical and histological responses of hepatotoxic rats fed *M. paradisiaca* supplemented diet. *Int J Pharmacol* 2011;7(4): 471–477.

[12] Jimmy EO, Okon MA. Periodic validation of high anti-diabetic potentials of unripe plantain in comparison with glibenclamide and Fansidar. *Am J Pharmacol Toxicol* 2012;7(1): 15–18.

[13] Asuquo EG, Udobi CE. Antibacterial and toxicity studies of the ethanol extract of *Musa paradisiaca* leaf. *Cogent Biol* 2016;2:1219248.

[14] Amutha K, Selvakumari U. Wound healing activity of methanolic stem extract of *Musa paradisiaca* Linn. (Banana) in Wistar albino rats. *Int Wound J* 2016;13:763–767.

- [15] Krishnan SSC, Subramanian IP, Subramanian SP. Isolation, characterization of syringin, phenylpropanoid glycoside from *Musa paradisiaca* tepal extract and evaluation of its antidiabetic effect in streptozotocin-induced diabetic rats. *Biomed Prev Nutr* 2014; 4:105–111.
- [16] Pereira A, Maraschi M. Banana (*Musa* spp) from peel to pulp: ethnopharmacology, source of bioactive compounds and its relevance for human health. *J Ethnopharmacol* 2015;160:149–163. <https://doi.org/10.1016/j.jep.2014.11.008>
- [17] Shodehinde SA, Oboh G. Antioxidant properties of aqueous extracts of unripe *Musa paradisiaca* on sodium nitroprusside induced lipid peroxidation in rat pancreas in vitro. *Asian Pac J Trop Biomed* 2013;3:449–457. [https://doi.org/10.1016/S2221-1691\(13\)60095-7](https://doi.org/10.1016/S2221-1691(13)60095-7)
- [18] Agyare C, Asase A, Lechtenberg M, Niehues M, Deters A, Hensel A. An ethnopharmacological survey and in vitro confirmation of ethnopharmacological use of medicinal plants used for wound healing in Bosomtwi-Atwima-Kwanwoma area. *Ghana J Ethnopharmacol* 2009;125:393–403. <https://doi.org/10.1016/j.jep.2009.07.024>
- [19] Joshi A, Sengar N, Prasad SK, Goel RK, Singh A, Hemalatha S. Wound-healing potential of the root extract of *Albizia lebbek*. *Planta Med* 2013;79:737–743. <https://doi.org/10.1055/s-0032-1328539>.
- [20] Murthy S, Gautam M, Goel S, Purohit V, Sharma H, Goel R. Evaluation of in vivo wound healing activity of *Bacopa monniera* on different wound model in rats. *Biomed Res Int* 2013;972028
- [21] Patil MVK, Kandhare AD, Bhise SD. Pharmacological evaluation of ethanolic extract of *Daucus carota* Linn root formulated cream on wound healing using excision and incision wound model. *Asian Pac J Trop Biomed* 2012;2:S646–S655
- [22] Ajiboye BO, Oloyede HOB, Salawu MO. Antihyperglycemic and antidyslipidemic activity of *Musa paradisiaca*-based diet in alloxan-induced diabetic rats. *Food Sci Nutr*. 2018; 6:137–145. <https://doi.org/10.1002/fsn3.538>
- [23] Oluwajuyitan TD, Ijarotimi OS. Nutritional, antioxidant, glycaemic index and antihyperglycaemic properties of improved traditional plantain-based (*Musa AAB*) dough meal enriched with tigernut (*Cyperus esculentus*) and defatted soybean (*Glycine max*) flour for diabetic patients. *Heliyon* 2019;5:e01504. <https://doi.org/10.1016/j.heliyon.2019.e01504>
- [24] Correa M, Mesomo MC, Pianoski KE, Torres YR, Corazza ML. Extraction of inflorescences of *Musa paradisiaca* L. using supercritical CO₂ and compressed propane. *J Supercrit Fluids* 2016;113:128–135
- [25] Mattenheimer H. Urinary enzyme measurements in the diagnosis of renal disorders. *Ann Clin Lab Sci*. 1981;11(3):189–201.
- [26] Reitman S, Frankel S. Glutamic – pyruvate transaminase assay by colorimetric method. *Am J Clin Pathol*. 1957;28:56–60.
- [27] Englehardt A. Measurement of alkaline phosphatase. *Aerztl Labor*. 1970;16:42.
- [28] Trinder H. A simple Turbidimetric method for the determination of serum cholesterol. *Ann Clin Biochem*. 1969;6:165.
- [29] Tietz NW. *Clinical guide to laboratory tests*. 3rd ed. Philadelphia: Saunders and Co. Publishers; 1995.
- [30] Grove TH. Effect of reagent pH on determination of high-density lipoprotein cholesterol by precipitation with sodium phosphotungstate-magnesium. *Clin Chem*. 1979; 25(4):560–4.
- [31] Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499–502.
- [32] Sinha AK. Colorimetric assay of catalase. *Anal Biochem*. 1972;47:389–94.

- [33] Misra HP. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem.* 1972;247(15):3170–5.
- [34] Beutler E, Duron O, Kelly BM. Improved method for the determination of blood glutathione. *J Lab Clin Med.* 1963;61:882–90.
- [35] Weichselbaum TE. An accurate and rapid method for the determination of protein in small amount of blood, serum. *Am J Clin Pathol.* 1995;16:40.
- [36] Ayepola OR, Chegou NN, Brooks NL, Oguntibeju OO. Kolaviron, a Garcinia biflavonoid complex ameliorates hyperglycemia-mediated hepatic injury in rats via suppression of inflammatory responses *BMC Complement. Altern. Med.* 2013;13p. 363.
- [37] Mossa A, Refaie A, Ramadan A, Bouajila Jalloul. Amelioration of Prallethrin-Induced Oxidative Stress and Hepatotoxicity in Rat by the Administration of *Origanum majorana* Essential Oil. *BioMed Research International* 2013 ;(2):85908.
- [38] Kim EE, Wyckoff HW. Reaction mechanism of alkaline phosphatase based on crystal structures. Two-metal ion catalysis. *J. Mol. Biol.*, 1991;218: 449-464.
- [39] Champe, PC, Havey RA. Cholesterol metabolism. In: Lippicott's illustrated reviews biochemistry. 2nd edition. Champe PC, HaveyRA.(eds). JB Lippincott, Kendallville, pp. 210-225 1994.
- [40] Airaodion AI, Ogbuagu EO, Ogbuagu U, Adeniji AR, Agunbiade AP, Airaodion EO. Hepatoprotective effect of *Parkiabiglobosa* on acute ethanol-induced oxidative stress in Wistar Rats. *International Research Journal of Gastroenterology and Hepatology.* 2019a ;4(6):1-10.
- [41] Airaodion AI, Akinmolayan JD, Ogbuagu EO, Esonu CE, Ogbuagu U. Preventive and therapeutic activities of methanolic extract of *Talinum triangulare* leaves against ethanol-induced oxidative stress in Wistar rats. *International Journal of Bio-Science and Bio-Technology.* 2019b; 11 (7): 85-96.
- [42] Shahzad M, Atif Y, Mehmood S, Muhammad KF, Nazish MA, Tanveer H, Muhammad S, Nuzhat Fatima. Effects of Exposure to Pesticides on Blood Serum Components and Butyrylcholinesterase (BChE) in Pesticide Vendors of Punjab Province, Pakistan. *Annals of King Edward Medical University Lahore Pakistan* 2016;22 (4). DOI: 10.21649/akemu.v22i4.1460
- [43] Araoud M, Neffeti F, Douki W, Hfaiedh H B, Akrouit M, Hassine M. Adverse effects of pesticides on biochemical and haematological parameters in Tunisian agricultural workers. *J. Expos. Scie. Environ. Epidem.* 2012; 22 (3): 243-247.
- [44] Khan DA, Bhatti MM, Khan FA. Adverse effects of pesticides residues on biochemical markers in Pakistani tobacco farmers. *International Journal of Clinical and Experimental Medicine.* 2008;1(3):274–282.
- [45] Jamal F, Haque SQ, Singh S. The Influence of Pesticides on Hepatic and Renal Functions in Occupational Sprayers of Rural Malihabad, Lucknow (India). *Toxicol open access* 2016;1:106. doi:10.4172/2476-2067.1000106.
- [46] Chinedum OE, Polycarp O. Use of unripe plantain (*Musa paradisiaca*) in the management of diabetes and hepatic dysfunction in streptozotocin induced diabetes in rats *Interventional Medicine and Applied Science* 2015;7 (1):9–16.
- [47] Panagiotakos B, Pitsavos C, Skoumas J, Chrysohoou, C. Toutouza M. Stefanadis, C.I. Importance of LDL/HDL ratio as a predictor for coronary heart disease events in patients with heterozygous familial hypercholesterolemia: A 15 year follow-up (1987-2002). *Curr Med Res Opin.* 2003;19:89–94.
- [48] Mallick CR, Maiti. D. Comparative study on antihyperglycemic and antihyperlipidemic effect of separate and composite extract of seed of *Eugenia jambolana* and root of *Musa*

paradisiaca in streptozotocin induced diabetic male albino rat. *Iran. J. Pharmacol. Ther.* 2006;5: 27-33.

[49] Parmar HS, Kar A. Medicinal values of fruit peels from *Citrus sinensis*, *Punica granatum*, and *Musa paradisiaca* with respect to alterations in tissue lipid peroxidation and serum concentration of glucose, insulin, and thyroid hormones. *J Med Food* 2008;11:376–381. <https://doi.org/10.1089/jmf.2006.010>

[50] Vijayakumar S et al (2017) Therapeutic effects of gold nanoparticles synthesized using *Musa paradisiaca* peel extract against multiple antibiotic resistant *Enterococcus faecalis* biofilms and human lung cancer cells (A549). *Microb Pathog* 102:173–183. <https://doi.org/10.1016/j.micpath.2016.11.029>

[51] Gardner R, Salvador A, Moradas-Ferreira P. Why does SOD overexpression sometimes enhance, sometimes decrease, hydrogen peroxide production? A minimalist explanation. *Free Radic. Biol. Med.* 2002; 32, 1351-1357.

[52] de Haan JB, Bladier C, Griffiths P, Kelner M, O'Shea RD, Cheung NS, Bronson RT, Silvestro MJ, Wild S, Zheng SS, Beart PM, Hertzog PJ, Kola I. Mice with a homozygous null mutation for the most abundant glutathione peroxidase, Gpx1, show increased susceptibility to the oxidative stress-inducing agents paraquat and hydrogen peroxide. *J Biol Chem* 1998;273:22528–22536

[53] Friday U, Chinedu I. Hypoglycemic, Hypolipidemic and Antioxidant Properties of *Musa paradisiaca*, Normalis (Plantain)- supplemented diet on Alloxan-induced diabetic albino rat. *Asian J. Biochem.*, 2016;11 (3): 162-167.

[54] Kaleem M, Kirmani D, Asifa M, Ahmedb Q. and Bilqees B. (2006). Biochemical effects of *Nigella sativa* L seeds in diabetic rats. *Indian Journal of Experimental Biology* (44): 745-748.

[55] Ji Xi, Zhang P, Armstrong NR, Gilliland LG. The three-dimensional structure of a glutathione S-transferase from the Mu gene class. Structural analysis of the binary complex of isoenzyme 3-3 and glutathione at 2.2-Å resolution. *Biochemistry* 1992;31(42): 10169–10184.