

# A preliminary study on allelopathy and potential allelochemicals of root exudates from *Solanum rostratum* Dunal.

**Abstract:**

**Aims:** In order to understand the role of allelopathy in invasion success, effects of root exudates from *Solanum rostratum* Dunal on germination and primary growth of wheat and cabbage seeds were studied by vitro bioassay. These results indicated the allelopathy of root exudates and would be helpful for understanding *Solanum rostratum* Dunal invasion .

**Study Design:** The roots were washed clean for three times with distilled water before they were gathered in a big bucket. The container was filled with distilled water, enough to submerge the roots, and air was aerated constantly with an air pump. The water extract was collected per 24hs and for four times totally. The water extract from *Solanum rostratum* was concentrated with a rotary evaporator at 40°C and deposited in a refrigerator at 4°C. All the root exudates were pooled around 1500mL, and the fresh weight of *Solanum rostratum* used for exudates collection was 1208g, the exudate concentration was then marked as 0.8g f w/mL.

**Place and Duration of Study:** *Solanum rostratum* was grown in the greenhouse on April 15th, 2019, its seeds were collected from the invaded fields in Chaoyang city (Liaoning province, China)

**Methodology:** Petri dish bioassay was applied to test the effects of root exudates of *Solanum rostratum* on the seeds of wheat and Chinese cabbage. For test of allelochemicals exudated through the root, The compounds in organic fraction of root exudates analysed by GC-MS.

**Results:** These results indicated the allelopathy of root exudates and would be helpful for understanding *Solanum rostratum* Dunal invasion .

**Conclusion:** Root exudates of *Solanum rostratum* contained some allelochemicals, which could inhibit the germination and radicle growth of wheat and Chinese cabbage, though the effects of root exudates on shoot growth are different, with a stimulation on Chinese cabbage while a inhibition on wheat. The difference indicated a selectivity of allelopathy effect of root exudates from *Solanum rostratum* Dunal.

**Key words:** *Solanum rostratum* Dunal; root exudates; Allelopathy

## 1. INTRODUCTION

Invasion of exotic species pose a great economic loss<sup>[1]</sup>, decreases in the abundances of native species and destroy composition and function of ecosystems in invaded regions<sup>[2-4]</sup> (Wilcove et al. 1998; Memmott et al. 2000; Grigulis et al. 2001; Wilcove et al. 1998). In order to find a solution to weed problems, an increasing body of research has been focused on the mechanism understanding invasion success,

Allelopathy refers to a direct positive or negative effects of a plant on another plant or on micro-organisms by allelochemicals released into the environment<sup>[5]</sup> (Bais, 2003), and has been reported to be one of main mechanisms for plant invasion<sup>[6-7]</sup> (Callaway, 2000; Ping YF, 2012). In “novel weapons hypothesis”, the allelochemicals, produced and released by invasive species, may contribute to their successful invasion<sup>[8-10]</sup> (Callaway, 2008; Vivanco, 2004; Liu C, 2019). The allelochemicals released by invasive species, which are noxious to native plants, can reduce the growth and even cause the death of susceptible native species, thus reducing competition and

increasing resource availability for the alien species. Invasive plants may gain an advantage over their competitors<sup>[11-14]</sup> (Bertin et al, 2003; Weir et al, Zhao XH, 2004; Tian XF, 2017).

*Solanum rostratum* Dunal (Buffalobur), native to North America and belong to the Solanaceae family, is an annual spiny weed (Whalen, 1979; Bassett and Munro, 1986). As an invasive weed, it has extended from central Mexico northward across the Great U.S.A. to some other countries of the world, including U.S.S.R., Australia, Canada and Korea<sup>[15]</sup> (Cho and Kim, 1997; Whalen, 1979b). In China, *Solanum rostratum* was first found in Chaoyang, Liaoning Province in 1981<sup>[16]</sup> (Guan et al. 1984), and was reported capable of invading into roadsides, riverbanks, wasted and cultivated fields in Jilin, Beijing, Hebei, Shanxi, and Xinjiang<sup>[17]</sup> (Wei et al. 2007). It has been reported that leaves, berries, and roots of the invasive species contain some noxious substances, which could poison the livestock to death<sup>[18]</sup> (Bab et al, 2004).

This research preliminarily studied the effects of root exudates from *Solanum rostratum* on effect of cabbage and wheat seeds.

## 2 MATERIALS AND METHODS

### 2.1 Collection of root exudates of *Solanum rostratum* Dunal

*Solanum rostratum* was grown in the greenhouse on April 15<sup>th</sup>, 2010, its seeds were collected from the invaded fields in Chaoyang city (Liaoning province, China). The plants were dug out and collected when they were at the bud stage.

The roots of 50 healthy plants with uniform growth, which fresh weight were 1208g, were washed clean for three times with distilled water before they were gathered in a big bucket. The container was filled with distilled water, enough to submerge the roots, and air was aerated constantly with an air pump. The water extract was collected per 24hs and for four times totally<sup>[19]</sup>. Finally, 1500 ml of root exudate extract was obtained, whichThe water extract from *Solanum rostratum* was concentrated with a rotary evaporator at 40°C and deposited in a refrigerator at 4°C. All the root exudates were pooled around 1500mL, and the fresh weight of *Solanum rostratum* used for exudates collection was 1208g, the exudate concentration was then marked as 0.8g f w/mL.

### 2.2 Effects of root exudates on seeds germination and growth

Petri dish bioassay was applied to test the effects of root exudates of *Solanum rostratum* on the seeds seed germination of wheat (*Triticum aestivum* L.) and Chinese cabbage (*Brassica campestris* L.). Wheat and cabbage seeds were purchased from Shenyang Agricultural University seed company. Prepare 4 extracts of root exudates with different concentrations: control group (CK): 0 g f w/mL (distilled water); I: 0.1 g f w/mL (27.5 mL distilled water plus 3.5 mL root exudates); II: 0.2g f w/mL (22.5 mL distilled water plus 7.5 root exudates); III: 0.4 g f w/mL (27.5 mL distilled water plus 3.5 root exudates). The root exudates of four concentrations were sterilized with microporous filter membrane (0.2μm), mixed with distilled water containing agar, which were used for seed germination after solidification, and added to a culture dish with a diameter of 15cm.

Sterilized Petri dishes (15 cm diameter) were filled with root exudates, which were diluted by distilled water into 4 different concentrations: control group (CK): 0 g f w/mL (distilled water); I: 0.1 g f w/mL (27.5 mL distilled water plus 3.5 mL root exudates); II: 0.2g f w/mL (22.5 mL distilled water plus 7.5 root exudates); III: 0.4 g f w/mL (27.5 mL distilled water plus 3.5 root

exudates). Before the mixture, the root exudates were filtered by millipore filter (0.2μm) and the distilled water was mixed with appropriate weight of agar, which was preheated and liquid. The dish was placed in a 30 degree angle with the tabletop, and the seeds of 20 Chinese cabbage after sterilization were placed in a straight line 1/3 in the culture dish, as well as 15 wheat seeds.

Bioassays of each concentration of extracts were conducted with 30 wheat or cabbage seeds equidistantly set in each glass Petri dishes. There were three replicates for treatment of each concentration, and the replicates were kept at room temperature and shaded against the light. Germination rates were recorded daily until no further seeds germinated for three days. Then the length of the radicle and shoot were measured with a ruler. A seed was considered as having germinated when the seed coat was broken and the radicle had emerged.

RI (response index) was determined according to Williamson (1988), as followed:

$$RI=1-C/T \ (T>C) \text{ or } RI=T/C-1 \ (T<C),$$

$$T=\text{Treatment response}, C=\text{Control response}$$

RI ranges from -1 to + 1, with positive values indicating stimulation by the treatments and negative values indicating inhibition, relative to the controls. The absolute values indicate the degrees of effects.

### 2.3 Potential allelochemicals in root exudates tested by GC-MS

Refer to Friebe's method and make some changes<sup>[20]</sup>, for test of allelochemicals exudated through the root, 200 mL root exudates was concentrated to about 50 mL, and was successively extracted with four organic solvents: Chloroform, ethyl acetate, acetone and ethyl ether, each 30mL for three times(totally 90 mL separately). The organic phase was collected as sample and concentrated to around 5mL with rotary evaporator.

The sample was analyzed by Aglient 5973N (GC-MS instrument), with a column HP-5 MS (30 m x 250 mm; film thickness=0. 25 μm). The column temperature was maintained at 50°C for 1 min, and then raised to 280°C (at the rate of 10°C per min) and maintained for 15mins.; electrode stem temperature was set at 150°C and ion source temperature at 230°C; The injector temperature was set to 250°C; E I was set at 70eV. a full scan (35-520m/z). 1 μL sample was injected by using a splitless injector, using helium as a carrier gas at a linear flow rate of 1 mL/min, and analyzed with a range of full scan (35-520m/z).

### 2.4 Statistical analysis

Probit analysis was conducted with Dps 2000 software. Effects of the extracts root exudates were tested through one-way ANOVA and tested by LSD at P<0.05. Significant difference (P<0.05) was marked with different letters in figures

## 3. RESULTS

### 3.1 Germination rates of wheat and Chinese cabbage

Germination rates of wheat and cabbage were overall decreased by the treatments of different concentrations of root exudates from *Solanum rostratum*, and the inhibitory effects became stronger as the concentration increased. Germination rates of wheat were: CK: 62.97%; I:51.83%; II:42.53%; III: 44.4%, and the inhibitory effect was significant (P<0.05) when the concentration

was above 0.1 g f w/ml (fig.1). Germination rates of Chinese cabbage are: CK:50%; I:41.67%; II:43.33%; III: 26.67%, and the germination rate was decreased significant ( $P < 0.05$ ) when the concentration was above 0.2 g f w/ml (fig.2).




Fig.1 Germination rates of wheat (*Triticum aestivum* L.) treated by root exudates of *Solanum rostratum* Dunal.

Note: Different lowercase letters indicate significant differences among species at  $P < 0.05$  according to ANOVAs (Tukey HSD test). The same as below.




Fig. 2 Germination rates of Chinese cabbage (*Brassica campestris* L.) treated by root exudates of *Solanum rostratum*

### 3.2 Radicle length of wheat and Chinese cabbage

Totally, radicle growth of wheat and cabbage were inhibited by various concentrations of root exudates from *Solanum rostratum*, and the inhibitory effects became greater as the concentration increased. Radicle length (cm) of wheat were as follows: 5.24(CK), 4.27(I), 4.29(II) and 1.62 (III), and the inhibitory effect was significant ( $P < 0.05$ ) when the concentration reached as high as 0.1 g f w/ml (fig.3). Radicle length (cm) of Chinese cabbage were as follows: 5.71(CK), 4.76(I), 3.41(II) and 2.64(III), and the radicle growth was significantly inhibited ( $P < 0.05$ ) when the concentration was above 0.2 g f w/ml (fig.4).




Fig.3 Radicle length of wheat treated by root exudates of *Solanum rostratum* Dunal.




Fig.4 Radicle length of Chinese cabbage treated by root exudates of *Solanum rostratum*

### 3.3 Shoot length of wheat and Chinese cabbage

Indicated by the results, effects of root exudates on shoot growth of wheat and cabbage came out to be different. The shoot growth of wheat was inhibited while the shoot growth of Chinese cabbage was stimulated. Shoot length (cm) of wheat were as follows: 3.11 (CK), 2.49 (I), 2.54 (II) and 1.75 (III), and the inhibitory effect was significant ( $P < 0.05$ ) when the concentration reached as high as 0.1 g f w/ml and the effects strengthened as concentration increased (fig.5). Shoot length (cm) of Chinese cabbage were as follows: 2.67 (CK), 2.99 (I), 2.91 (II) and 3.33 (III), and the radicle growth was stimulated by root exudates, but the effects was negatively significant ( $P > 0.05$ ) (fig.6).




Fig.5 Shoot length of wheat treated by root exudates of *Solanum rostratum* Dunal.




Fig.6 Shoot length of Chinese cabbage treated by root exudates of *Solanum rostratum* Dunal.

### 3.4 Response index of wheat and Chinese cabbage

RI values of wheat were negative, which indicates that root exudates inhibited the germination and growth of radicle and shoot. When the concentration was as high as 0.4g fw/mL, obvious difference of absolute RI values occurred in sequence: RI (radicle length) > RI (shoot length) > RI (germination rates) (fig.7). Unlikely to wheat, RI value (shoot length) of Chinese cabbage was positive, while the RI values of germination rates and radicle length are negative. The RI values manifested that the growth of shoot of cabbage was stimulated but the germination and radicle growth were inhibited by root exudates. And the inhibitory effect on radicle growth was stronger, suggested by absolute RI value (fig.8).




Fig.7 Response index of root exudates from *Solanum rostratum* Dunal on wheat




Fig.8 Response index of root exudates from *Solanum rostratum* Dunal on Chinese cabbage

### 3.5 Analysis of allelochemicals by GC-MS

In organic fraction of root exudates, 23 different compounds were detected, including 14 sorts of saturated and in saturated alkane, 7 esters, 1 amine and 1 phenols. From all the compounds were selected 7 esters, 1 amine and 1 phenol, which were potential allelochemicals, and their retention time were marked in total ion flow graph of GC-MS (Fig.9). The chemical names of the potential allelochemicals were listed in table 1.




Fig.9 Total ion flow graph of organic fraction of root exudates by GC-MS

Tab.1 Potential allelochemicals in root exudates of *Solanum rostratum* Dunal

<i>Chemical names of potential allelochemicals</i>	<i>Retention Time (min)</i>	<i>Peak Area (%)</i>	<b>Molecular Formular</b>
1-Butanol,3-methyl-, acetate	3.715	53.18	83
1-Butanol,2-methyl-, acetate	4.048	7.82	90
Thenylidiamine	14.698	3.58	86
1,2-Benzenedicarboxylic acid, bis (2-methylpropyl) ester	16.509	1.66	83
Pentadecanoic acid, 14-methyl-, methyl ester	17.076	0.26	83
Eptadecanoic acid,16-methyl-, methyl ester	18.920	0.13	95

Phenol,2,2'-methylene bis[6-(1,1-dimethylethyl)-4-methyl-	21.537	0.11	98
<b>Bis(2-ethylhexyl)phthalate</b>	22.487	0.15	82

#### 4. DISCUSSION

Our research revealed that root exudates of *Solanum rostratum* contained some allelochemicals, which could inhibit the germination and radicle growth of wheat and Chinese cabbage, though the effects of root exudates on shoot growth are different, with a stimulation on Chinese cabbage while a inhibition on wheat. The difference indicated a selectivity of allelopathy effect of root exudates from *Solanum rostratum* Dunal. In the natural world, the allelochemicals released into the soil could inhibit the seed germination and growth of the other species, and reduce the ability of gaining resources of other plants<sup>[21-23]</sup> (Molly et al, 2002; Rose, 1972; Weiner et al, 1997).

The compounds in organic fraction of root exudates analysed analyzed by GC-MS included 4 straight chain esters, 2 phthalate esters, 1 amine and 1 phenol. Of those compounds, the phthalate esters, amine (also a sulfide) and phenol belonged to potential allelochemical class listed by Rice<sup>[24]</sup> (1984). And in some other Solanaceae plants, similar compounds have been determined as allelochemicals<sup>[25-27]</sup> (Wang, 2003; Geng 2009; Hou, 2007). As for the straight chain esters, their structures were similar to some allelochemicals determined in eggplant, such as Hexadecane acid methyl ester and heptadecane acid methyl ester<sup>[25]</sup>(Wang, 2003). Though the functional groups of the substances analysed analyzed as allelochemicals were similar to those allelochemicals, the branched-chain groups were still different.

The results of our research showed that *Solanum rostratum* potentially inhibit the growth of the surrounding plants, which would be important in evaluating its invasion success. However, the major effective compounds functioning of allelopathy of *Solanum rostratum* Dunal were still not confirmed, and there were still new compounds found<sup>[10]</sup> (Liu et al. 2019). 29 constituents of the essential oil extracted from *Solanum rostratum* exhibited significant suppressive action on seed germination and seedling development of test species<sup>[28]</sup>. Further works, such as the allelopathy of other organs and allelopathy in the field, interaction mechanism between allelochemicals and soil microorganisms, between allelochemicals and parasitic plants, were still needed to be determined.

#### 5. CONCLUSION

Root exudates of *Solanum rostratum* contained some allelochemicals, which could inhibit the germination and radicle growth of wheat and Chinese cabbage, though the effects of root exudates on shoot growth are different, with a stimulation on Chinese cabbage while a inhibition on wheat. The difference indicated a selectivity of allelopathy effect of root exudates from *Solanum rostratum* Dunal.

#### ACKNOWLEDGEMENT

This research was based on the research platform of the Key Laboratory of Global Change and Biological Invasion in Liaoning Province, and supported by the Liaoning Natural Science Foundation Guidance Plan Project (2019-ZD-0713), Shenyang Young and Middle-aged Technological Innovation Talent Support Program Project (RC170540) and And national key R&D project topics (2017YFC1503105).



## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES (全部替换吧)

- Bais, Harsh, P., et al. 2003. Allelopathy and Exotic Plant Invasion: From Molecules and Genes to Species Interactions.[J]. Science, 301(5638):1377-1380.
- Bertin C, Yang XH, Weston LA. 2003. The role of root exudates and allelochemicals in the rhizosphere. Plant Soil, 256:67-83.
- Callaway R M, Aschehong E T. 2000. Invasive plants versus their new and old neighbors: A mechanism for exotic invasion [J]. Science, 290:521-523.
- Callaway, R.M., Cipollini, D., Barto, K., Thelen, G.C., Hallett, S.G., Prati, D., Stinson, K. and Klironomos, J. 2008 Novel weapons: invasive plant suppresses fungal mutualists in America but not in its native Europe. Ecology, 89: 1043-1055.
- Hou Yong-xia, ZHOU Bao-li, WU Xiao-ling, FU Ya-wen. 2007. Allelopathy of Root Exudates of Pepper Journal of Shenyang Agricultural University, 38(4):504-507.
- GENG Guang-dong, ZHANG Su-qin, and CHENG Zhi-hui. 2009. Allelopathy and Allelochemicals of Root Exudates in Hot Pepper Acta Horticulturae Sinica, 36 (6): 873-878.
- Grigulis, K., Sheppard, A.W., Ash, J.E. & Groves, R.H. 2001. The comparative demography of the pasture weed *Echium plantagineum* between its native and invaded ranges. Journal of Applied Ecology, 38:281-290.
- Guan, G. Q., D. C. Gao, W. Y. Li, J. Ye, X. G. Xin, and S. D. Li. 1984. A quarantine weed: *Solanum rostratum*. Plant Quarantine, 4:25-28.
- Liu C, Tian J, An T, et al. 2019. Secondary Metabolites from *Solanum rostratum* and Their Antifeedant Defense Mechanisms against *Helicoverpa armigera*[J]. Journal of Agricultural and Food Chemistry, 68(1):188-197.
- Memmott, J., Simon, S.V., Paynter, Q., Sheppard, A.W. & Syrett, P. 2000. The invertebrate fauna on broom, *Cytisus scoparius*, in two native and two exotic habitats. Acta Oecologia, 21: 213-222.
- Molly E. Hunter and Eric S. Menges. 2002. Allelopathic effects and root distribution of *Ceratiola ericoides* (empetraceae) on seven rosemary scrub species American Journal of Botany, 89(7): 1113-1118.
- Pimentel, D., Lach, L., Zuniga, R. & Morrison, D. (2000) Environmental and economic costs of nonindigenous species in the United States. BioScience, 50, 53-65.
- Ping YF, Zhu JW, Zhang ZG. 2012. Effects of *Solanum rostratum* on seed germination and seedling growth of tomato [J]. Agricultural Research in the Arid Areas, 3:20-25.
- Rose M A, Harper J L. 1972. Occupation of biological space during seedling establishment [J] Journal of Ecology, 60:77-78.
- Tian XF, Bo QU. 2017. Application of DNA Barcoding in Inspection and Quarantine of the Invasive Plant *Solanum rostratum* [J]. Journal of Weed Science, 35(1):30-35.
- Vivanco, J.M., Bais, H.P., Stermitz, F.R., Thelen, G.C. & Callaway, R.M. 2004. Biogeographical variation in community response to root allelochemistry: novel weapons and exotic invasion. Ecology Letters, 4:285-292.

- Wang Fang. 2003. The mechanism of eggplant (*Solanum melongena* L.) replanting problem [Ph. D. Dissertation]. Beijing: China Agricultural University (in Chinese).—
- Wei, S. H., C. X. Zhang, Y. Liu, H. J. Huang, Q. H. Meng, H. L. Cui, and X. J. Li. 2007. Invasive weed species buffalobur (*Solanum rostratum*) and its risk assessment. Chinese Agric. Sci. Bull. 23(3):347–351.
- Weiner J, Wright DB, Castro S. 1997. Symmetry of below-ground competition between *Kochia scoparia* individuals [J]. OIKOS, 79: 85–91.
- Weir T. L., Park S. W., Vivanco J. M. 2004. Biochemical and physiological mechanisms mediated by allelochemicals. Curr. Opin. Plant Biol., 7: 472–79
- Whalen, M. D. 1979. Taxonomy of *Solanum* section *Androceras*. Gentes. Herb., 11: 359–426.
- Wilcove, D. S., Rothstein D., Dubow, J., Phillips, A. & Losos, E. (1998) Quantifying threats to imperiled species in the United States. BioScience, 48:607–615.
- Williamson G. B., Richardson D. 1988. Bioassays for allelopathy measuring treatment responses with independent controls. Journal of Chemical Ecology, 14(1):181–187.
- Zhao XH, Zhang GL, Song Z, et al. 2017. Effects of *Solanum rostratum* Invasion on Soil Properties in Different Soil Types [J]. Chinese Journal of Agrometeorology, 38(2):76–87.

1. Pimentel D, Lach L, Zuniga R, Morrison D. Environmental and economic costs of nonindigenous species in the United States. BioScience. 2000;50:53–65.
2. Wilcove DS, Rothstein D, Dubow J, Phillips A, Losos E. Quantifying threats to imperiled species in the United States. BioScience. 1998;48:607–615.
3. Memmott J, Fowler SV, Paynter Q, Sheppard AW, Syrett P. The invertebrate fauna on broom, *Cytisus scoparius*, in two native and two exotic habitats. Acta Oecologia. 2000;21:213–222.
4. Grigulis K, Sheppard AW, Ash JE, Groves RH. The comparative demography of the pasture weed *Echium plantagineum* between its native and invaded ranges. J Appl Ecol. 2001;38:281–290.
5. Bais HP, Vepachedu R, Gilroy S, Callaway RM, Vivanco JM. Allelopathy and exotic plant invasion: from molecules and genes to species interactions. Science. 2003;301(5638):1377–1380.
6. Callaway RM, Aschehong ET. Invasive plants versus their new and old neighbors: a mechanism for exotic invasion. Science. 2000;290:521–523.
7. Ping YF, Zhu JW, Zhang ZG. Effects of *Solanum rostratum* on seed germination and seedling growth of tomato. Agricultural Research in the Arid Areas. 2012;3:20–25.
8. Callaway RM, Cipollini D, Barto K, Thelen GC, Hallett SG, Prati DS, et al. Novel weapons: invasive plant suppresses fungal mutualists in America but not in its native Europe. Ecology. 2008;89(4):1043–1055.
9. Vivanco JM, Bais HP, Stermitz FR, Thelen GC, Callaway RM. Biogeographical variation in community response to root allelochemistry: novel weapons and exotic invasion. Ecology Letters. 2004;7(4):285–292.
10. Liu C, Tian J, An T, Lv FN, Jia PF, Zhou MJ, et al. Secondary metabolites from *Solanum rostratum* and their antifeedant defense mechanisms against *Helicoverpa armigera*. J Agr Food Chem. 2019;68(1):188–197.

11. Bertin C, Yang XH, Weston LA. The role of root exudates and allelochemicals in the rhizosphere. *Plant Soil*. 2003;256(1):67-83.
12. Weir TL, Park SW, Vivanco JM. Biochemical and physiological mechanisms mediated by allelochemicals. *Curr Opin Plant Biol*. 2004;7(4):472-79
13. Zhao XH, Zhang GL, Song Z, Zhang RH, Fu WD. Effects of *Solanum rostratum* invasion on soil properties in different Soil types. *J Agrometeorol*, 2017;38(2):76-87.
14. Tian XF, Ou B. Application of DNA barcoding in inspection and quarantine of the invasive plant *Solanum rostratum*. *Weed Science*, 2017;35(1):30-35.
15. Zhao J, Lislle SM, Lou A, Mario VM, Nicholas AT. (2013). Population structure and genetic diversity of native and invasive populations of *Solanum rostratum* (Solanaceae).. *Plos One*, 2013;8(11):e79807.
16. Guan GQ, Gao DC, Li WY, Ye J, Xin XG, Li SD. A quarantine weed: *Solanum rostratum*. *Plant Quarantine*. 1984;4:25-28.
17. Wei SH, Zhang CX, Liu Y, Huang HJ, Meng QH, Cui HL, et al Invasive weed species buffalobur (*Solanum rostratum*) and its risk assessment. *Chinese Agric Sci Bull*. 2007;23(3):347-351.
18. Satyal P, Maharjan S, Setzer WN. Volatile constituents from the leaves, fruits (berries), stems and roots of *Solanum xanthocarpum* from nepal. *Nat Prod Commun*, 2015;10(2):361-364.
19. Jin YN, Liu WX, Yang GQ, Wan FH, Wang JJ. Comparison of two methods in evaluating the allelopathic potential of root exudates from *Ageratina adenophora* (sprengel). *Chinese Agricultural Science Bulletin*. 2010;26(22):297-300.
20. Friebe A, Klever W, Sikora R, Schnabl H. Allelochemicals in Root Exudates of Maize. Springer US. 1998
21. Molly E, Hunter, Menges ES. Allelopathic effects and root distribution of *Ceratiola ericoides* (Empetraceae) on seven rosemary scrub species *Am Journal Bot*. 2002;89(7):1113-1118.
22. Rose MA, Harper JL. Occupation of biological space during seedling establishment. *J Ecol*. 1972;60(1):77-78.
23. Weiner J, Wright DB, Castro S. Symmetry of below-ground competition between *Kochia scoparia* individuals. *OIKOS*. 1997;79: 85-91.
24. Rice EL. Allelopathy 2th ed. London, Academic Press; 1984.
25. Wang F. The mechanism of eggplant (*Solanum melongena* L.) replanting problem. Beijing: China Agricultural University. 2003
26. Geng GD, Zhang SQ, Cheng ZH. Allelopathy and allelochemicals of root exudates in hot pepper *Acta Horticulturae Sinica*, 2009;36(6):873-878.
27. Hou YX, Zhou BL, Wu XL, Fu YW. Allelopathy of Root Exudates of Pepper *Journal of Shenyang Agricultural University*, 2007 38(4):504-507.
28. Zhou SX, Zhu XZ, Shi K, Han CX, Kuchkarova N, Zhang C, et al. Chemical composition and allelopathic potential of the invasive plant *Solanum rostratum* Dunal essential oil. *Flora - Morphology Distribution Functional Ecology of Plants*. 2021;274:151730.