

The Role of Diet in the Pathogenesis of Erectile Dysfunction

Orcid # 0000-0003-0007-5582

Publons # E-4889-2016

Researchgate: https://www.researchgate.net/profile/Shashi_Agarwal

Abstract

Erectile dysfunction (ED), characterized by a man's constant or recurrent disability to obtain and/or maintain penile erection sufficient to achieve and complete satisfactory sexual intercourse, is a common ailment. Its incidence and prevalence increases with age. ED is caused by several factors -psychological factors, organic factors, and metabolic factors. ED may also be iatrogenic and related to some drugs, pelvic surgery, or radiotherapy. The main etiology is vascular in nature, with risk factors that are common to other cardiovascular diseases. Several drugs, mechanical and prosthetic devices, and surgical procedures are used to alleviate ED. Lifestyle modulation is increasingly being added as first-line treatments for ED. These include lack of physical activity and cigarette smoking. Obesity and an unhealthy quality of diet are also lifestyles that affect ED. This manuscript **narratively** reviews the impact of excessive caloric intake, various dietary ingredients, and special diets on ED.

Keywords: erectile dysfunction, diet, cardiovascular disease, plant-based diet, alcohol, dietary macronutrients, dietary micronutrients

Introduction

Erectile dysfunction (ED) is defined as man's constant or recurrent disability to obtain and/or maintain penile erection sufficient to achieve and complete satisfactory sexual intercourse (Fourth International Consultation on Sexual Medicine)¹. It is a common disease². It is expected to affect over 322 million men globally by 2025³. Its incidence and prevalence increases with age - ED affects >50% of the elderly men⁴. ED is often accompanied by comorbid conditions such as diabetes mellitus (T2DM), hypertension (HTN), hypercholesterolemia, and obesity⁵. ED is twice as common in men with metabolic syndrome⁶. Anywhere from 20–85% of diabetic patients also have ED⁷. In addition, Wang et al. reported that hypertension was associated with an increased risk of ED (summary OR = 1.58)⁸. Its risk factors (hypertension, obesity, and smoking) are common with other cardiovascular disease (CVD)⁹, especially coronary artery disease (CAD). and ED is an independent predictor of CVD events¹⁰. The reverse is also true – patients with CAD often have ED¹¹.

The severity of erectile dysfunction is often described as mild, moderate, or severe according to the five-item International Index of Erectile Function (IIEF) questionnaire, with a score of 1–7 indicating severe, 8–11 moderate, 12–16 mild–moderate, 17–21 mild and 22–25 no erectile dysfunction¹². ED may be caused by psychological factor (absence of sexual education, phobias,

and presence of situational factors), organic factors (vascular such as atherosclerosis, venous insufficiency), neurological factors (such as stroke, spinal cord injuries, herniated disc), and metabolic factors (such as diabetes mellitus, hypogonadism, hyperprolactinemia). ED may also be iatrogenic and may be caused by some drugs (antidepressants, tranquilizers, non-selective β -blockers, antiandrogens), pelvic organ surgery or radiotherapy¹³. Almost 85% of men suffer from ED after invasive urological procedures, such as radical prostatectomy¹⁴. A plethora of evidence now clearly indicates that the predominant cause of ED seen in clinical practice is vascular^{15,16}.

ED is usually treated with oral, (including phosphodiesterase type 5 inhibitors such as sildenafil, vardenafil, and tadalafil; apomorphine; and synthetic prostaglandin E1 (alprostadil), phentolamine, and papaverine intraurethral. It may also be treated with intra-cavernosal medications, vacuum devices, or penile prosthesis^{17,18}. Injection of stem cells or platelet-rich plasma is also being used, mainly experimentally¹⁹. Herbal treatment (Panax ginseng, Pygogenol, Prelox and Tribulus terrestris) has also been tried but adequate human experiments documenting efficacy and safety are lacking²⁰. Lifestyle changes can prevent progression or improve regression of ED and is increasingly being added as first-line treatment for ED²¹. Deleterious lifestyles include lack of physical activity, obesity, unhealthy diets, and cigarette smoking. Erectile dysfunction results in a reduced self-esteem, psychological frustration, poor quality of life, and stress^{22,23}. Although not a killer disease, ED is a marker of increased risk of CVD, and all-cause mortality²⁴.

Discussion

Lifestyle modification help improve ED^{21,25-27}. A systematic review of 6 randomized controlled trials demonstrated beneficial effects of lifestyle interventions on erectile function after follow-up of 2–24 months²⁸. Lifestyles such as physical activity help prevent and improve ED²⁹. Cigarette smoking increases the risk of erectile dysfunction (Odds Ratio=1.4)³⁰ and smoking cessation helps reduce the risk³¹. Similarly, in a large study of 31,724 men who were free of ED at baseline, a 40% increased risk of ED was noted with the development of obesity³². Both weight loss and a healthy diet have been associated with improvements in erectile function³³. The European Association of Urology states that “lifestyle changes and risk factor modification must precede or accompany any erectile dysfunction treatment”³⁴.

Obesity

Obesity is significantly related to the development of ED^{32,35-37}. The Health Professionals Follow-up Study in the United States, including 31,724 men free of ED at baseline, showed a 40% increased risk of developing ED with obesity³². Prospective studies of variable duration from 5 to 25 years of follow-up reported that overweight or obese men had an increased probability (70%–96% higher) of developing ED compared with normal weight men³⁵⁻³⁷. Moreover, the severity of ED has been shown to be correlated with comorbid obesity, and the presence of the latter results in a reduced responsiveness to phosphodiesterase 5 (PDE5) inhibitors³⁸. In another study, 79% of men presenting with ED had a BMI of 25 kg/m² or higher and obese men, over 30 kg/m², had 3 times greater risk for sexual dysfunction than the general population³⁹. Furthermore, the degree of ED in obese patients is more severe than that in non-

obese patients, resulting an increased refractory rate to PDE5 inhibitors in severely obese patients⁴⁰. According to a meta-analysis, the overall refractory rate for sildenafil, tadalafil, and vardenafil is 30% to 35%, which is reduced to less than 20% when improving comorbid conditions⁴¹. Weight reduction with bariatric surgery significantly improves ED in obese men⁴². An animal study demonstrated that bariatric surgery improves glucose tolerance and elevates intra-cavernosal pressure and endothelial nitric oxide synthase and nervous nitric oxide synthase expressions in Otsuka Long-Evans Tokushima fatty rats⁴³. Moreover, Kun et al. observed that gastric bypass surgery decreases serum cholesterol and triglycerides levels and cavernosal intima-media thickness and increases endothelial function score in a human study⁴⁴.

Diet influences obesity⁴⁵. Obesity is defined by body mass index (BMI). A normal BMI is between 18.5 kg/m² to 24.9 kg/m². A BMI of 25 kg/m² to 29.9 kg/m² is considered overweight, while at or >30 kg/m² is diagnosed as obese⁴⁶. The 10-y odds of ED was 2.0 comparing men with a BMI $\geq$ 28 with men with a BMI <28 at baseline⁴⁷. A 2016 data showed that the RR of ED for obese men (BMI $\geq$ 30) was almost twice that of men with an ideal BMI (<25) (RR = 1.9)⁴⁸. Obesity results in higher levels of pro-inflammatory cytokines such as tumor necrosis factor- α and interleukin-6⁴⁹. Obese individuals have more reactive oxygen species, increased insulin resistance, and this reduces vascular nitric oxide (NO), resulting in endothelial dysfunction^{50,51}. Obesity is also associated with low-testosterone levels has been established in healthy men⁵².

Measures indicating abdominal adiposity have been suggested as being superior to BMI when evaluating ED risk⁵³. Fillo et al observed that men with abdominal obesity had a higher incidence rate of ED and the incidence rate was elevated in proportion to the degree of obesity⁵⁴. Abdominal obesity is ascertained by several anthropometric measurements⁵⁵⁻⁵⁷. Waist circumference (WC) should ideally be <102 cm in males and <88 cm in females when measured to the nearest 0.1 cm at the umbilical level in a standing position⁵⁵. The waist-hip ratio is normally 0.85 or less for women and 0.9 or less for men⁵⁶. Less commonly used is the waist-height ratio – this is calculated by dividing the WC by height. A ratio > 0.5 is consistent with central obesity⁵⁷. In a study from Slovakia, 73% of men older than 40-years-old with abdominal obesity had some degree of ED⁵⁸. Visceral adipose tissue is more metabolically active than peripheral subcutaneous fat. It is often associated with decreased glucose tolerance, reduced insulin sensitivity and adverse lipid profiles. It is associated with more inflammation, reduced testosterone levels and worse endothelial function⁵⁹. Weight loss helps⁶⁰. Patients can avoid or alleviate ED include, by reducing weight (5%–10%) or achieving a BMI $\leq$ 30 kg m²⁶⁰. In 110 obese men without diabetes, hypertension or dyslipidemia, ED was strongly correlated with waist/hip ratio and was significantly improved with weight loss and increased activity, which lowered glucose, insulin, waist/hip ratio, blood pressure and triglycerides, and increased endothelial NO production⁶¹. Another study in obese men, also showed that weight loss induced by diet and vigorous activity resulted in increased insulin sensitivity and reduced circulating insulin, and this was strongly associated with increased NO production⁶².

Fruits and Vegetables

Growing evidence supports an improvement in endothelial function and blood pressure after increased dietary flavonoid intake⁶³⁻⁶⁵ suggesting that flavonoids might be more likely than other

dietary factors to improve erectile function. Flavonoids are present in many plant-based foods/beverages, including fruit, vegetables, tea, herbs, and wine^{66,67}, and exert anti-inflammatory effects, inhibit LDL oxidation and endothelial NADPH oxidase, modulate endothelial nitric oxide (NO) synthase activity, and augment NO status⁶⁸. The beneficial effects on ED were confirmed by Wang et al. who found that there was a 10% risk reduction with each additional daily serving of fruit/vegetable consumed⁶⁹. Fruits and vegetables are rich in antioxidants⁷⁰. Antioxidants play an important role in modulating intra-cavernosal blood flow, and smooth muscle relaxation⁷¹. Zhang et al. that antioxidants improve ED by protection of NO bioavailability⁷².

Whole grains/Fiber/Nuts

The benefits of intake of whole grains and nuts is derived from the benefits of Mediterranean diet on ED. In one study, intake of 100 g pistachio nuts for 3 weeks resulted in an improvement in the IIEF-15 score (International Index of Erectile Function (IIEF) Questionnaire)⁷³. The IIEF-15 evaluates erectile dysfunction and treatment outcomes in clinical trials. Mean IIEF-15 score was 36 ± 7.5 before the diet and 54.2 ± 4.9 after the diet. Mean peak systolic velocity values before and after the pistachio diet were 35.5 ± 15.2 and 43.3 ± 12.4 cm, respectively⁷⁴. This study further supports the beneficial role of nuts in ED.

Fish

Research has revealed that omega-3 fatty acids exert beneficial effects in the development and progression of atherosclerosis⁷⁵. Fish is a good source of omega 3 polyunsaturated fats (PUFA), namely eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)⁷⁶. In a review, Zher et al. found that n-3 PUFA supplementation successfully improved endothelial dysfunction in individuals with traditional risk factors for atherosclerotic CV disease⁷⁷. Endothelial dysfunction plays an important role in the pathogenesis of ED⁷⁸. In an erectile dysfunction rat model, omega 3 fatty acids improved intra-cavernosal pressure⁷⁹. Omega 3 fatty acids in a dose of 500–1000 mg taken daily by men under age 60 years has been suggested to help improve ED^{80,81}. The direct association between fish intake and ED is still not specifically been studied.

Coffee/Tea/Chocolate

Coffee is a rich source of caffeine, antioxidants and anti-inflammatory compounds and has been implicated to have a potentially beneficial role against chronic diseases^{82,83}. Benefits have also been noted in patients with ED⁸⁴. In a cross-sectional analysis using the National Health and Nutrition and Examination Survey from 2001–2004, a nationally representative sample of the US noninstitutionalized male population (3724 men (≥ 20 years old)), the equivalent of 2–3 cups of coffee per day was associated with a lower likelihood of ED⁸⁵. Coffee is rich in polyphenols and may also increase testosterone levels^{86–88}. The result may be relaxation of the cavernous smooth muscle and improved blood flow through penile arteries⁸⁹. However, in the Health Professionals Follow-Up Study, (a prospective analysis of 21,403 men aged 40–75 years old) followed for 10 years with a 34% rate of patients with incident ED ($n = 7,298$), did not support an association between coffee and ED⁹⁰. Tea is known to increase NO and has beneficial effects in coronary artery disease^{91,92}. In animal studies, there is reduction in atherosclerotic progression in

cavernous tissue in aged rats following chronic ingestion of catechin-rich beverages like green tea⁹³. ED is associated with reduced NO production in the systemic vasculature. NO is a powerful indicator of flow mediated dilation (FMD). A decreased FMD is associated with erectile dysfunction⁹⁴. Chocolate consumption improves FMD. Ingestion of chocolate with 55% cocoa, improves FMD⁹⁵. Dark chocolate intake (with low sugar content) should therefore be beneficial in ED patients.

Alcohol

Alcohol and its relationship with cardiovascular diseases is well known⁹⁶⁻⁹⁸. A similar relationship exists between alcohol intake and ED. Cheng et al reported in a meta-analysis that one or two alcoholic drinks daily appears to reduce ED (Odds Ratio or OR=0.79)⁹⁹. In a subsequent meta-analysis of 24 studies (154,295 patients), Wang et al. also found that light to moderate alcohol consumption (<21 drinks/week) was associated with a decreased risk of erectile dysfunction (OR = 0.71)¹⁰⁰. However, regular (ever vs. never) and high alcohol consumption (>21 drinks/week) had no significant influence on the prevalence of ED (regular: OR = 0.87; high: OR = 0.99)¹⁰⁰. Moderate alcohol increases vascular NO in laboratory animals¹⁰¹ while excessive alcohol suppresses vascular NO and causes undesired structural alterations in the penile endothelium of the mouse^{102,103}. Overall, the limited data indicates that alcohol should be restricted to mild to moderate amounts (one to two drinks daily), which have been shown in a recent meta-analysis to reduce ED.

Red Meat/Saturated and Unsaturated Fat

There are no direct studies associating red meat or saturated fat intake with ED. However, given that a high fat meal increases inflammation, decreases vascular NO, increases oxidative stress, and decreases FMD, it may tend to promote atherosclerosis and ED¹⁰⁴⁻¹⁰⁶.

Vitamins/Micronutrients

The role of several vitamins and minerals have been studied in the pathogenesis of ED. The relationship between vitamin D levels and ED has been frequently questioned^{107,108}. A recent meta-analysis did not provide a strong relationship between vitamin D and the risk of ED¹⁰⁹.

Folic acid deficiency and/or supplementation may influence ED. In a meta-analysis (6 studies with 982 ED patients and 860 healthy subjects) Zhang et al. demonstrated that folic acid deficiency was an independent risk factor for ED¹¹⁰. Their work also suggested that folic acid supplement may have potentially positive effects in the treatment of ED patients. In another study, ED in diabetic patients treated with the combination of tadalafil and folic acid 5 mg improved significantly as compared with the placebo group¹¹¹. The degree of folic acid deficiency may reflect the severity of ED¹¹². It appears to be more commonly seen in younger individuals¹¹². This association may be mediated by homocysteine and higher than usual doses of folic acid may be necessary to normalize serum homocysteine levels¹¹³. Some other minerals and vitamins deserve mention. Calcium supplementation causes a small but significant decrease of blood pressure, the effect being greater in individuals with insufficient dietary intake¹¹⁴. Vitamin E is an antioxidant and increases the circulating levels of vitamin C¹¹⁵ and both may be

beneficial in ED patients. L-Arginine in a large dose (5 g per day) has been reported to have some benefit on ED¹¹⁶.

Special Diets

Dietary patterns with high content of whole grain foods and legumes and vegetables and fruits, and that limit red meat, full-fat dairy products, and food and beverages high in added sugars are associated with a reduced risk of ED¹¹⁷. The prime example, and probably the most studies, is the Mediterranean diet (MedD). This diet refers to a traditional dietary pattern of people residing around the Mediterranean Sea (Greece, Crete, and Southern Italy) and is regarded as one of the healthiest diets¹¹⁸. The MedD is characterized by a high intake of fruits, vegetables, legumes, monounsaturated fatty acids (essentially extra virgin olive oil), a moderate intake of fish and wine, and a low intake of red meat¹¹⁹. It is associated with higher levels of omega-3's, folic acid, antioxidants, and anti-inflammatory ingredients¹²⁰. MedD has a beneficial role on erectile function, both diabetic and non-diabetic men^{121,122}. In a study by Esposito et al. after 2 years, men on a Mediterranean diet consumed more fruits, vegetables, nuts, whole grain, and olive oil when compared with men on the control diet¹²³. IIEF score increased up to 22 in 13 men in the intervention group and only in 2 men in the control group¹²³. Several other studies have replicated the benefits in ED by adhering to a MedD¹²⁴⁻¹²⁶. These benefits on atherosclerosis in general, and ED can be mediated through multiple biological pathways, including a reduction of oxidative stress, inflammation, amelioration of endothelial dysfunction and insulin sensitivity, - all help increase NO release in the penis arteries¹²⁷.

Conclusion

Most patients with ED share similar risk factors that exist for coronary artery disease¹²⁸⁻¹³¹. They often have asymptomatic coronary artery stenosis¹³². The reverse is also true - the prevalence of ED is relatively high in patients with CAD¹³³. The number of studies involved in clarifying the relationship of several dietary ingredients are sparse. However, it is increasingly clear that plant-based diets, such as the Mediterranean diet, are vascular friendly^{122,123}. They help reduce the risk of ED and help mitigate its progression. The mechanisms include amelioration of insulin resistance, low grade inflammation and endothelial dysfunction¹³⁴.

Acknowledgements: None

Funding: None

Conflict of Interest: None

References

1. McCabe MP, Sharlip ID, Atalla E, et al. Definitions of Sexual Dysfunctions in Women and Men: A Consensus Statement From the Fourth International Consultation on Sexual Medicine 2015. *J Sex Med.* 2016;13:135–43.
2. Shamloul R, Ghanem H. Erectile dysfunction. *Lancet.* 2013 Jan;381((9861)):153–65.

3. Campbell JD, Milenkovic U, Usta MF, et al. The good, bad, and the ugly of regenerative therapies for erectile dysfunction. *Transl Androl Urol*. 2020;9:S252–S261. doi: 10.21037/tau.2019.10.06.
4. Kessler A, Sollie S, Challacombe B, et al. The global prevalence of erectile dysfunction: A review. *BJU Int*. 2019;124:587–99. doi: 10.1111/bju.14813.
5. Sanchez E, Pastuszak AW, Khera M. Erectile dysfunction, metabolic syndrome, and cardiovascular risks: facts and controversies. *Transl Androl Urol*. 2017 Feb;6((1)):28–36.
6. Besiroglu H, Otunctemur A, Ozbek E. The relationship between metabolic syndrome, its components, and erectile dysfunction: a systematic review and a meta-analysis of observational studies. *J Sex Med*. 2015 Jun;12((6)):1309–18.
7. Kouidrat Y, Pizzol D, Cosco T, Thompson T, Carnaghi M, Bertoldo A, et al. High prevalence of erectile dysfunction in diabetes: a systematic review and meta-analysis of 145 studies. *Diabet Med*. 2017 Sep;34((9)):1185–92.
8. Wang XY, Huang W, Zhang Y. Relation between hypertension and erectile dysfunction: a meta-analysis of cross-section studies. *Int J Impot Res*. 2018 Jun;30((3)):141–6.
9. Gandaglia G, Briganti A, Jackson G, Kloner RA, Montorsi F, Montorsi P, et al. A systematic review of the association between erectile dysfunction and cardiovascular disease. *Eur Urol*. 2014 May;65((5)):968–78.
10. Thompson IM, Tangen CM, Goodman PJ, Probstfield JL, Moinpour CM, Coltman CA. Erectile dysfunction and subsequent cardiovascular disease. *JAMA* 2005;294:2996–3002.
11. Rodriguez JJ, Al Dashti R, Schwarz ER. Linking erectile dysfunction and coronary artery disease. *Int J Imp Res*. 2005;17:S12–S18. doi: 10.1038/sj.ijir.3901424.
12. Rosen RC, Cappelleri JC, Gendrano N 3rd. The International Index of Erectile Function (IIEF): a state-of-the-science review. *Int J Impot Res*. 2002 Aug;14(4):226–44. doi: 10.1038/sj.ijir.3900857.
13. Yafi FA, Jenkins L, Albersen M, et al. Erectile dysfunction. *Nature reviews Disease primers*. 2016;2:16003. doi: 10.1038/nrdp.2016.3.
14. Rew KT, Heidelbaugh JJ. Erectile dysfunction. *Am Fam Physician*. 2016;94:820–27.
15. Kloner RA. Erectile dysfunction as a predictor of cardiovascular disease. *Int J Impot Res* 2008;20:460–5.
16. Fung MM, Bettencourt R, Barrett-Connor E. Heart disease risk factors predict erectile dysfunction 25 years later: the Rancho Bernardo Study. *J Am Coll Cardiol* 2004;43:1405–11.
17. Diniz A. F. A., Ferreira R. C., de Souza I. L. L., Silva B. A. Ionic Channels as Potential Therapeutic Targets for Erectile Dysfunction: A Review. *Frontiers in Pharmacology*. 2020;11:p. 1120. doi: 10.3389/fphar.2020.01120.
18. Kim S., Cho M. C., Cho S. Y., Chung H., Rajasekaran M. R. Novel Emerging therapies for erectile dysfunction. *The World Journal of Men's Health*. 2021;39(1):p. 48. doi: 10.5534/wjmh.200007.
19. Krzastek SC, Bopp J, Smith RP, Kovac JR. Recent advances in the understanding and management of erectile dysfunction. *F1000Res*. 2019;8:F1000 Faculty Rev-102. Published 2019 Jan 25. doi:10.12688/f1000research.16576.

20. Ho CC, Tan HM. Rise of herbal and traditional medicine in erectile dysfunction management. *Curr Urol Rep*. 2011 Dec;12(6):470-8. doi: 10.1007/s11934-011-0217-x.
21. Maiorino MI, Bellastella G, Esposito K: Lifestyle modifications and erectile dysfunction: what can be expected? *Asian J Androl*. 2015;17(1):5–10. 10.4103/1008-682X.137687.
22. Shiri R, Koskimäki J, Tammela TL, et al. Bidirectional relationship between depression and erectile dysfunction. *J Urol*. 2007;177:669–73. doi: 10.1016/j.juro.2006.09.030.
23. Peate I Breaking the silence: helping men with erectile dysfunction. *Br J Community Nurs* 2012;17:310, 312, 314–7. doi:10.12968/bjcn.2012.17.7.310.
24. Araujo AB, Travison TG, Ganz P, Chiu GR, Kupelian V, et al. Erectile dysfunction and mortality. *J Sex Med*. 2009;6:2445–54.
25. Hannan JL, Maio MT, Komolova M, Adams MA. Beneficial impact of exercise and obesity interventions on erectile function and its risk factors. *J Sex Med*. 2009;(6 Suppl 3):254–261.
26. Meldrum DR, Gambone JC, Morris MA, Esposito K, Giugliano D, Ignarro LJ. Lifestyle and metabolic approaches to maximizing erectile and vascular health. *Int J Impot Res*. 2012;24:61–68.
27. Sivaratnam L, Selimin DS, Abd Ghani SR, et al. Behavior-Related Erectile Dysfunction: A Systematic Review and Meta-Analysis. *J Sex Med* 2021;18:121–143.
28. Gupta BP, Murad MH, Clifton MM, Prokop L, Nehra A, Kopecky SL. The effect of lifestyle modification and cardiovascular risk factor reduction on erectile dysfunction: a systematic review and meta-analysis. *Arch Intern Med* 2011;171:1797–803.
29. Cheng JY, Ng EM, Ko JS, et al. : Physical activity and erectile dysfunction: meta-analysis of population-based studies. *Int J Impot Res*. 2007;19(3):245–52. 10.1038/sj.ijir.3901521.
30. Bacon CG, et al. A prospective study of risk factors for erectile dysfunction. *J Urol*. 2006;176:217–221.
31. Brackett NL, Lynne CM, Ibrahim E, Ohl DA, Sonksen J. Treatment of infertility in men with spinal cord injury. *Nat Rev Urol*. 2010;7:162–172.
32. Bacon CG, Mittleman MA, Kawachi I, et al. : Sexual function in men older than 50 years of age: results from the health professionals follow-up study. *Ann Intern Med*. 2003;139(3):161–8. 10.7326/0003-4819-139-3-200308050-00005.
33. Justin La, Natalie H. Roberts, Faysal A.Yafi. Diet and Men's Sexual Health. *Sexual Medicine Reviews*. Volume 6, Issue 1, January 2018, Pages 54-68.
34. Hatzimouratidis K, et al. Guidelines on male sexual dysfunction: erectile dysfunction and premature ejaculation. *Eur Urol*. 2010;57:804–814.
35. Feldman HA, Johannes CB, Derby CA, Kleinman KP, Mohr BA, et al. Erectile dysfunction and coronary risk factors: prospective results from the Massachusetts male aging study. *Prev Med*. 2000;30:328–38.
36. Shiri R, Koskimäki J, Hakama M, Häkkinen J, Huhtala H, et al. Effect of life-style factors on incidence of erectile dysfunction. *Int J Impot Res*. 2004;16:389–94.
37. Fung MM, Bettencourt R, Barrett-Connor E. Heart disease risk factors predict erectile dysfunction 25 years later: the Rancho Bernardo Study. *J Am Coll Cardiol*. 2004;43:1405–11.

38. Demir O, Akgul K, Akar Z, Cakmak O, Ozdemir I, Bolukbasi A, et al. Association between severity of lower urinary tract symptoms, erectile dysfunction and metabolic syndrome. *Aging Male*. 2009;12:29–34.
39. Kaya E, Sikka SC, Gur S. A comprehensive review of metabolic syndrome affecting erectile dysfunction. *J Sex Med*. 2015;12:856–875.
40. Patel JP, Lee EH, Mena CI, Walker CN. Effects of metformin on endothelial health and erectile dysfunction. *Transl Androl Urol*. 2017;6:556–565.
41. McMahon CN, Smith CJ, Shabsigh R. Treating erectile dysfunction when PDE5 inhibitors fail. *BMJ*. 2006;332:589–592.
42. Aleid M, Muneer A, Renshaw S, George J, Jenkinson AD, Adamo M, et al. Early effect of bariatric surgery on urogenital function in morbidly obese men. *J Sex Med*. 2017;14:205–214.
43. Choi YS, Lee SK, Bae WJ, Kim SJ, Cho HJ, Hong SH, et al. Bariatric surgery improves the cavernosal neuronal, vasorelaxation, and contraction mechanisms for erectile dysfunction as result of amelioration of glucose homeostasis in a diabetic rat model. *PLoS One*. 2014;9:e104042.
44. Kun L, Pin Z, Jianzhong D, Xiaodong H, Haoyong Y, Yuqian B, et al. Significant improvement of erectile function after Roux-en-Y gastric bypass surgery in obese Chinese men with erectile dysfunction. *Obes Surg*. 2015;25:838–844.
45. Hensrud DD. Diet and obesity. *Curr Opin Gastroenterol*. 2004 Mar;20(2):119-24. doi: 10.1097/00001574-200403000-00012.
46. World Health Organization. Body mass index - BMI. 2018. <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>.
47. Feldman HA, Johannes CB, Derby CA, Kleinman KP, Mohr BA, Araujo AB. Erectile dysfunction and coronary risk factors: prospective results from the Massachusetts male aging study. *Prev Med* 2000;30:328–38.
48. Cassidy A, Franz M, Rimm EB. Dietary flavonoid intake and incidence of erectile dysfunction. *Am J Clin Nutr*. 2016;103(2):534-541. doi:10.3945/ajcn.115.122010.
49. Giugliano F, Esposito K, Di Palo C, Ciotola M, Giugliano G, Marfella R et al. Erectile dysfunction associates with endothelial dysfunction and raised proinflammatory cytokines in obese men. *J Endocrinol Invest* 2004; 27: 665–669.
50. Ungvari Z, Parrado-Fernandez C, Csiszar A, de Cabo R . Mechanisms underlying caloric restriction and life span regulation: implications for vascular aging. *Circ Res* 2008; 102: 519–528.
51. Muniyappa R, Quon MJ . Insulin action and insulin resistance in vascular endothelium. *Curr Opin Clin Nutr Metab Care* 2007; 10: 523–530.
52. Kalyani RR, Dobs AS. Androgen deficiency, diabetes, and the metabolic syndrome in men. *Curr Opin Endocrinol Diabetes Obes*. 2007;14:226–34.
53. Corona G, Rastrelli G, Filippi S, Vignozzi L, Mannucci E, Maggi M. Erectile dysfunction and central obesity: an Italian perspective. *Asian J Androl*. 2014;16(4):581-591. doi:10.4103/1008-682X.126386.

54. Fillo J, Levčikova M, Ondrusova M, Breza J, Labas P. Importance of Different Grades of Abdominal Obesity on Testosterone Level, Erectile Dysfunction, and Clinical Coincidence. *Am J Mens Health*. 2017 Mar;11(2):240-245. doi: 10.1177/1557988316642213.
55. Ross R, Neeland IJ, Yamashita S, et al. Waist circumference as a vital sign in clinical practice: a Consensus Statement from the IAS and ICCR Working Group on Visceral Obesity. *Nat Rev Endocrinol*. 2020;16(3):177-189. doi:10.1038/s41574-019-0310-7.
56. <https://www.ncbi.nlm.nih.gov/mesh?Cmd=DetailsSearch&Term=%22Waist-Hip+Ratio%22%5BMeSH+Terms%5D>.
57. Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obes Rev*. 2012 Mar;13(3):275-86. doi: 10.1111/j.1467-789X.2011.00952.x.
58. Fillo J, Breza J, Levčikova M, Luha J, Vachulova A, Durdík Š, et al. Occurrence of erectile dysfunction, testosterone deficiency syndrome and metabolic syndrome in patients with abdominal obesity. Where is a sufficient level of testosterone? *Int Urol Nephrol*. 2012;44:1113–1120.
59. Esposito K, Giugliano D. Obesity, the metabolic syndrome, and sexual dysfunction in men. *Clin Pharmacol Ther*. 2011;90:169–73.
60. Moyad MA, Park K. What do most erectile dysfunction guidelines have in common. No evidence-based discussion or recommendation of heart-healthy lifestyle changes and/or Panax ginseng? *Asian J Androl*. 2012;14:830–41.
61. Esposito K, Giugliano F, Di Palo C, Giugliano G, Marfella R, D'Andrea F et al. Effect of lifestyle changes on erectile dysfunction in obese men: a randomized controlled study. *JAMA* 2004; 291: 2978–2984.
62. Roberts CK, Nosratola D, Vaziri MD, Barnard J . Effect of diet and exercise intervention on blood pressure, insulin, oxidative stress, and nitric oxide availability. *Circulation* 2002; 106: 2530–2532.
63. Schewe T, Steffen Y, Sies H. How do dietary flavanols improve vascular function? A position paper. *Arch Biochem Biophys* 2008;476:102–6.
64. Buscemi S, Rosafio G, Arcoleo G, Mattina A, Canino B, Montana M. Effects of red orange juice intake on endothelial function and inflammatory markers in adult subjects with increased cardiovascular risk. *Am J Clin Nutr* 2012;95:1089–95.
65. Jennings A, Welch AA, Fairweather-Tait SJ, Kay C, Minihane AM, Chowienczyk P. Higher anthocyanin intake is associated with lower arterial stiffness and central blood pressure in women. *Am J Clin Nutr* 2012;96:781–8.
66. Hooper L, Kroon PA, Rimm EB, Cohn JS, Harvey I, Le Cornu KA. Flavonoids, flavonoid-rich foods, and cardiovascular risk: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2008;88:38–50.
67. Bondonno CP, Yang X, Croft KD, Considine MJ, Ward NC, Rich L. Flavonoid-rich apples and nitrate-rich spinach augment nitric oxide status and improve endothelial function in healthy men and women: a randomized controlled trial. *Free Radic Biol Med* 2012;52:95–102.

68. Loke WM, Hodgson JM, Proudfoot JM, McKinley AJ, Puddey IB, Croft KD. Pure dietary flavonoids quercetin and (–)-epicatechin augment nitric oxide products and reduce endothelin-1 acutely in healthy men. *Am J Clin Nutr* 2008;88:1018–25.
69. Wang F, Dai S, Wang M, Morrison H. Erectile dysfunction and fruit/vegetable consumption among diabetic Canadian men. *Urology*. 2013 Dec;82(6):1330-5. doi: 10.1016/j.urology.2013.07.061.
70. Halvorsen BL, Carlsen MH, Phillips KM, Bohn SK, Holte K, Jacobs DR et al. Content of redox-active compounds (ie, antioxidants) in foods consumed in the United States. *Am J Clin Nutr* 2006; 84: 95–135.
71. Azadzoi KM, Schulman RN, Aviram M, Siroky MB. Oxidative stress in arteriogenic erectile dysfunction: prophylactic role of antioxidants. *The Journal of Urology*. 2005;174(1):386–393.
72. Zhang Q, Radisavljevic ZM, Siroky MB, Azadzoi KM. Dietary antioxidants improve arteriogenic erectile dysfunction. *International Journal of Andrology*. 2011;34(3):225–235.
73. <https://www.mdapp.co/international-index-of-erectile-function-iiief-questionnaire-calculator-569/>.
74. Aldemir M, Okulu E, Neşelioğlu S, Erel O, Kayıgil O. Pistachio diet improves erectile function parameters and serum lipid profiles in patients with erectile dysfunction. *Int J Impot Res*. 2011 Jan-Feb;23(1):32-8. doi: 10.1038/ijir.2010.33.
75. Jain AP, Aggarwal KK, Zhang PY. Omega-3 fatty acids and cardiovascular disease. *Eur Rev Med Pharmacol Sci*. 2015;19(3):441-5.
76. <https://ods.od.nih.gov/factsheets/Omega3FattyAcids-HealthProfessional/>.
77. Zehr KR, Walker MK. Omega-3 polyunsaturated fatty acids improve endothelial function in humans at risk for atherosclerosis: A review. *Prostaglandins Other Lipid Mediat*. 2018 Jan;134:131-140. doi: 10.1016/j.prostaglandins.2017.07.005.
78. Verveniotis A, Siasos G, Oikonomou E, Tsigkou V, Papageorgiou N, et al. The Impact of Omega 3 Fatty Acids in Atherosclerosis and Arterial Stiffness: An Overview of their Actions. *Curr Pharm Des*. 2018;24(17):1865-1872. doi: 10.2174/1381612824666180321095022.
79. Shim JS, Kim DH, Bae JH, Moon du G. Effects of Omega-3 Fatty Acids on Erectile Dysfunction in a Rat Model of Atherosclerosis-induced Chronic Pelvic Ischemia. *J Korean Med Sci*. 2016 Apr;31(4):585-9. doi: 10.3346/jkms.2016.31.4.585.
80. Kris-Etherton PM, Harris WS, Appel LJ . American Heart Association Nutrition Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation* 2002; 106: 2747–2757.
81. Buga GM, Gold ME, Fukuto JM, Ignarro LJ . Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17: 187–193.
82. O’Keefe JH, Bhatti SK, Patil HR, et al. Effects of habitual coffee consumption on cardiometabolic disease, cardiovascular health, and all-cause mortality. *J Am Coll Cardiol*. 2013;62(12):1043–1051.
83. Freedman ND, Park Y, Abnet CC, et al. . Association of coffee drinking with total and cause-specific mortality. *N Engl J Med*. 2012;366(20):1891–1904.

84. Lopez DS, Wang R, Tsilidis KK, et al. . Role of caffeine intake on erectile dysfunction in US men: results from NHANES 2001–2004. *PLoS One* 2015;10(4):e0123547.
85. Lopez DS, Wang R, Tsilidis KK, et al. . Role of caffeine intake on erectile dysfunction in US men: results from NHANES 2001–2004. *PLoS One* 2015;10(4):e0123547.
86. Sansone A, Romanelli F, Gianfrilli D, et al. . Endocrine evaluation of erectile dysfunction. *Endocrine*. 2014;46(3):423–430.
87. Paton CD, Lowe T, Irvine A. Caffeinated chewing gum increases repeated sprint performance and augments increases in testosterone in competitive cyclists. *Eur J Appl Physiol*. 2010;110(6):1243–1250.
88. Kelly DM, Jones TH. Testosterone: a vascular hormone in health and disease. *J Endocrinol*. 2013;217(3):R47–R71.
89. Adebisi A, Adaikan PG. Effect of caffeine on response of rabbit isolated corpus cavernosum to high K⁺ solution, noradrenaline and transmural electrical stimulation. *Clin Exp Pharmacol Physiol*. 2004;31(1–2):82–85.
90. Lopez DS, Liu L, Rimm EB, Tsilidis KK, de Oliveira Otto M, Wang R, Canfield S, Giovannucci E. Coffee Intake and Incidence of Erectile Dysfunction. *Am J Epidemiol*. 2018 May 1;187(5):951-959. doi: 10.1093/aje/kwx304.
91. Persson IA, Josefsson M, Persson K, Andersson RG . Tea flavanols inhibit angiotensin-converting enzyme activity and increase nitric oxide production in human endothelial cells. *J Pharm Pharmacol* 2006; 58: 1139–1144.
92. Mukamal KJ, Maclure M, Muller JE, Sherwood JB, Mittleman MA . Tea consumption and mortality after acute myocardial infarction. *Circulation* 2002; 105: 2476–2481.
93. Neves D, Assunção M, Marques F, Andrade JP, Almeida H. Does regular consumption of green tea influence expression of vascular endothelial growth factor and its receptor in aged rat erectile tissue? Possible implications for vasculogenic erectile dysfunction progression. *Age (Dordr)*. 2008 Dec;30(4):217-28. doi: 10.1007/s11357-008-9051-6.
94. Kovács I, Császár A, Toth J, Siller G, Farkas A, Tarján J, Horváth J, Koller A. Correlation between flow-mediated dilation and erectile dysfunction. *J Cardiovasc Pharmacol*. 2008 Feb;51(2):148-53. doi: 10.1097/FJC.0b013e31815e8514.
95. Faridi Z, Njike VY, Dutta S, Ali A, Katz DL . Acute dark chocolate and cocoa ingestion and endothelial function: a randomized controlled crossover trial. *Am J Clin Nutr* 2008; 88: 58–63.
96. Esselstyn CB, Jr, Gendy G, Doyle J, et al. A way to reverse CAD? *J Fam Pract*. 2014;63:356–364.
97. Esselstyn CB. A plant-based diet and coronary artery disease: a mandate for effective therapy. *J Geriatr Cardiol*. 2017;14(5):317-320. doi:10.11909/j.issn.1671-5411.2017.05.004.
98. Arnett D.K., Blumenthal R.S., Albert M.A. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American heart association task force on clinical practice guidelines. *J Am Coll Cardiol*. 2019 Mar 7 doi: 10.1016/j.jacc.2019.03.010.

99. Cheng JYW, Ng EML, Chen RL, Ko JSN . Alcohol consumption and erectile dysfunction: meta-analysis of population-based studies. *Int J Impot Res* 2007; 19: 343–352.
100. Wang XM, Bai YJ, Yang YB, Li JH, Tang Y, Han P. Alcohol intake and risk of erectile dysfunction: a dose-response meta-analysis of observational studies. *Int J Impot Res*. 2018 Nov;30(6):342-351. doi: 10.1038/s41443-018-0022-x.
101. Abou-agag LH, Khoo NK, Binsack R, White CR, Darley-USmar V, Grenett HE et al. Evidence of cardiovascular protection by moderate alcohol: role of nitric oxide. *Free Radic Biol Med* 2005; 39: 540–548.
102. Abou-agag LH, Khoo NK, Binsack R, White CR, Darley-USmar V, Grenett HE et al. Evidence of cardiovascular protection by moderate alcohol: role of nitric oxide. *Free Radic Biol Med* 2005; 39: 540–548.
103. Aydinoglu F, Yilmaz SN, Coskun B, Daglioglu N, Ogulener N . Effects of ethanol treatment on the neurogenic and endothelium-dependent relaxation of corpus cavernosum smooth muscle in the mouse. *Pharmacol Rep* 2008; 60: 725–734.
104. Nappo F, Esposito K, Cioffi M, Giugliano G, Molinari AM, Paolisso G et al. Postprandial endothelial activation in healthy subjects and in type 2 diabetic patients: role of fat and carbohydrate meals. *J Am Coll Cardiol* 2002; 39: 1145–1150.
105. Esposito K, Ciotola M, Sasso FC, Cozzolino D, Saccomanno F, Assaloni R et al. Effect of a single high-fat meal on endothelial function in patients with the metabolic syndrome: role of tumor necrosis factor- α . *Nutr Metab Cardiovasc Dis* 2007; 17: 274–279.
106. Johnson BD, Padilla J, Harris RA, Wallace JP . Vascular consequences of a high-fat meal in physically active and inactive adults. *Appl Physiol Nutr Metab* 2011; 36: 368–375.
107. Dumbraveanu I, Banov P, Arian I, Ceban E. The Correlations of Clinical and Biochemical Indices of Vitamin D with Erectile Dysfunction. *J Med Life*. 2020 Apr-Jun;13(2):144-150. doi: 10.25122/jml-2020-0009.
108. Crafa A, Cannarella R, Condorelli RA, La Vignera S, Calogero AE. Is There an Association Between Vitamin D Deficiency and Erectile Dysfunction? A Systematic Review and Meta-Analysis. *Nutrients*. 2020 May 14;12(5):1411. doi: 10.3390/nu12051411.
109. Wei Y., Chen P., Chen Q., Zhu H. Serum vitamin D levels and erectile dysfunction: A systematic review and meta-analysis. *Andrologia*. 2019;51:e13211. doi: 10.1111/and.13211.
110. Zhang Y, Zhang W, Dai Y, et al. Serum Folic Acid and Erectile Dysfunction: A Systematic Review and Meta-Analysis. *Sex Med* 2021;9:100356.
111. Hamidi Madani A, Asadolahzade A, Mokhtari G, Shahrokhi Damavand R, Farzan A, Esmaeili S. Assessment of the efficacy of combination therapy with folic acid and tadalafil for the management of erectile dysfunction in men with type 2 diabetes mellitus. *J Sex Med*. 2013 Apr;10(4):1146-50. doi: 10.1111/jsm.12047.

112. Xu J, Wang C, Zhang X, Ouyang J, Zhang J. Serum folic acid levels and erectile dysfunction: A meta-analysis and systematic review. *Andrologia*. 2021 May;53(4):e14003. doi: 10.1111/and.14003.
113. Safarinejad MR, Safarinejad S, Shafiei N . Role of methylenetetrahydrofolate reductase gene polymorphisms (C677T, A1298C, and G1793A) in the development of early onset vasculogenic erectile dysfunction. *Arch Med Res* 2010; 41: 410–422.
114. van Mierlo LAJ, Arends LR, Streppel MT, Zeegers MPA, Kok FJ, Grobbee DE et al. Blood pressure response to calcium supplementation: a meta-analysis of randomized controlled trials. *J Hum Hypertens* 2006; 20: 571–580.
115. Hamilton IM, Gilmore WS, Benzie IF, Mulholland CW, Strain JJ . Interactions between vitamins C and E in human subjects. *Br J Nutr* 2000; 84: 261–267.
116. Zorgniotti AW, Lizza EF . Effect of large doses of the nitric oxide precursor, L-arginine, on erectile dysfunction. *Int J Impot Res* 1994; 6: 33–35.
117. Wang F, Dai S, Wang M, Morrison H. Erectile dysfunction and fruit/vegetable consumption among diabetic Canadian men. *Urology*. 2013;82:1330–5.
118. Willett W.C., Sacks F., Trichopoulou A., Drescher G., Ferro-Luzzi A., Helsing E., Trichopoulos D. Mediterranean diet pyramid: A cultural model for healthy eating. *Am. J. Clin. Nutr.* 1995;61(Suppl. S6):1402S–1406S.
119. Shen J, Wilmot KA, Ghasemzadeh N, Molloy DL, Burkman G, Mekonnen G, Gongora MC, Quyyumi AA, Sperling LS. Mediterranean Dietary Patterns and Cardiovascular Health. *Annu Rev Nutr.* 2015;35:425-49. doi: 10.1146/annurev-nutr-011215-025104.
120. Mentella M.C., Scaldaferri F., Ricci C., Gasbarrini A., Miggiano G.A.D. Cancer and Mediterranean Diet: A Review. *Nutrients*. 2019;11:2059. doi: 10.3390/nu11092059.
121. Giugliano F, Maiorino MI, Bellastella G, Autorino R, De Sio M, et al. Adherence to Mediterranean diet and erectile dysfunction in men with type 2 diabetes. *J Sex Med.* 2010;7:1911–7.
122. Esposito K, Giugliano F, De Sio M, Carleo D, Di Palo C, et al. Dietary factors in erectile dysfunction. *Int J Impot Res.* 2006;18:370–4.
123. Esposito K, Ciotola M, Giugliano F, De Sio M, Giugliano G, et al. Mediterranean diet improves erectile function in subjects with the metabolic syndrome. *Int J Impot Res.* 2006;18:405–10.
124. Russo GI, Broggi G, Cocci A, Capogrosso P, Falcone M, Sokolakis I, Gül M, Caltabiano R, Di Mauro M. Relationship between Dietary Patterns with Benign Prostatic Hyperplasia and Erectile Dysfunction: A Collaborative Review. *Nutrients*. 2021 Nov 19;13(11):4148. doi: 10.3390/nu13114148.
125. Esposito K, Giugliano F, Maiorino MI, Giugliano D. Dietary factors, Mediterranean diet and erectile dysfunction. *J Sex Med.* 2010 Jul;7(7):2338-45. doi: 10.1111/j.1743-6109.2010.01842.x.
126. Esposito K, Ciotola M, Giugliano F, De Sio M, Giugliano G, D’Armiento M. Mediterranean diet improves erectile function in subjects with the metabolic syndrome. *Int J Impot Res* 2006;18:405–10.

127. Lopez-Garcia E, Hu FB. Nutrition and the endothelium. *Curr Diab Rep.* 2004;4:253–9. ; Esposito K, Giugliano D. Diet and inflammation: a link to metabolic and cardiovascular diseases. *Eur Heart J.* 2006;27:15–20.
128. Baumann F, Hehli D, Makaloski V, Schumacher M, Schönhofen H, Diehm N. Erectile dysfunction - overview from a cardiovascular perspective. *Vasa.* 2017 Aug;46((5)):347–53.
129. Chaudhary RK, Shamsi BH, Chen HM, Tan T, Tang KF, Xing JP. Risk factors for erectile dysfunction in patients with cardiovascular disease. *J Int Med Res.* 2016 Jun;44((3)):718–27.
130. García-Cruz E, Leibar-Tamayo A, Romero J, Piqueras M, Luque P, Cardeñosa O, et al. Metabolic syndrome in men with low testosterone levels: relationship with cardiovascular risk factors and comorbidities and with erectile dysfunction. *J Sex Med.* 2013 Oct;10((10)):2529–38.
131. Uddin SM, Mirbolouk M, Dardari Z, Feldman DI, Cainzos-Achirica M, DeFilippis AP, et al. Erectile dysfunction as an independent predictor of future cardiovascular events. *Circulation.* 2018 Jul;138((5)):540–2.
132. Lee JY, Lee SR, Lee SY. Prevalence of Asymptomatic Coronary Artery Stenosis Based on Coronary Computed Tomography Angiography in Adults with Erectile Dysfunction: A Cross-Sectional Study. *Med Princ Pract.* 2020;29(6):565-571. doi: 10.1159/000508876.
133. Foroutan SK, Rajabi M. Erectile dysfunction in men with angiographically documented coronary artery disease. *Urol J.* 2007 Winter;4(1):28-32.
134. Esposito K, Giugliano D. Lifestyle/dietary recommendations for erectile dysfunction and female sexual dysfunction. *Urol Clin North Am.* 2011;38:293–301.