

# Insights into the Metabolites Conferring Pathogenicity of *Xanthomonas oryzae* and its inhibition by *Trichoderma longibrachiatum* EF5

## ABSTRACT

**Aims:** The present study was aimed to evaluate the effect of volatile and soluble metabolites of *Trichoderma longibrachiatum* EF5 against *Xanthomonas oryzae* pv. *oryzae*, and to identify the metabolites produced by *Xanthomonas oryzae* pv. *oryzae* in culture filtrate.

**Study design:** *In vitro* bioassay with CRD.

**Place and Duration of Study:** Biocatalysts laboratory, Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore, 2019-2020.

**Methodology:** Inverted plate, bipartite plate and agar well method was done under *in vitro* to observe the efficacy of *T. longibrachiatum* EF5 VOC against *X. oryzae* pv. *oryzae*. Analysis of *Xanthomonas oryzae* pv. *oryzae* metabolites by using Gas Chromatography and Mass Spectrometry (GC-MS).

**Results:** *T. longibrachiatum* EF5 VOC and metabolites completely inhibited the growth of *X. oryzae* pv. *oryzae* in inverted plate assay, whereas in bipartite and agar well diffusion assays unmeasurable growth of *X. oryzae* pv. *oryzae* was observed. The metabolites or Diffusible signal factors such as butyrolactone, propionic acid derivatives, phenyl acetic acid, hydrofurans, picoxystrobin, benzoic acid derivatives were produced by *X. oryzae* pv. *oryzae* in the growing medium. The role of these metabolites revealed that they are involved in pathogenicity, virulence, quorum sensing and the synthesis of antioxidant.

**Conclusion:** *T. longibrachiatum* EF5 volatile and soluble metabolites can be used as biocontrol agent against *X. oryzae* pv. *oryzae*.

**Keywords:** [Volatile organic compounds, *Trichoderma longibrachiatum*, *Xanthomonas oryzae* pv. *oryzae*, diffusible signal factors]

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is one of the major food crops which was infected by different pathogens such as fungal, bacterial and viruses at all stages that affect the yield and grain quality. Among them, the most common disease, both at nursery and main field is bacterial blight caused by *Xanthomonas oryzae* pv. *oryzae* (Arshad *et al.*, 2015). The pathogen *Xanthomonas oryzae* pv. *oryzae* causes infection at all phases of growth with characteristic symptoms. It invades plants through natural pores and wounds, causing water-soaked lesion. The lesion extends from leaf tip as V shaped wavy margin and cause yellowing and results in death of the plant (Nino-Liu *et al.*, 2006). *X. oryzae* pv. *oryzae* has different mechanisms to cause infection such as hypersensitive response and pathogenicity (*hrp*) genes, secondary metabolites and toxins, type II and III secretory system, extracellular enzymes, polysaccharides and diffusible signal factors. Different management strategies have been implemented to combat the disease such as use of chemicals, resistant varieties and biocontrol agents (Sun *et al.*, 2005; Rajeswari *et al.*, 2005).

Plant growth promoting rhizospheric microorganisms were identified and employed as antagonist against *X. oryzae* pv. *oryzae* (Gnanamanickam, 2009). Fungal and bacterial microorganisms either directly kill the pathogen or indirectly induce defense in plants. These organisms produce antibiotics, lytic enzymes and other mechanisms by which they reduce the pathogen growth (Bardin *et al.*, 2015). Previous studies have shed light on bacterial antagonists such as *Bacillus* and *Pseudomonas* against *X. oryzae* pv. *oryzae* and *Serratia* sp. etc. that induced systemic resistance (Vleeschauwer *et al.*, 2009).

Endophytes encompassed of fungal and bacteria which live inside the plants without causing harm to the plant (Rodriguez *et al.*, 2008). Such endophytes can be isolated from internal parts of plant without contaminating with epiphytes. They enter plants through natural pores and wounds and colonize the plants acropetally to all parts (Hallmann *et al.*, 1997). Further, they assist the host for uptake of nutrients, inducing defense and production of plant growth hormones (Moronta-Barrios *et al.*, 2018). In the same way, as direct application of biocontrol agents, these organisms produce volatile organic compounds (VOCs) which act as a medium for disease suppression by direct inhibition of plant pathogens and indirectly by inducing defense response and plant growth promotion (Ryu *et al.*, 2003). These VOCs from microorganisms are produced naturally and alters during interaction with other organisms such as microbes, nematodes and plants. Ryu *et al.* (2004) reported the first VOC compound 2, 3-butanedial which induce defense against *Pectobacterium carotovorum* ssp. *carotovorum* in *Arabidopsis thaliana*. Many synthetic VOCs were exploited against plant pathogens for their antimicrobial activity such as dimethyl disulfide, 1-undecene, benzaldehyde, benzothiazole, dimethyl trisulfide, cyclohexanol, decanal, 2-ethyl-1-hexanol, methyl pyrazine and some mid- and long-chain alkanes, alkenes and alcohols (Kai *et al.*, 2009). VOCs from *B. subtilis* reduced *Escherichia coli* motility and increased resistance to antibiotics (Kim *et al.*, 2013). *Bacillus subtilis* D13 volatiles reduced the motility of *X. oryzae* pv. *oryzae* and altered the surface morphology with concentrated cytoplasm. Among the 12 VOCs profiled in GC-MS, 0.48 mg decyl alcohol and 2.4 mg 3,5,5-trimethylhexanol inhibited the growth of pathogens (Xie *et al.*, 2016). However, despite a large number of reports on this topic, there are no available reports on endophytic fungal strains for controlling rice bacterial blight. The study was aimed to exploit the virulence factors of *Xanthomonas oryzae* pv. *oryzae* and the volatiles and metabolites mediated inhibition by fungal antagonist *Trichoderma longibrachiatum* EF5

## **2. MATERIAL AND METHODS**

### **2.1 MICROORGANISMS AND CULTURE CONDITIONS**

Endophytic fungus *Trichoderma longibrachiatum* EF5 isolated from rice leaves was obtained from Biocatalysts Lab, Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore. The pathogen *Xanthomonas oryzae* pv. *oryzae* (Xoo) was collected from Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore. The fungus and bacteria were maintained in Petri plates containing potato dextrose agar (PDA) and nutrient agar (NA) at  $28 \pm 2$  °C, respectively.

## **2.2 PROFILING SOLUBLE METABOLITES OF *X. oryzae* pv. *oryzae* RESPONSIBLE FOR PATHOGENICITY**

The pathogen, *X. oryzae* pv. *oryzae* (Xoo) was grown in 250 ml Potato Dextrose (PD) broth. Three replications were maintained and the set up was kept at  $28 \pm 2$  °C for 3 days. Potato Dextrose broth without inoculation served as a control. After incubation the culture was centrifuged at 6000rpm for 20 min at 4°C. The filtered broth was mixed with equal quantity of ethyl acetate and kept in shaker for overnight, concentrated using vacuum flash evaporator. Consequently, the crude metabolite extracted in 1 ml methanol was further used for GCMS analysis after passing through a 0.2µm syringe filter (Meena, 2017).

The purified crude methanolic extract was subjected to GC-MS analysis in a Perkin Elmer GC-MS Clarus® SQ 8 equipped with DB-5MS (Agilent, USA) capillary standard non-polar column with dimensions of 0.25mm OD x 0.25 µm ID x 30 m length. The instrument was set to an initial temperature of 40°C, and the injection port temperature was ensured at 220°C, interface temperature set 250°C, source kept at 220°C, oven temperature-programmed as 75°C for 2 min, 150°C at 10°C/min, up to 250°C at 10°C per min. The GC conditions were as followed: 1:12 split, helium carrier at 20 psi. The MS conditions were: positive ion mode, electron impact spectra at 70 eV. The mass spectral scan range was set at 50 to 600 Da. The MS peaks were determined by their scatter pattern. The linear regression coefficient was used to calculate the concentrations in the samples from peak areas obtained in the chromatographs. The bioactive molecules were identified by comparison of mass spectra with NIST 08 Mass Spectra Library (National Institute of Standards and Technology). The name, molecular weight, and structure were ascertained from NIST, PubChem, and HMDB databases (Leylaie and Zafari, 2018).

## **2.3 ANTAGONISTIC ASSAY OF VOLATILE ORGANIC COMPOUNDS (VOCs) OF EF5 AGAINST Xoo**

The VOC mediated antagonistic assay was performed both by inverted and bipartite plate assays. An 8 mm disc of *T. longibrachiatum* EF5 was inoculated in bottom plate containing PDA and incubated for 3 days at  $28 \pm 2$ ° C. After 3 days, 48 h old culture of *Xanthomonas oryzae* pv. *oryzae* was streaked on another bottom plate and placed above to

the antagonist plate for the exposure of VOC and sealed with parafilm to prevent the VOC from escaping. A plate without antagonist but containing PDA alone was used as a control. Three replications were maintained and it was completely randomized. The plates were incubated for 3 days at  $28\pm 2^{\circ}$  C to examine the effect of VOC (Meena, 2017). While in bipartite plate, 8 mm disc of *T. longibrachiatum* EF5 was placed at periphery of one compartment and incubated at  $28\pm 2^{\circ}$  C for 2 days. After incubation, 48 h old culture of *X. oryzae* pv. *oryzae* was streaked on another compartment, sealed and incubate at  $28\pm 2^{\circ}$  C for 2 days. Three replications with completely randomized design were maintained.

## **2.4 SOLUBLE METABOLITES (SMs) OF EF5 AGAINST *Xoo***

The antagonistic effect of SMs against *Xoo* was assayed by seeded agar plate technique (Meena, 2017). *T. longibrachiatum* EF5 was grown in PD broth for 10 days and the filtrate was obtained by separating the mycelial mat through Whatman filter paper. The filtrate was then passed through 0.2 $\mu$ m filter. The non-volatile metabolite was mixed with warm PDA medium at 25 per cent concentration and plated in Petri plate. The pathogen load was adjusted to 0.1 OD<sub>600</sub> and 100 $\mu$ l was poured on each agar well. PDA medium without metabolites served as control. These plates were maintained at  $28\pm 2^{\circ}$  C till colony in control plate was visible. Three replications with completely randomized design were maintained.

## **2.5 EFFICACY OF EXTRACTED CRUDE METABOLITES**

*T. longibrachiatum* EF5 was grown in PD broth for 10 days and the filtrate was obtained by separating the mycelial mat through Whatman filter paper. The filtrate was then passed through 0.2 $\mu$ m membrane filter. Then it was mixed with equal volume of ethyl acetate and kept in shaker overnight. The mixture was placed in a separating funnel, and allowed to stand until the aqueous and organic phases are separated. The organic phase was collected from the separating funnel. Crude metabolite was separated in a vacuum flask evaporator using methanol. Approximately, 100 $\mu$ L of *X. oryzae* pv. *oryzae* culture (OD<sub>600</sub>=0.1) was spread on a PDA plate. Four uniform wells were drilled using sterile cork borer and 50  $\mu$ l of crude metabolite was placed in each agar well, incubated at  $28\pm 2^{\circ}$  C. The formation of zone around the well was measured (Rani, 2017).

## **3. RESULTS AND DISCUSSION**

### **3.1 DIFFUSIBLE SOLUBLE METABOLITES CONFERRING *XOO* PATHOGENICITY**

In the present study, we profiled the soluble metabolites and diffusible soluble factors responsible for virulence in *Xanthomonas oryzae* pv. *oryzae*. The soluble metabolites or diffusible soluble factors (DSF) responsible for pathogenicity were investigated in GC-MS. More than 40 compounds produced by *Xoo* cultured in PDB medium were detected; nevertheless, PDB medium can produce many compounds likewise. The same compounds

produced by *Xoo* and PDB medium were deducted, and 33 compounds specifically produced by *Xoo* were identified, including acids, alcohols, ketones, benzene derivatives, and esters. The metabolites present in *Xoo* were indicated in Table.1.

Among the DSF compounds mentioned above, butyrolactone involved in quorum sensing and biofilm formation. Compounds such as 1H-indene (Chiarini *et al.*, 1980), nonanal (Kubo *et al.*, 1995) and phenyl acetic acid (Kim *et al.*, 2004) whereas Picoxystrobin is commercially using as a fungicide. Hence the pathogenicity and virulence factors of *Xoo* have to be quenched down by an appropriate antagonist. The results expedited key metabolites such as benzoic acid and other benzene derivatives. The yellow pigmentation of *Xoo* was due to xanthomonadiins which is produced from 3-Hydroxybenzoic Acid and 4-Hydroxybenzoic acid. Previous reports suggest that hydroxyl benzoic acid is one of the virulent factors (Zhou *et al.*, 2013) for *Xanthomonas* and it also protects the bacteria from photooxidative damage (Rajagopal *et al.*, 1997). The results of the investigation also confirm the presence of Xanthomonadiin biosynthetic pathway and the strain was virulent. During pathogenesis, *Xanthomonas campestris* pv. *manihotis* produced blight inducing toxin 3(methylthio) propionic acid (MTP acid) in cassava leaves. The necrotic tissues also contain sulphur containing compounds thiopropionic acid is more volatile than MTP acid (Perreux *et al.*, 1986). Likewise, the present study reported that identification of Pyrrolo[1,2-a] pyrazine-1,4-dione, hexahydro-3-(2-methylpropyl), in the metabolites of *Xoo* expected that these compounds may be derivative to MTP. Similar compounds identified by Noda *et al.* (1980) in ethyl acetate extract of *Xoo* were 3-methylthiopropionic acid, trans-3-methylthioacrylic acid, phenylacetic acid, isovaleric acid, succinic acid and fumaric acid. When the culture suspension was treated on rice leaves it induced necrosis and chlorosis at higher concentration (2000 µg/ml). The results, further confirmed the presence of succinimide, phenyl acetic acid and other phenolic derivatives.

Another phytotoxic compound produced by *X. albilineans* was albidin which is the major pathogenicity factor to cause symptoms in sugarcane (Royer *et al.*, 2004). Similar to previous reports on virulence factors, *Xoo* also registered Picoxystrobin. Few soluble compounds such as 2is (2S, 4S)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran-borate may act in signal transduction pathway and in inter and intra species communication as reported in *Vibrio harveyi* (Chen *et al.*, 2002), whereas in *S. typhimurium* it is (2R,4S)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran acts as auto inducer (Miller *et al.*, 2004). The results also revealed butyrolactone and other fatty acid derivatives, which are involved in cell to cell communication and bio-film formation. This fatty acid family regulates QS signaling and involved in DSF synthesis (Hu *et al.*, 2018). Numerous reports are available on fatty acid derived AHL 3-hydroxypalmitic acid methyl ester by Gram negative bacteria *Ralstonia*

*solanacearum* (Flavier *et al.*, 1997). Phenylacetic acid produced in *Xoo* culture is argued for depressive growth of young rice seedling roots as supported by Egawa *et al.*, 1967. It can be concluded that the metabolites produced by *Xoo* were utilized for its growth, survival, cell-cell communication, signaling, virulence and pathogenicity (Hu *et al.*, 2018).

In our investigation, we attempted to control *Xoo* pathogenicity by an endophytic antagonistic fungus *T. longibrachiatum* EF5.

### **3.2 TRICHODERMA LONGIBRACHIATUM EF5 ON THE VIRULENCE OF *Xoo***


The effect of strain EF5 volatiles on *Xoo* virulence was evaluated (Figure.1a, b). The results showed that the VOC blends emitted by EF5 exerted strongest inhibition on *Xoo* cells as evident by inverted plate assay. The pathogen was not grown in the VOC exposed plate whereas in control, yellow colony was observed. While, when the VOC exposure was stopped and kept for further incubation period, *Xoo* growth was initiated. Further, bipartite plate assay also demonstrated the VOC mediated growth inhibition of *Xoo*, where minute growth was observed. VOCs of *T. longibrachiatum* EF5 expressed strong inhibition activity in inverted plate assay. In the bipartite assay considerable reduction in growth was noticed. Since, both the methods were evaluated for its volatile mediated inhibition, this change in inhibition might be due to the diffusion of volatiles within the plate and its interaction with pathogen. In inverted plate, VOC from *T. longibrachiatum* EF5 directly interacted with the pathogen which was placed ventrally whereas in bipartite, the VOC has to diffuse through the plate to another compartment. The inhibition might be also due to changes in the VOC production by *T. longibrachiatum* EF5 on due course of time during the interaction. These volatiles inhibit the normal growth of pathogen with change in morphology and color of the pathogen. Similar result was revealed by Xie *et al.* (2016) in which *B. cereus* D13 strongly inhibited *X. oryzae* pv. *oryzae* growth and reduced the motility and virulence. Co-cultivation of *Xanthomonas* sp. with D13 modified the cytoplasm with transformed exterior morphology under ultramicroscopic study. This may cause leakage of cell content and disrupt the normal physiological process. When *Xanthomonas* sp. was exposed to VOCs produced by other microorganisms in the soil might reduce the disease incidence and motility of the pathogen. The high vapor pressure of volatiles made it to pass through the soil pores and air allows communicating in short and long distance. Research states that volatiles of *Trichoderma* have both antifungal (Meena *et al.*, 2017; Li *et al.*, 2018) and antibacterial activity (Li *et al.*, 2019). Our previous study stated that *Trichoderma longibrachiatum* EF5 produced VOCs such alcohols, esters, aldehydes, ketones and terpenes, more specifically longifolene, cedrane, caryophyllene and cuprenene (Sridharan *et al.*, 2020). The result suggested that the synergistic action of VOC blend inhibited the growth of *X. oryzae* pv. *oryzae*. Nevertheless,

Single VOC is not much effective than VOC blends. The bacterial pathogen *Xoo* was significantly inhibited by the soluble metabolites. Since, no growth was visible in the agar well showed that soluble metabolites suppressed the growth of the pathogen by direct interaction (Figure.1c). The crude metabolite of *T. longibrachiatum* EF5 expressed a halo zone of 8 mm around the agar well by suppressing the growth of pathogen. The halo region is the indication of direct action of crude metabolite against *Xoo* (Figure. 1d). Many antagonistic bacteria inhibited the *X. oryzae* pv. *oryzae* such as fluorescent *Pseudomonas* (Shivalingaiah and Umesha, 2013), *Bacillus* sp., *B. subtilis*, *Pseudomonas putida* and *Enterobacter* sp. (Yousefi *et al.*, 2018).


**Table1. Non-volatile compounds profiled in *Xanthomonas oryzae* pv. *oryzae***

RT	Compound	Area	RT	Compound	Area
3.314	Butyrolactone	0.716	7.265	1H-Indene, 1-methylene-	1.75
3.514	Desulphosinigrin	0.267	7.375	Dodecane	4.289
3.556	Galacto-heptulose	0.268	12.297	2,4-Di-tert-butylphenol	1.632
3.779	Ethyl cyanoacetate	0.275	12.572	Benzoic acid	0.331
3.959	Benzaldehyde	0.482	12.687	2(3H)-Furanone, dihydro-5-phenyl-	0.483
4.123	Phenol	0.381	13.132	Pentadecane, 5-methyl-	0.366
4.469	Decane	0.469	16.514	3-Methyl-1,4-diaza bicyclo[4.3.0]nonan-2,5-dione,N-acetyl-	0.334
5.134	Benzyl methallyl sulphide	0.368	17.919	Heptadecane, 3-methyl-	0.391
5.134	Benzeneethanamine	0.371	20.756	Diethyltrisulphide	1.264
5.314	Ethanone, 1-(1H-pyrrol-2-yl)-	0.348	20.921	Pyrrolo[1,2-a] pyrazine-1,4-dione, hexahydro-3-(2-methylpropyl)-	2.112
5.377	3-Acetyl-1H-pyrroline	0.501	21.645	L-(+)-Ascorbic acid 2,6-dihexadecanoate	2.513
5.975	Nonanal	0.528	21.661	n-Hexadecanoic acid	2.45
6.140	Benzaldehyde dimethyl acetal	0.304	24.852	Picoxystrobin	0.928
6.205	Phenylacetic acid, cyclobutyl ester	0.622	25.013	L-Ascorbic acid, 6-octadecanoate	0.791
6.405	Pyrimidine	0.321	25.027	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	0.494
6.424	Succinimide	0.345	25.407	Dasycarpidan-1-methanol, acetate (ester)	0.787
6.725	Cycloheptatrienone	0.639	34.756	Cyclohexanol	0.862

**Figure1. Effect of volatile and soluble metabolites of *Trichoderma longibrachiatum* EF5 against *Xanthomonas oryzae* pv. *oryzae***







Inhibition of *Xanthomonas oryzae* pv. *oryzae* growth in (I) inverted plate assay – complete growth inhibition by VOCs, (II) bipartite plate assay – restricted growth mediated by VOCs, (III) seeded agar assay – growth inhibition by soluble metabolites, (IV) crude metabolite well assay – growth inhibition and formation of halo zone

#### 4. CONCLUSION

In the present investigation, Thus, volatile and soluble metabolites from *T. longibrachiatum* EF5 might act as fumigant, thereby suppressing Xoo growth. Based on the metabolite concentration the growth of the pathogen was completely inhibited in crude metabolite but unmeasurable growth was observed in culture filtrate. Hence it can be concluded that, there is great potential of developing biological bactericide with the endophytic fungi *T. longibrachiatum* EF5 and its metabolites for management of bacterial leaf blight disease of rice.

#### REFERENCE

1. Arshad HMI, Naureen S, Saleem K, Ali S, Jabeen T, Babar MM. Morphological and biochemical characterization of *Xanthomonas oryzae* pv. *oryzae* isolates collected from Punjab during 2013. *Adv Life Sci*. 2015. 3: 125–130.
2. Bardin M, Ajouz S, Comby M, Lopez-Ferber M, Graillet B, Siegwart M. Is the efficacy of biological control against plant diseases likely to be more durable than that of chemical pesticides? *Frontiers in Plant Science*. 2015. 6:566.
3. Chen X, Schauder S, Potier N, Van Dorsselaer A, Pelczar I, Bassler BL, and Hughson FM. Structural identification of a bacterial quorum-sensing signal containing boron. *Nature*. 2012. 415(6871): 545-549.
4. Ewbank E, Maraite H. Conversion of methionine to phytotoxic 3-methylthiopropionic acid by *Xanthomonas campestris* pv. *manihotis*. *J Gen Microbiol*. 1990. 136(7):1185-1189.
5. Fernando WGD, Ramarathnam R, Krishnamoorthy AS, Savchuk SC. Identification and use of potential bacterial organic antifungal volatiles. *Soil Biol Biochem*. 2005. 37:955–964.
6. Flavies, Albert B, Steven J, Clough, Mark, Schell A, Timothy, Denny P. Identification of 3-hydroxypalmitic acid methyl ester as a novel autoregulator controlling virulence in *Ralstonia solanacearum*. *Mol microbiol*. 1997. 26(2): 251-259.
7. Gnanamanickam SS. An overview of progress in biological control, 8<sup>th</sup> ed. *Biological Control of Rice Diseases*. Springer. 2009.
8. Hallmann J, Quadt-Hallmann A, Mahaffee WF, Kloepper JW. Bacterial endophytes in agricultural crops. *Can. J. Microbiol*. 1997. 43: 895-914
9. Jun Yuan, Waseem Raza, Qirong Shen, Qiwei Huang. Antifungal Activity of *Bacillus amyloliquefaciens* NJN-6 Volatile Compounds against *Fusarium oxysporum* f. sp. *cubense*. *Appl Environ Microbiol*. 2012. 78(16): 5942–5944
10. Kai M, Haustein M, Molina F, Petri A, Scholz B, Piechulla B. Bacterial volatiles and their action potential. *Appl Microbiol Biot*. 2009. 81: 1001–1012.
11. Kim KS, Lee S, Ryu CM. Interspecific bacterial sensing through airborne signals modulates locomotion and drug resistance. *Nat Commun*. 2013. 4: 1809.
12. Leylaie S, Zafari D. Antiproliferative and antimicrobial activities of secondary metabolites and phylogenetic study of endophytic *Trichoderma* species from *Vinca* plants. *Front Microbiol*. 2018 9: 1484.
13. Li N, Islam MT, Kang S. Secreted metabolite-mediated interactions between rhizosphere bacteria and *Trichoderma* biocontrol agents. *PLoS ONE*. 2019. 14(12).
14. Li N, Alfiky A, Wang W, Islam M, Nourollahi K, Liu X, et al. Volatile compound-mediated recognition and inhibition between *Trichoderma* biocontrol agents and *Fusarium oxysporum*. *Front. Microbiol*. 2018. 9:2614.
15. Zhou L, Huang LW, Wang JY, Sun S, Chen G, Poplawsky A, He YW. The Rice Bacterial Pathogen *Xanthomonas oryzae* pv. *oryzae* Produces 3-Hydroxybenzoic Acid and 4-

- Hydroxybenzoic Acid via XanB2 for Use in Xanthomonadin, Ubiquinone, and Exopolysaccharide Biosynthesis. *MPMI*. 2013. 26(10): 1239–1248
16. Meena M, Swapnil P, Zehra A, Dubey MK, Upadhyay RS. Antagonistic assessment of *Trichoderma* spp. by producing volatile and non-volatile compounds against different fungal pathogens. *Arch. Phytopathol. Plant Prot.* 2017. 50: 629–648.
  17. Miller ST, Xavier KB, Campagna SR, Taga ME, Semmelhack MF, Bassler BL, et al. *Salmonella typhimurium* recognizes a chemically distinct form of the bacterial quorum-sensing signal AI-2. *Mol Cell.* 2004.15(5):677-87.
  18. Moronta-Barrios F, Gionechetti F, Pallavicini A, Marys E, Venturi V. Bacterial microbiota of rice roots: 16S-based taxonomic profiling of endophytic and rhizospheric diversity, endophytes isolation and simplified endophytic community. *Microorganisms*. 2018. 6(14).
  19. Nino-Liu D, Ronald P, Bogdanove A. *Xanthomonas oryzae* pathovars: model pathogens of a model crop. *Mol. Plant Pathol.* 2006. 7: 303–324.
  20. Noda T, Soto Z, Kobayashi H, Iwazakis S, Okuda S. Isolation and structural elucidation of phytotoxic substances produced by *Xanthomonas campestris* pv. *oryzae*. *Ann of the Phytopathol Society of Japan*. 1980. 46(663466).
  21. Perreaux D, Maraite H, Meyer JA, Detection of 3(methylthio) propionic acid in cassava leaves infected by *Xanthomonas campestris* pv. *manihotis*. *Physiol Mol PI Pathol*. 1986. 28(3):323-328
  22. Rajagopal L, Sundari CS, Balasubramanian D, Sonti R. The bacterial pigment xanthomonadin offers protection against photodamage. *FEBS Lett.* 1997. 415: 125–128.
  23. Rani R, Sharma D, Chaturvedi M, Yadav JP. Antibacterial Activity of Twenty Different Endophytic Fungi Isolated from *Calotropis procera* and Time Kill Assay. *Clin Microbiol.* 2017. 6: 280
  24. Rodriguez RJ, Henson J, Van Volkenburgh E, Hoy M, Wright L, Beckwith F, Kim Y, Redman RS. Stress tolerance in plants via habitat-adapted symbiosis. *International Society of Microb Ecol.* 2008. 2: 404– 416
  25. Royer M, Costet L, Vivien E, Bes M, Cousin A, Damais A et al. Albicidin pathotoxin produced by *Xanthomonas albilineans* is encoded by three large PKS and NRPS genes present in a gene cluster also containing several putative modifying, regulatory, and resistance genes. *Mol. Plant Microbe Interact.* 2004. 17(4):414-27.
  26. Ryu CM, Farag MA, Hu CH, et al. Bacterial volatiles promote growth in *Arabidopsis*. *Proceedings of the National Acad Sci.* 2004. 4927–4932.
  27. Ryu CM, Farag MA, Hu CH, et al. Bacterial volatiles promote growth in *Arabidopsis*. *Proceedings of the National Acad Sci.* 2003. 4927–4932.
  28. Schirmbock M, Lorito M, Wang YL, Hayes CK, Arisan-Atac I, Scala F, et al. Parallel formation and synergism of hydrolytic enzymes and peptaibol antibiotics, molecular mechanisms involved in the antagonistic action of *Trichoderma harzianum* against phytopathogenic fungi. *Appl Environ Microbiol.* 1994. 60(12):4364-70.
  29. Xie S, Zang s, Wu H, Uddin F, Gao R. Antibacterial effects of volatiles produced by *Bacillus* strain D13 against *Xanthomonas oryzae* pv. *oryzae*. *Mol Plant Pathol.* 2016. 19(1):49-58
  30. Shivalingaiah, Umesha. *Pseudomonas fluorescens* inhibits the *Xanthomonas oryzae* pv. *oryzae*, the bacterial leaf blight pathogen in rice. *Canadian J. Plant Protect.* 2013. 1(5): 147
  31. Sun QH, Hu J, Huang GX, Ge C, Fang RX, He CZ. Type-II secretion pathway structural gene xpsE, xylanase- and cellulase secretion and virulence in *Xanthomonas oryzae* pv. *oryzae*. *Plant Pathol.* 2005. 54: 15–21.
  32. Vleeschauwer DD, Chernin L, Hofte MM. Differential effectiveness of *Serratia plymuthica* IC1270-induced systemic resistance against hemibiotrophic and necrotrophic leaf pathogens in rice. *BMC Plant Biol.* 2009. 9:9

33. Yousefi H, Hassanzadeh N, Behboudi K, Firouzjahi F. Identification and determination of characteristics of endophytes from rice plants and their role in biocontrol of bacterial blight caused by *Xanthomonas oryzae* pv. *oryzae*, Hellenic Plant Prot J 2018. 11(1): 19-33
34. Hu Z, Dong H, Ma JC, Yu Y, Li KH, Guo QQ, et al. Novel *Xanthomonas campestris* long-chain-specific 3-oxoacyl-acyl carrier protein reductase involved in diffusible signal factor synthesis. mBio. 2018. 9:e00596-18
35. Kubo A, Lunde CS, Kubo I. Antimicrobial activity of the olive oil flavor compounds. J Agric Food Chem. 1995;43(6):1629–1633. doi: 10.1021/jf00054a040.
36. Chiarini A, Ferranti A, Giovanninetti G, Matteuzzi D. Attività antimicrobica di alcuni idrazoni contenenti unità benzofuranica e 1H-indenica [Antimicrobial activity of various hydrazones containing benzofuran and 1H-indene units]. Farmaco Sci. 1980. 35(5):413-7.
37. Kim Y, Cho JY, Kuk JH, Moon JH, Cho JI, Kim YC, Park KH. Identification and antimicrobial activity of phenylacetic acid produced by *Bacillus licheniformis* isolated from fermented soybean, Chungkook-Jang. Curr Microbiol. 2004. 48(4):312-7.