

Nutritional and sensory profiles of functional breads produced from partial substitution of wheat flour by pigeon pea flour (*Cajanus cajan*)

ABSTRACT

The present study aims to formulate and develop functional breads from wheat and pigeon pea flour and evaluate their nutritional and sensory qualities. Indeed, different composite bread formulations have been made by partial substitution of wheat flour with pigeon pea flour in varying proportions (10%, 20% 25% 50%), and the nutritional and sensory characteristics of the different breads produced were evaluated. Results indicated that the protein content of the composite breads significantly increases ($p < 0.005$) according to the pigeon pea flour incorporation rate. However, the evaluation of sensory characteristics indicated that only composite breads result from the incorporation rate of 10% of pigeon pea flours have organoleptic characteristics appreciated by the panel of tasters. These results therefore underline that the perception and opinion of consumers are very important parameters to be taken into account in any innovation process in the field of food technology.

Keywords: functional breads, pigeon pea, sensory profile, consumers perception, Benin

1. INTRODUCTION

Bread is described as a fermented product obtained from wheat flour, yeast, water and salt through a series of process involving mixing, kneading, checking, shaping and baking [1, 2]. Nowadays, bread consumption is very popular, but the low protein content of wheat flour has been a major concern in its utilization [3; 4]. Alam et al. [2] reported that the need of consumers to eat quality and healthy foods known as functional foods, which contains ingredients that can provide additional health benefits, is increasing nowadays. Therefore, researches should more focused on the production of specialty breads (functional foods) made from wheat flour and other functional ingredients such as proteins. In fact, according to the FAO [5], more than 850 million people suffer from undernourishment in the world. In sub-Saharan Africa, food problem arises in quantity and quality for more than a third of the population [6]. Faced with this situation, it has been suggested that the diversity of food from plants origin which abound in African countries could be a solution [7]. Unfortunately, the large part of this important resource is underexploited or neglected [8]. However, several researches reported that these plant resources have considerable potential that could increase food production and contribute to poverty reduction [9]. Among these plant species, legumes are nutritious foods that could replace proteins from animal origin [10]. Legumes are also source of dietary fiber, with high levels of vitamins and minerals [11]. Pigeon pea (*Cajanus cajan* (L.) is an important legume cultivated in the tropics, including semi-arid areas [12]. It is mainly grown for its grains whose nutritional value, is comparable to that of beans (*Phaseolus vulgaris*) [13]. It is also known as an excellent source of protein (21.7%), a good source of energy, vitamins and essential amino acids such as lysine, phenylalanine, valine, leucine and isoleucine [12]. The seeds are rich in fatty acids, such as linoleic and palmitic acids; and are also a good source of iron and calcium [14]. Then, the fortification of bread with legume flours particularly in regions where protein utilization is inadequate, could help reduce malnutrition and encourage farmers to grow more legume [15]. Then, the aim of this study was to formulate and develop functional breads from wheat flours and pigeon pea flour and to evaluate their nutritional and sensory qualities, as well as consumers overall acceptability.

2. MATERIAL AND METHODS

2.1. Raw materials and preparation of composite flours

Wheat flour, pigeon pea flour, sugar, iodized salt, yeast, spice and other general ingredients were purchased from the local market. The composite flour was prepared by using wheat and pigeon pea flours in different proportions as described in Table 1.

Table 1. Quantity of ingredients (g) used for different bread samples

Sample	Wheat flour	Pigeon pea flour	Salt	Sugar	Fat/Shorting	Yeast	Spice	Water
Control	100	00	2	6	4	2	1	65
Sample A	90	10	2	6	4	2	1	65
Sample B	80	20	2	6	4	2	1	65
Sample C	75	25	2	6	4	2	1	65
Sample D	50	50	2	6	4	2	1	65

2.2. Bread making

The method used for bread samples making is those described by Islam et al. [15] and Ndife et al. [16] as follow: wheat flour and pigeon pea flour were mixed and blended with other baking ingredients in a mixer. After kneaded into consistent dough, it was molded and placed in a pre-oiled baking bowl. The dough was after that proofed for 45 to 60 min at 35°C and 85% relative humidity and then baked in a reel oven for 35 min at 217°C.

2.3. Physicochemical and sensorial analyses

Physicochemical composition of the bread samples such as protein and ash contents were determined by methods described by AOAC [17]. 30 panelists carried out sensory evaluation of the composite bread samples on hedonic scale for different parameters such as colour, aroma, taste, texture and relative acidity as described by Ihekoronye and Ngoddy [18].

2.4. Statistical analysis

The data generated from these studies were analyzed using Statistical Analysis Software (SAS) and SYSTAT 5.05. The statistical analyses carried out were mean, standard deviation and analysis of variance.

3. RESULTS AND DISCUSSION

Table 2 presented the results of the protein and ash contents in the various functional breads produced. The analysis of these results indicated a significant variability ($p < 5\%$) at the protein and Ash contents level in the different breads produced. Indeed, the highest levels of protein and ash are obtained with the breads having the highest proportions of pigeon pea flour. This increase in the protein and ash contents in breads produced could be due to the protein and ash contents of the pigeon pea flour used. Similar results are reported by Ndife et al. [16] and Alam et al. [2] in breads obtained by partial substitution of wheat flour by soybean flour. Diallo et al. [19] also reported an increase in protein content in breads obtained from partial substitution of wheat flour by Voandzou flour (*Vigna subterranea* L. verdc).

Table 2. Protein and ash contents of different functional breads produced

Samples	Protein (%)	Ash (%)
A	15.75±1.06 ^a	3.81 ± 0.32 ^a
B	16.27±0.38 ^b	3.84 ± 0.22 ^a
C	17.22±1.38 ^c	3.91 ± 0.12 ^b
D	17.50±0.70 ^c	3.92 ± 0.31 ^b
Control	8.90±1.28 ^d	2.06±0.12 ^c

Values are mean ($n = 3$). The means followed by different superscript letters in the same column are significantly different according to ANOVA and Tukey's multiple comparison tests

The sensory analysis of the different types of bread produced revealed that the incorporation of pigeon pea flour modified the sensory characteristics such as color, aroma, texture, taste and acidity of the breads obtained, compared to the control (Figures 1-5). However, the effect of these sensory modifications is depending on the proportion of pigeon pea flour used. Only the "A" bread samples have sensory characteristics close to the control samples (Figures 4 and 5).

These results therefore show that, whatever the parameter considered (color, aroma, texture, taste and acidity), the breads obtained with pigeon pea flour incorporation rates of 25% and 50% presented poor organoleptic characteristics (pronounced color and aroma, more rigid texture, perceptible acidity), compared to the control. However, very pleasant acidity and texture were observed in the breads obtained with an incorporation rate of 10% pigeon pea flour. Similar results were reported by Olanipekun et al. [20] who pointed out that the incorporation of other types of flours in the manufacture of wheat bread affects the overall acceptability of the breads produced. The same remarks were reported by Ouazib [21] during the evaluation of the effect of the partial substitution of wheat flour by chickpea flour on bread quality.

The results of this study show that despite the high protein levels observed in the breads obtained from the incorporation rates of 50%, 25% and 20% pigeon peas, the results of the sensory analyses are not reassuring as to their consumer acceptability. Indeed, for a new product to be adopted, it must first be acceptable to consumers [22] and this acceptability must be assessed by taking into account sensory information and non-sensory information [23].

However, according to Masson [22], studying the acceptability of a new food product to place it on the market is not enough to predict its adoption, because the adoption of an innovation by consumers always involves a temporal dimension which makes its study more complicated. Thus, marketing research has most often involved unique and immediate measures of attitude, preference or purchase intention [24]. However, food science research has shown that immediate preferences are a weak predictor of final choice [25] and consumption behavior [26], because preference evaluations change over time [27]. Similarly, the different degrees of perceived novelty of the innovation can influence its evaluation and adoption by consumers [22]. Indeed, it has been shown that new products are tried more by people led by their own norms and values (individualists), than by people led by the values and norms of society (conformists). Thus, if the difference between innovation and the products of the category is large, the greater the difference in the answers between the conformists and the individualists will be. This explains the fact that slightly different innovations are adopted more quickly because the compatibility is greater. As a result, everyday consumer goods very often have a stronger purchase intention for products perceived as very new.



Figure 1. Sensorial profile of functional breads produced from partial substitution (50%) of wheat flour by pigeon pea flour




Figure 2. Sensorial profile of functional breads produced from partial substitution (25%) of wheat flour by pigeon pea flour




Figure 3. Sensorial profile of functional breads produced from partial substitution (20%) of wheat flour by pigeon pea flour




Figure 4. Sensorial profile of functional breads produced from partial substitution (10%) of wheat flour by pigeon pea flour




Figure 5. Sensorial profile of breads produced with wheat flour (Control)

4. CONCLUSION

Functional breads from wheat and pigeon pea flours were found to have a good nutritional profile. However, they presented poor organoleptic characteristics (pronounced color and aroma, more rigid texture, perceptible acidity), except for the breads obtained with an incorporation rate of 10% of pigeon pea flour with a very pleasant acidity and texture. These results therefore underline that the perception and opinion of consumers are very important parameters to be taken into account in any innovation process in the field of food technology. Nevertheless, in relation to the composition of breads, further investigations should be made to assess their fibre and carbohydrates contents.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- 1- Dewettinck K, Bockstaele FV, Kuhne B, Van WD, Courtens T, Gellynck X. Nutritional value of bread: Influence of processing, food interaction and consumer perception. *Rev. J. Cereal Sci.* 2008 ; 48 : 243 - 257.
- 2- Alam J, Talukder MU, Rahman MN, Prodhon UK, Obidul HAK. Evaluation of the nutritional and sensory quality of functional breads prepared from whole wheat and soybean flour. *Annals Food Science and Technology.* 2013 ; 14 (2) : 171-175.
- 3- Bakke A, Vickers Z. Consumer liking of refined and whole wheat breads. *J. Food Sci.* 2007; 72: S473-S480.
- 4- Jideani V, Onwubali F. Optimisation of wheat-sprouted soybean flour bread using response surface methodology. *Afr. J. Biotechnol.* 2009 ; 8 : 6364-6373.
- 5- FAO. L'état de l'insécurité alimentaire dans le monde. Rome 2002, 4p.
- 6- Kennedy G, Nantel G, Shetty P. The scourge of "hidden hunger": global dimensions of micronutrient deficiencies. *Food Nutrition and Agriculture.* 2003; 32: 8-16.
- 7- Chauvet L. Socio-political instability and the allocation of international aid by donors. *European Journal of Political Economy.* 2003; 19(1): 33-59.
- 8- Padulosi S, Hoeschle-Zeledon I. Underutilized plant species: what are they? *LEISA.* 2004; 20(1): 5-6.
- 9- Mayes SD, Calhoun SL, Aggarwal R, Baker C, Mathapati S, Anderson R, Petersen C. Explosive, oppositional, and aggressive behavior in children with autism compared to other clinical disorders and typical children. *Res Autism Spectrum Disord.* 2012; 6: 1-10.
- 10- Adebawale OJ, Maliki K. Effect of fermentation period on the chemical composition and functional properties of pigeon pea seed flour. *International Research Journal.* 2011; 18:1329-1333.
- 11- Almeida Costa GE, Da Silva Q.M.K., Pissini MRS., Costa De Oliveira A. Chemical composition, dietary fibre and resistant starch contents of raw and cooked pea, common bean, chickpea and lentil legumes. *Food Chemistry* 2006; 96: 327-330.
- 12- Wu N, Fu K, Fu YJ, Zu YG, Chang FR, Chen YH, Liu XL, Kong Y, Liu W, Gu CB. Antioxidant activities of extracts and main components of pigeon pea *Cajanus cajan* (L.) Millsp. leaves. *Molecule.* 2009; 14(3): 1032-1043.
- 13- Niyonkuru DN. La culture du pois cajan. *Saïd Cameroun* 2002. 23 p.
- 14- Fossou RK, Kouassi NK, Kouadjo GCZ, Zako SMIB, Zeze A. Diversité de *Rhizobia* dans un champ cultivé de pois d'angle (*Cajanus cajan* L.) (légumineuses) à Yamoussoukro (centre côte d'ivoire). *Agronomie Africaine.* 2012 ; 24 (1) : 29-38.
- 15- Islam T, Chowdhury A, Islam M, Islam S. Standardization of Bread Preparation from Soy Flour. *Int. J. Sustain. Crop Prod.* 2007 ; 2 : 15-20.
- 16- Ndife J, Abbo E. Functional Foods: Prospects and Challenges in Nigeria. *J.Sci.Technol.* 2009 ; 1 : 1-6.
- 17- AOAC. Official Methods of analysis. 15th ed. Association of Official Analytical Chemists 1990. Washington, DC.
- 18- Ihekoronye AI, Ngoddy PO. Integrated Food Science and Technology for the Tropics. 2nd ed. Macmillan Publishers Ltd 1985. London.
- 19- Diallo SK, Soro D, Kone KY, Assidjo NE, Yao K B, Gnagri D. Fortification et substitution de la farine de blé par la farine de Voandzou (*Vigna subterranea* L. verdc) dans la production des produits de boulangerie. *International Journal of Innovation and Scientific Research.* 2015 ; 18 (2) : 434-443
- 20- Olanipekun BF, Abioye VF, Oyelade OJ, Osemobor CO. Potentials of Pigeon Pea-wheat Flour Mixes in Bread Production. *AFSJ.* 2018; 4(2): 1-8.
- 21- Ouazib M. Effet de traitements sur les paramètres nutritionnels et fonctionnels du pois chiche produit localement : impact sur les propriétés rhéologiques, physicochimiques et sensorielles de pain à base de pois chiche. Thèse de Doctorat d'Université, Sciences biologiques, Sciences des aliments, Université A. MIRA-BEJAIA, 2017. 153p.

- 22- Masson J. Effets de la modification d'un attribut constitutif d'un produit alimentaire sur son adoption par les consommateurs : le cas du vin a teneur réduite en alcool. Thèse de Doctorat, Montpellier Supagro. Centre International d'Etudes Supérieures en Sciences Agronomiques, 2010, 286p.
- 23- Deliza R, MacFie HJH. The generation of sensory expectation by external cues and its effects on sensory perception and hedonic ratings: a review. *Journal of Sensory Studies*.1996 ; 11 (2) : 103-128.
- 24- D'Hauteville F. Un modèle d'acceptation du nouveau produit par le consommateur : cas du vin allégé en alcool", Thèse de Doctorat en Sciences de Gestion, Université Montpellier II. 1994. French.
- 25- Köster EP, Couronne T, Leon F, Levy C, Marcelino A S. Repeatability in hedonic sensory measurement: a conceptual exploration. *Food Quality and Preference*. 2002 ; 14 (2) : 165-176.
- 26- Vickers Z, Holton E. A comparison of taste test ratings, repeated consumption, and post consumption ratings of different strengths of iced tea, *Journal of Sensory Studies*. 1998; 13 (2): 199-212.
- 27- Chung S-J,Vickers Z. Long-term acceptability and choice of teas differing in sweetness, *Food Quality and Preference*. 2007; 18 (7) : 963-974.