

The potential of piperidine alkaloids-rich extracts from *Prosopis* spp. to combat plant pests and diseases

ABSTRACT

The genus *Prosopis* comprises 44 species of spiny small trees adapted to arid and semiarid regions of the planet that have been used by human populations as a source of wood, food and medicine. *Prosopis* spp. possess high quantities of phenolic compounds and piperidine alkaloids with antioxidant, neurotoxic, antimicrobial, antimalarial, insecticidal and allelopathic activities. Juliprosopine is the major and more characterized piperidine alkaloid present in the genus *Prosopis*. In this review we present the potential of *Prosopis* spp. extracts and piperidine alkaloids for the control of plant pests and diseases and discuss the possibility of their use as an alternative in organic agriculture.

Keywords: Prosopis spp. extracts; piperidine alkaloids; phytopathogen; plant diseases.

1. INTRODUCTION

Since agriculture emerged, weeds, insects, and microorganisms, which co-evolved with plants in the ecosystem, have turned out in plant pests and diseases. The ability of pathogenic microorganism to colonize the host have increased with the domestication, and systematic-extensive monoculture of plants. Today, these pests and diseases have become a threat to our-food-production safety. This is due to the natural phenomena of resistance present in weeds [1], insects [2], bacteria [3], and fungi [4] as a consequence of the indiscriminate application of bactericides, fungicides and pesticides agents (frequently mixed and applied together) in crops over time [5, 6]. All these practices have conducted to the creation of global programs oriented to reducing and making rational use of synthetic biocides in plant pests and diseases control, such as the Integrated Pest Management (IPM, FAO), the International Plant Protection Convention (IPPC, FAO) and the Fungicide Resistance Action Committee (FRAC).

Natural products emerge as an alternative to the control of insects and phytopathogenic microorganisms with minimal residuality and low environmental impact [7]. In addition, natural biocides appear as the main option in the growing organic agriculture market (USDA Organic). According to the Research Institute of Organic Agriculture (FiBL), 71.5 million hectares were organically managed in 2018, and the offer for biopesticides is increasing every year (OMRI). In this respect, plants represent a vast source of new compounds with biocidal properties due to they have extraordinary ability to synthesizing organic molecules to cope with phytopathogens [5, 8].

This review provides a summary of the available biological activities of the extracts of *Prosopis* spp., such as antibacterial, antifungal, insecticide and allelopathic activity, as well as information about the bioactive compounds that have been identified in the genus.

2. CHARACTERISTICS FEATURES OF *Prosopis* spp.

The genus *Prosopis* belongs to the family Leguminosae (Fabaceae), subfamily Mimosaceae, and is composed of 44 species usually distributed in arid, tropical and subtropical regions from Asia, Africa, and the Americas [9-11]. Members of *Prosopis* spp. have been used with different purposes such as; wood as a source of fuel; the pods as a source of carbohydrates and protein for livestock and human populations, and the leaves, seeds, and gum as a medicinal source [12-14]. Ecologically, *Prosopis* spp. are evergreen xerophyte trees or shrubs well adapted to arid zones that spread and multiply easily, and they might become invasive species in the places where they are introduced if not properly controlled [15, 16]. *P. juliflora* is the most studied species from a pharmacological and phytochemical point of view. Its alkaloids display a wide range of biological activities [15, 17-20]. Nevertheless, the emphasis on *Prosopis* spp. extracts as a potential source of biopesticides has not been recently reviewed. Piperidine alkaloids have also been identified in leaves, pods, roots, and flowers of other *Prosopis* species, such as *P. glandulosa* [21, 22], *P. ruscifolia* [23], *P. africana* [24], *P. affinis* [11], *P. alata*, *P. argentina*, *P. chilensis*, *P. flexuosa* and *P. pugionata* [9]. Two kinds of piperidine alkaloids could be found in *Prosopis* spp. based on presence or absence of an indolizidine ring in their structures.

2.1 *Prosopis juliflora*

Prosopis juliflora is an evergreen shrub or tree commonly called “mesquite” or “algarroba”, originally from America and nowadays introduced and widespread in many countries around the world. *P. juliflora* is an important species in semi-desert areas, playing an important role as shelter, improving soils via nitrogen fixation, increasing organic matter via leaf litter addition, changing soil structure and microbial populations. Its flowers produce a large amount of pollen as such as they are highly melliferous [12, 25, 26]. For humans, it is used as a revegetation species [27], as a source of wood for fuel and construction [28] and the pods are used as cattle food [29]. In the countries where *P. juliflora* is introduced, it can become an insidious species [16, 17] due to the allelochemicals present in the different tissues that can change the native ecosystem structure [14]. In traditional medicine, the leaves and bark have been employed to treat cold, sore throat, inflammation, digestive problems, skincare and wound healing [30-32]. Its extracts displayed antioxidant [26, 31, 33], antimalarial [34], antibacterial [35-37], antifungal [30, 38], antimites [39], and herbicidal [40] activities. Also, the pharmacological properties of its extracts, alkaloid fractions and compounds have been extensively reviewed [13, 14, 41-43].

2.2 *Prosopis africana*

P. africana, commonly known as African mesquite, is the only *Prosopis* species that grows in the African continent. It is a small to large tree with thornless twigs and a fast-growing root system [44]. The pods and leaves have been used as a source of animal food due to their high protein content; the wood, as a source of charcoal, construction poles, and wooden tools; and the roots and bark as a source of medicine for human ailments, such as toothache, [20, 45, 46], gingivitis and periodontal disease treatment [24], as well as wound healing [47]. Ecologically is an important soil fixing nitrogen plant [46]. The stem, bark and root extracts have shown several biological activities, such as antibacterial, antifungal [24, 47, 48] and antimalarial activity [44]. Some studies have described the presence of alkaloids and phenolic compounds in this species [24, 49].

Unlike other *Prosopis* species, *P. africana* has a low germination rate of seeds and low fruit production [45]. Its use has been so extensive, that in some regions of Africa it is a threatened species.

2.3 *Prosopis glandulosa*

Prosopis glandulosa, popularly known as honey mesquite, is a medium sized tree with spiny branches native from the north of Mexico to the south of EE.UU [50]. *P. glandulosa* has been employed as firewood and in the traditional medicine for eye infections, open wounds, dermatological and stomach ailments. There exist two varieties of *P. glandulosa*; var. *glandulosa* and var. *torreyana*. The former is present in the more humid areas toward Gulf of Mexico and the latter prefers the drier climate near the Pacific. Anatomically, *P. glandulosa* var. *glandulosa* present larger leaves than var. *torreyana* and there exist thorny populations of *P. glandulosa* var. *glandulosa* that can be classified as var. *prostrata* [10]. The polar extracts (ethanolic and aqueous) of the leaves have displayed antifungal activity and the alkaloid rich fraction derived of the extracts recorded higher activity [21, 22, 50]. Besides, the aqueous extract has been used to generate silver nanoparticles with antibacterial and antifungal activity [51, 52].

2.4 *Prosopis ruscifolia*

Prosopis ruscifolia, also known as “vinal” in South America, is a thorny tree native to northern Argentina [23]. The methanolic bark extract exhibits antibacterial activity against *Staphylococcus aureus* (ATCC 8095 and INEI 2213), *Enterococcus faecium* (INEI2464), *Klebsiella pneumoniae* (ATCC 10031) [53] and the methanolic leaf extract shows antifungal activity. The antifungal components were identified as juliflorine (**1**), prosopiflorine (**4**), cyclitol and a triterpene saponin [54, 55]. Indolizidine alkaloids have been detected *in situ* in the cuticle and palisade mesophylls of the leaflets, as well as in petiole and rachis parenchyma. On the other hand, in pods, are present in the cuticle, epidermal cells, mesocarp parenchyma cells and the phloem [23].

3. CHEMICAL CONSTITUENTS OF *Prosopis* spp.

3.1 Stem bark

Characterization of the bark extracts from *P. juliflora* have revealed the presence of high content of flavonoids, as 4'-O-methyl-gallocatechin, (+)-caofechins and (-)-mezquitol, the latter is the most abundant [56, 57]. Also, the 7,3',4'-trihydroxy-3-methoxyflavanone was identified in the *P. africana* bark [49]. Additionally, *Prosopis* spp. bark contains alkylated piperidine alkaloids (Fig. 1). [58] isolated the N-methylcassine (**12**) from methanolic bark extract of *P. nigra*, *P. ruscifolia* and *P. vinalillo* through an acid-base extraction followed of preparative Thin Layer Chromatography (TLC). Besides, [11] isolated by first time and structurally characterized the N-methyl-2-isocassine (**12b**), N-methyl-6-isocassine (**12c**), and N-methyl-6-isocarnavaline (**13**) from *P. affinis* bark, using mass spectrometry (MS), infra-red (IR) and nuclear magnetic resonance (NMR) data. In addition, juliprosopine (**1**) and juliprosine (**2**) have been found in *P. flexuosa* exudate [9].

3.2 Flowers and Pollen

The flowers of *Prosopis* spp. are highly melliferous since they produce an enormous pollen quantity. Phytochemical studies using Dragendorff's reagent and direct analysis in real time (DART)-MS of ethanolic flower extract from *P. juliflora*, showed the presence of the alkaloids julifloridine (**7**), prosopine (**5**), prosopinine (**6**) and prosafrinine (**10**) (Fig.1) [59]. Also, a wide

variety of phenolic compounds have been detected in the hydroethanolic pollen extract, such as apigenine, luteolin and cinnamic acid derivatives, as well as flavonol, quercetin and genistein glycosides [26], making *P. juliflora* pollen an interesting source of natural antioxidant compounds and alkaloids.

3.3 Root

[21] identified the alkaloids 3-isocassine (**11**), N-methyl-3-isocassine (**12a**) and N-methylcassine (**12**) from the ethanolic root bark extract of *P. glandulosa* var. *torreyana* using IR spectra and ¹H- ¹³C NMR (Fig. 1). Also, alkaloids have been detected in the methanolic, ethanolic and aqueous root extract of *P. africana* with Dragendorff's, Mayer's and Wagner's reagents [24, 44].

3.4 Leaves

Most of the phytochemical studies have been carried out in the aerial parts of *Prosopis* spp. In *P. juliflora* leaves extracts have been detected alkaloids, flavonoids, tannins, saponins, anthraquinones, and coumarins [31, 33, 37, 41, 43, 60-62]. Similarly compounds that are present in the leaves are also present in the *P. juliflora* spines. [32] reported the presence of flavonoids, saponins, glycosides, terpenoids and coumarins in the aqueous and ethanolic thorn extracts by colorimetric methods, and alkaloids were found only in the aqueous extract. Studies of the relative abundance reveal that flavonoids (16%), pectin substances (4.9%) and alkaloids (3.6%) are the three main compounds in *P. juliflora* leaves [41].

Alkaloids have been found in *P. alba* [63], *P. laevigata* [64], *P. glandulosa* [22], *P. ruscifolia* [23], *P. alpataco*, *P. argentina*, *P. chilensis*, *P. flexuosa* and *P. pugionata* [9]. *Prosopis* spp. alkaloids have attracted more research interest than all other secondary metabolites detected in *Prosopis* spp. leaves due to their pharmacological and biological activities [41].

Characterization of *P. juliflora* leaf alkaloids through spectroscopic and spectrometric techniques has been carried out. The analysis of MS, IR as well as ¹H and ¹³C NMR spectra revealed the structure of juliflorine (juliprosopine) (**1**) and their stereoisomers juliflorinine (**1a**) and julifloricine (**1b**; **1c**), as well as the jullifloridine (**7**) and the juliprosinene (**3**) [65, 66]. Using the same techniques, [67] elucidated the juliprosine (**2**) and the isojuliprosine (**2a**) structures. Later, [68] reported the structures of secojuliprosopinal (**8**), 3''''-oxo-juliprosopine (**14**), 3-oxo-juliprosine (**15**) 3'-oxo-juliprosine (**16**) using Electrospray ionization (ESI)-MS and ¹H-¹³C NMR and julifloravizole was described by ¹H-¹³C NMR correlations [69]. In recent years, [35] and [59, 70] identified the alkaloids prosopiflorine (**4**), prosopine (**5**), prosopinine (**6**), projuline (**20**), prosafrinine (**10**) and N-methyl julifloridine (**9**) with DART-MS. The juliprosopine (**1**) was also identified in the aerial parts of *P. pugionata* using NMR and MS techniques [9]. [22] isolated and identified by IR, ESI-HRMS and ¹H-¹³C NMR the prosopilosidine (**18**) and their stereoisomers isoprosopilosidine (**18a**, **18b**) as well as the prosopilosine (**19**) and their stereoisomers isoprosopilosine (**19a**, **19b**), all compounds from the ethanolic leaf extract of *P. glandulosa* var. *glandulosa*. Tryptamine and juliprosopine (**1**) were also present, the latter is the representative alkaloid of the genus *Prosopis* [21]. In addition, the alkaloids prosopine (**5**), prosopinine (**6**) and its stereoisomer prosophylline (**6a**), prosafrinine (**10**) and prosafrine (**20**) were isolated from *P. africana* leaves (Fig. 1) [71].

3.5 Pods

The *Prosopis* spp. pods have been used as a source of protein to feed cattle and human populations, nevertheless, if they are used as the only source of nourishment, could provoke

a neurotoxic disease called “twisted face” or “cara torta”, especially in ruminant cattle [72, 73]. Nevertheless, *Prosopis* spp. pods are a rich source of polyphenolic compounds with antioxidant activity and a high variety of piperidine alkaloids [20]. In *P. juliflora* pods have been detected the same alkaloids that those found in the leaves but in different concentrations. The relative abundance with DART-MS in *P. juliflora* showed that the major alkaloid in leaves is juliprosopine (1) followed by juliprosine (2), both alkaloids with indolizidine moiety; while in pods and flowers the most abundant is julifloridine (7) followed by prosopinine (6) belonging to the group of alkaloids without indolizidine moiety [35, 59]. It is important to point out that prosopinoline (21) was only detected in pods, while N-methyl julifloridine (9) was found only in the leaves [59]. In *P. flexuosa* pods, juliprosine (2) and juliprosopine (1) are the major alkaloids according with HPLC-HRMS analysis [72] (Fig. 1).

In addition to alkaloids, *Prosopis* spp. pods are a rich source of polyphenolic compounds. *P. alba* pods contain Q-dihexoside rhamnoside, vitexin and isovitexin, and in *P. nigra* pods are present anthocyanins like cyanidin rhamnosyl hexoside, peonidin-3-hexoside, malvidin dihexoside and vicianin II [20]. The total phenols and flavonoid contents in *P. farcta* pods were estimated as 61.5 mg gallic acid equivalent and 17 mg quercetin equivalent per gram of dry plant material, respectively. The major volatile constituents are the palmitic acid, 9,12-octadecadienoic acid ethyl ester, cembrene A, myristic acid, farnesyl acetate, nonanal and α -terpinyl acetate [74].




Fig. 1. Piperidine alkaloids from *Prosopis* spp.

4. BIOLOGICAL ACTIVITIES OF *Prosopis* spp. EXTRACTS AND PHYTOCHEMICALS

Plants are adapted to naturally produce secondary metabolites to deal with biotic and abiotic stress. Compounds like terpenes, flavonoids, phenols, and alkaloids are responsible for this plant's chemical defense to cope with adverse conditions. Phytochemicals are biologically active and the antimicrobial metabolites are especially interesting since they could reduce the antimicrobial resistance (AMR) effect with a low environmental residually [8, 75].

4.1 Antifungal activity

Despite plants having naturally developed a defense mechanism to avoid fungal diseases, crops around the world are infected by several phytopathogenic fungi that overcome those defenses causing plant disease, low productivity, and crop losses. Synthetic fungicides are currently used to reduce this problem with the disadvantages of resistance phenomena [4]. In this context, *Prosopis* spp. derived compounds and extracts could represent another natural alternative to reduce the use of synthetic fungicides to the control of the pathogenic fungi. [48] showed that the aqueous bark extract of *P. Africana*, and to a lesser extent the aqueous leaf extract, inhibited the *in vitro* mycelial growth and reduced the sporulation and spore length of *Curvularia lunata* isolated from *Jatropha curcas* (Table 1). [76] used aqueous extract of *P. juliflora* leaves (2% w/v) to the control late leaf spot (LLS; *Phaeoisariopsis personata*) and rust (*Puccinia arachidis*) in groundnut (*Arachis hypogaea*) under greenhouse conditions and in the field. Foliar application of *P. juliflora* leaf extract reduced foliar disease severity in the greenhouse, whereas the combination of the extracts with chlorothalonil increased the pod yields by 81-98% in the field. This strategy reduced three times the use of chlorothalonil and allowed the management of these diseases in field conditions. [77] tested seven commercial fungicides amended with the *P. juliflora* alkaloid fraction against the seed-borne *Fusarium verticillioides* isolated from maize, and the use of the alkaloid fraction significantly reduced the percentage of the synthetic fungicides to reach the same mycelial inhibition as that produced by the highest concentration of the pure fungicide. The application of *P. juliflora* aqueous extracts to seeds and seedlings grown in pots showed potential to reduce root infecting fungi (*Fusarium* spp., *Rhizoctonia solani*, and *Macrophomina phaseolina*) and improves germination in cowpea (*Vigna unguiculata*) and mungbean (*Vigna radiata*) [78]. Furthermore, the leaf powder added to the soil was effective in controlling the root rot disease in okra plants [79] (Table 1).

In addition, aqueous extract of *P. glandulosa* leaves at 5% completely inhibited the radial growth of *C. gloeosporioides* and dramatically reduced to *F. oxysporum*, *R. oryzae*, *R. stolonifera* (Table 1). At this concentration, the extract also decreased the disease symptoms (necrosis, soft rot, dehydration) produced by *C. gloeosporioides*, *F. oxysporum*, *R. oryzae*, and *R. stolonifer* on strawberry, cherry tomato, and baby carrot [50]. *Colletotrichum* spp. is an important genus of plant pathogens that generates pre and post-harvest losses in a wide variety of crops [80, 81]. Resistance reports to the benzimidazoles fungicides benomyl, thiabendazole, and thiophanate methyl had been noticed in *C. musae* isolated from banana and *C. gloeosporioides* from mango [82, 83]. Additionally, *C. gloeosporioides* resistance to the post-harvest application of imazalil fungicide has been reported in mango [84]. Also, isolated strains of *C. cereale* and *C. coffeanum* had presented benzimidazole resistance, and there exist reports of *C. graminicola* resistant to strobilurin related to QoI fungicides. [38] tested the *in vitro* antifungal activity of methanolic leaf extracts (10% w/v) of several plant species against *C. musae* isolated from banana, and *P. juliflora* extract displayed the highest inhibitory effect on conidial germination and mycelial growth. In a similar way, [85] reported the *in vitro* antifungal activity of methanolic and acetone extract (10-20% w/v) from *P. juliflora* leaves against *C. gloeosporioides* isolated from mango, as well as a reduction in the

anthracnose disease symptoms on mango fruits (Table 1). In addition, the aqueous extract (24% w/v) and the alkaloid rich fraction from *P. juliflora* leaves inhibited in vitro radial growth of *Alternaria alternata* isolated from tobacco plants [69].

Finally, the methanolic extract of the aerial parts (leaves and limbs) of *P. ruscifolia* displayed antifungal activity against *Aspergillus fumigatus*, *A. niger*, *A. nomius*, *A. flavus*, and *A. parasiticus*. Further analysis of this extract revealed the presence of alkaloids, such as juliprosopine (**1**) and juliprosinene (**3**) (Fig. 1); pinitol and a triterpene saponin [54]. The juliprosopine (**1**) and prosopilosidine (**18**) isolated from *P. glandulosa* var. *glandulosa* showed in vitro antifungal activity against *Cryptococcus neoformans* (ATCC90113) and *A. fumigatus* (ATCC 90906), respectively [22, 86].

The potential of *Prosopis* spp. extracts or chemical constituents to the control of phytopathogenic fungi could be potentiated through the formulation of nanoparticles. Silver nanoparticles (AgNPs) obtained from aqueous leaf extract of *P. glandulosa* inhibited the in vitro and in vivo growth of *F. solani* in cotton plants [52], and zinc oxide nanoparticles (ZnONPs) synthesized with aqueous extract from *P. farcta* pods exhibited antifungal activity against 10 clinical isolates of *C. albicans* [87].

4.2 Antibacterial activity

The inhibitory effect of *Prosopis* spp. extracts against plant pathogenic bacteria is not as wide as that showed against fungi. In vitro antibacterial activity from leaves and root alcoholic extracts of *P. juliflora* have been reported against *Pseudomonas aureginosa* at doses of 10% (w/v) [60, 88, 89]. [90] also reported in vitro antibacterial activity of aqueous leaf extract against *Xanthomonas campestris* pathovars *malvacearum*, *phaseoli*, and *vesicatoria* from cotton, French bean and tomato, respectively. Also, the *P. juliflora* methanolic extract inhibited the growth of *X. campestris* [91, 92], and *Agrobacterium rhizogenes* [92]. Other studies have shown the antibacterial activity of the *P. juliflora* leaf methanolic extract [36], the *P. juliflora* alkaloid fraction [73] and the *P. farcta* pod ethanolic extract against human pathogenic bacteria, suggesting that *Prosopis* spp. extracts could display activity against many other phytopathogenic bacteria. Furthermore, the AgNPs from *P. glandulosa* leaf extract have potential antibacterial activity as well [51].

4.3 Insecticide activity

Plants produced secondary metabolites to cope with the frequent interactions with the environment, including herbivores (insects, nematodes, vertebrates) [5]. Insects cause major crop losses that have increased since the green-revolution [2]. In a similar way as in fungi, bacteria, and weeds, pesticide resistance emerged as early as the synthetic insecticide control was massively implemented [93], this was accompanied by the residual toxicity and the non-target organism effect. The hexane leaf extract of *P. juliflora* was lethal against *Macrotermes* spp. at 25, 50, 75 and 100 mg L⁻¹. The mortality percentage increased with the concentration in a time dependent manner [62]. In nursery conditions, the aqueous leaf extract is efficient in the control of *Tetranychus bastosi* on *Jatropha curcas* at LC50 and LC90 values of 53.4% and 85.3%, respectively. Low residuality of the extract and absence phytotoxic effects were observed in *J. curcas* plants [39]. Aqueous leaf extract applied in the field can restrain the population of the green leafhopper (*Nephotettix virescens*), that transmits the rice tungro virus (RTV) in rice plants [94]. [95] reported a deterrent effect of the aqueous and ether extracts from leaves, bark, and fruits of *P. juliflora* against *Callosobruchus chinensis* beetle. The ether leaf extract (10% w/v) displayed the highest adult mortality percentages (66.6%), the smoke treatment from fruit and leaves powder induced 70% of mortality while the aqueous bark extract (10%) reduced the egg laying. The

methanolic seed extract from *P. juliflora* increased the mortality of *Helicoverpa armigera* larvae from 10 to 80% at 1 and 80 mg L⁻¹, respectively. The combination of seed extract and spore-crystal mixture of *Bacillus thuringiensis* kurstaki had a synergistic effect increasing larvae mortality and decreasing adult longevity and fecundity [96].

4.4 Allelopathic activity

The *P. juliflora* extracts also displayed allelopathic activity that could be exploited to control weeds in some crops. The aqueous extracts (25%, p/v) of leaves, bark, and roots from *P. juliflora* reduced the seedling emergence of *Mimosa tenuiflora* around 30%. Additionally, the leaves extract also reduced the seedlings height and stem diameter [97]. Also, [98] reported the reduction in the germination rate of *Cynodon dactylon* seeds and the radicle length in seedlings with the exposition of aqueous leaf extract from *P. juliflora* (10-60 g L⁻¹) in a concentration dependent manner. [40] assessed the application of aqueous extracts from *P. juliflora* leaves, steam, and bark on wheat plants (*Triticum aestivum*) cultivated in the field to the control of weeds. All extracts reduced the weeds density 60 days after sowing, but the wheat yield and quality parameters were reduced as the extract concentration increased (up to 40%). This pointed out that *P. juliflora* extracts cannot be used in all kinds of crops, and it is necessary to establish the maximum doses to inhibit the growth of the weed without affecting the growth of the crop. [68] tested the effect of some alkaloids isolated from *P. juliflora* leaves, such as secojuliprosopinal (**8**), 3'''-oxo-juliprosopine (**14**), and the mix 3-oxo-juliprosopine (**16**)/3'-oxo-juliprosopine (**15**) on seeds germination and seedlings growth of *Lepidium sativum* (Fig. 1). All alkaloids inhibited the shoots and roots growth, whereas the mixture of 3-oxo-juliprosopine/3'-oxo-juliprosopine displayed the strongest activity on shoots (IC₅₀: 200 µM) and roots (IC₅₀: 100 µM). These results suggest that both piperidine alkaloids with and without indolizidine ring are responsible for the allelopathic activity of the *Prosopis* spp. extracts.

Table 1. Antimicrobial, insecticide and allelopathic activities from *Prosopis* spp. tissues, extracts and alkaloids.

Tissue and extracts	Activity	Assays	Ref
<i>Prosopis africana</i>			
Aqueous bark extract	Antifungal against:	In vitro	[48]
Aqueous leaf extract	<i>Curvularia lunata</i> isolated from <i>Jatropha curcas</i>	Mycelial growth by food poison method	
<i>Prosopis juliflora</i>			
Leaf powder; alone and mixed with <i>Verticillium chlamydosporium</i> , <i>Paecilomyces lilacinus</i> , and <i>Bacillus subtilis</i>	Antifungal against: <i>Macrophomina phaseolina</i> , <i>Rhizoctonia solani</i> , and <i>Fusarium solani</i>	In vivo	[79]
Aqueous leaf extract and methanolic leaf extract	<i>Colletotrichum musae</i> isolated from banana	In vitro	[38]
		Mycelial growth by disk diffusion method and conidia germination	
Aqueous leaf extract, methanolic leaf extract and acetone leaf extract	Antifungal against: <i>Colletotrichum gloeosporioides</i> isolated from mango	In vitro	[85]
		Mycelial growth by food poison method and conidia formation	
		In vivo	
		Disease severity on mango (<i>Mangifera indica</i>) fruits infected	

Aqueous leaf extract and aqueous leaf extract + chlorothalonil	Antifungal against: <i>Phaeoisariopsis personata</i> and <i>Puccinia arachidis</i> from groundnut	In vitro Conidial germination In vivo Disease severity in groundnut (<i>Arachis hypogaea</i> cv. TMV 2) plants on greenhouse and field	[76]
Successive extraction: Petroleum ether leaf extract (N/A), hexane leaf extract (N/A), chloroform leaf extract (N/A), methanolic leaf extract Combination of methanolic leaf extract or alkaloid fraction + fungicide (Copper oxychloride, methyl-thiophanate, chlorothalonil, carbendazim and strobilurins) Aqueous leaf extract Aqueous stem extract Aqueous flowers extract	Antifungal against: Seed-borne <i>Fusarium verticilloides</i> from maize	In vitro Mycelial growth by food poison method	[77]
Aqueous leaf extract Aqueous stem extract Aqueous flowers extract	Antifungal against: <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> , and <i>Macrophomina phaseolina</i>	In vivo Disease severity in Cowpea (<i>Vigna unguiculata</i>) and Mungbean (<i>Vigna radiata</i>) seeds and plants in greenhouse	[78]
Aqueous leaf extract Successive extraction: Petroleum ether extract (N/A), benzene extract (N/A), chloroform extract (N/A), ethanolic extract, methanolic extract Alkaloid fraction	Antifungal against: <i>Alternaria alternata</i> from tobacco	In vitro Mycelial growth by food poison method	[69]
Ethanol leaf extract	Antibacterial against: <i>Pseudomonas</i> sp., <i>P. aeruginosa</i> , <i>Salmonella typhimurium</i> , <i>Klebsiella pneumonia</i> , <i>E. coli</i> , <i>Staphylococcus epidermis</i> , <i>S. aureus</i> , <i>Micrococcus luteus</i> , <i>Streptococcus</i> sp., <i>B. subtilis</i> .	In vitro Agar disk diffusion method	[60]
Methanol, ethanol and butanol leaf extract Methanol leaf extract, petroleum spirit fraction (N/A), chloroform fraction, ethyl acetate fraction, aqueous fraction	Antibacterial against: <i>P. aeruginosa</i> , <i>B. subtilis</i> , <i>Enterococcus faecalis</i> , <i>E. coli</i> , <i>K. pneumonia</i> , <i>Salmonella typhi</i> , <i>S. typhimurium</i> , <i>S. aureus</i> , <i>S. epidermidis</i> , <i>Streptococcus pyogenes</i> .	In vitro Agar disk diffusion method	[88]

80% Ethanol leaf extract 80% Ethanol root bark extract	Antibacterial against: <i>P. aeruginosa</i> , <i>E. coli</i>	In vitro Agar disk diffusion method	[89]
Aqueous leaf extract	Antibacterial against: <i>Xanthomonas campestris</i> pvs. <i>malvacearum</i> , <i>phaseoli</i> and <i>vesicatoria</i> from cotton, french bean and tomato, respectively.	In vitro Agar well diffusion method	[90]
Methanol, ethanol, ethyl acetate (N/A), hexane, chloroform, dichloromethane and petroleum ether leaf extracts.	Antibacterial against: <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i> from tomato	In vitro Agar disk diffusion method	[91]
Aqueous leaf extract	Antibacterial against: <i>Xanthomonas campestris</i> , <i>Agrobacterium rhizogenes</i> and <i>Aspergillus fumigatus</i>	In vitro Agar well diffusion method	[92]
Methanol leaf extract	Antibacterial against: <i>Klebsiella</i> sp., <i>S. aureus</i> , <i>Streptococcus</i> sp., <i>Bacillus</i> sp., <i>E. coli</i> .	In vitro Agar well diffusion method	[36]
Hexane leaf extract	Insecticidal against: <i>Macrotermes</i> spp.	In vitro No-choice bioassay method	[62]
Aqueous leaf extract	Insecticidal against: <i>Tetranychus bastosi</i>	In vitro No-choice bioassay method In vivo Counting of adult females and control efficiency on the fecundity and residual efficiency on the fecundity and mortality of adult females in <i>Jatropha curcas</i> plants under nursery conditions	[39]
Aqueous leaf extract	Insecticidal against: <i>Nephotettix virescens</i>	In vivo Counting population of <i>N. virescens</i> and percentage of rice tungro disease incidence in nursery and field	[94]
Aqueous leaf extract Ether leaf extract Leaf powder Leaf smoke	Insecticidal against: <i>Callosobruchus chinensis</i>	In vitro Control efficiency on the egg laying, adult emergence, rate of development and adult mortality in <i>Phaseolus mungo</i> grains	[95]
Methanol seed extract	Insecticidal against:	In vitro	[96]

Methanol seed extract + <i>Bacillus thuringiensis</i> <i>kurstaki</i> (Btk) HD1 strain	<i>Helicoverpa armigera</i> from non Bt cotton fields	Lethal concentration (LC ₅₀), antifeedant bioassay, control efficiency on insect development	
Aqueous leaf extract Aqueous bark extract Aqueous root extract	Allelopathic against: <i>Mimosa tenuiflora</i>	In vitro Seedlings emergency percentage, stem height and diameter, leaves number and dry matter weight of <i>Mimosa tenuiflora</i> seedlings	[97]
Aqueous leaf extract	Allelopathic against: <i>Cynodon dactylon</i>	In vitro Seedlings germination percentage and early seedling development	[98]
Aqueous leaf extract Aqueous stem extract Aqueous root extract	Allelopathic against: Wheat (<i>Triticum aestivum</i>)	In vivo Weed density, fresh weed biomass, dry weed biomass, chlorophyll content, leaf area index, leaf area duration, crop growth rate, net assimilation rate, plant height and number of tillers, spike length, number of grains, 1000-grain weight, grain yield, biological yield, harvest index and grain protein content	[40]
Methanol leaf extract Alkaloid fraction (3''''-oxo-juliprosopine, secojuliprosopinal and 3-oxo-juliprosine/3'-oxo-juliprosine) <i>Prosopis ruscifolia</i>	Allelopathic against: <i>Lepidium sativum</i>	In vitro Determination of roots and shoots growth and IC ₅₀	[68]
Leaf and steam extracts Successive extraction: Hexane extract (N/A), dichloromethane extract (N/A), ethyl acetate extract (N/A), methanol extract Fraction with juliflorine, juliprosinene, pinitol, and a triterpene saponin <i>Prosopis glandulosa</i> Aqueous leaf extract	Antifungal against: <i>Aspergillus fumigatus</i> , <i>A. niger</i> , <i>A. nomius</i> , <i>A. flavus</i> , and <i>A. parasiticus</i>	In vitro Mycelial growth by bioautography and plate microdilution	[54]
	Antifungal against: <i>Colletotrichum gloeosporioides</i> , <i>Fusarium oxysporum</i> , <i>Rhizopus oryzae</i> and <i>R. stolonifer</i>	In vitro Mycelial growth by food poison method In vivo Disease severity in strawberry (<i>Fragaria x ananassa</i>), cherry, tomato	[50]

Ethanol leaf extract; Alkaloid fraction; juliprosopine Ethanol root extract; Alkaloid fraction; 3- isocassine (N/A), N-methyl- 3-isocassine (N/A), N- methylcassine(N/A) Ethanol leaf extract; Alkaloid fraction; prosopilosidine, prosopilosine, isoprosopilosine, isoprosopilosidine, juliprosopine Nanoparticles from the aqueous leaf extract	Antifungal against: <i>Cryptococcus neoformans</i> (ATCC90113), <i>A. fumigatus</i> (ATCC 90906)	(<i>Solanum lycopersicum</i> var. <i>cerasiforme</i>) and baby carrot (<i>Daucus carota</i>) In vitro Mycelial growth by plate microdilution	[21]
	Antifungal against: <i>C. neoformans</i> (ATCC90113), <i>A. fumigatus</i> (ATCC 90906)	In vitro Mycelial growth by plate microdilution In vivo Male mice (Swiss Webster strain) infected with <i>C.</i> <i>neoformans</i>	[22]
	Antifungal against: <i>Fusarium solani</i> (T-ICA04) from transgenic insect- resistant cotton	In vivo Determination of physiological and morphological parameters and disease severity in Bollgard® cotton transgenic seedlings	[52]

N/A; no activity.

5. CONCLUSION

Prosopis spp. are multipurpose trees distributed worldwide. The extracts and piperidine alkaloids from these species have demonstrated antimicrobial activity against fungal and bacterial phytopathogens, as well as insecticide activity against causal agents of plant pest and diseases. Therefore, it is important to highlight the potential of phytochemicals produced by these tree species as an alternative to decrease the use of synthetic biocides and the environmental pollution related to their excessive application. Although the synergism of *Prosopis* spp. extracts with fungicides or beneficial microorganisms allows the efficient use of compounds with biocidal activity, it is necessary to carry out studies in field conditions to validate the results reported under *in vitro* and greenhouse conditions.

In the market, there exist a wide variety of plant derived biocides and their demand is increasing since the spread of the organic agriculture and the necessity of an integrated pest management for a healthy agriculture. Hence, the evaluation of pure compounds and the standardization of extracts based on bioactive compounds are imperative to the formulation of biocides products derived from *Prosopis* spp.

REFERENCES

1. Burgos NR, Tranel PJ, Streibig JC, Davis VM, Shaner D, Norsworthy JK et al. Review: Confirmation of resistance to herbicides and evaluation of resistance levels. *Weed Sci.* 2013;61:4-20. DOI: <https://doi.org/10.1614/WS-D-12-00032.1>
2. Khan MA, Khan Z, Ahmad W, Paul B, Paul S, Aggarwal C, Akhtar MS. Insect pest resistance: an alternative approach for crop protection. In: Hakeem K editor. *Crop Production and Global Environmental Issues*. Springer, Cham. 2015. DOI: https://doi.org/10.1007/978-3-319-23162-4_11
3. Sundin GW, Wang N. Antibiotic resistance in plant-pathogenic bacteria. *Annu Rev Phytopathol.* 2018;56:161-180. DOI: <https://doi.org/10.1146/annurev-phyto-080417-045946>
4. Fisher MC, Hawkins NJ, Sanglard D, Gurr SJ. Worldwide emergence of resistance to antifungal drugs challenges human health and food security. *Science.* 2018;360(6390):739-742. DOI: <https://doi.org/10.1126/science.aap7999>
5. Guleria S, Tiku AK. Botanicals in pest management: current status and future perspectives. In: Peshin R, Dhawan AK editors. *Integrated pest management: innovation-development process*. Springer, Dordrecht. 2009. DOI: https://doi.org/10.1007/978-1-4020-8992-3_12
6. Thanner S, Drissner D, Walsh F. Antimicrobial resistance in agriculture. *mBio.* 2016;7(2):1-7. DOI: <https://doi.org/10.1128/mBio.02227-15>
7. Copping LG, Duke SO. Natural products that have been used commercially as crop protection agents. *Pest Manag Sci: formerly Pestic Sci.* 2007;63(6):524-554. DOI: <https://doi.org/10.1002/ps.1378>
8. Chandra H, Bishnoi P, Yadav A, Patni B, Mishra AP, Nautiyal AR. Antimicrobial resistance and the alternative resources with special emphasis on plant-based antimicrobials—a review. *Plants (Basel)* 2017;6(2):16. DOI: <https://doi.org/10.3390/plants6020016>
9. Tapia A, Feresin GE, Bustos D, Astudillo L, Theoduloz C, Schmeda-Hirschmann G. Biologically active alkaloids and a free radical scavenger from *Prosopis* species. *J Ethnopharmacol.* 2000;71(1-2):241-246. DOI: [https://doi.org/10.1016/s0378-8741\(00\)00171-9](https://doi.org/10.1016/s0378-8741(00)00171-9)
10. Pasiecznik NM, Harris PJ, Smith SJ. Identifying tropical *Prosopis* species: a field guide. Coventry: HDRA; 2004. URL: https://www.researchgate.net/publication/228714633_Identifying_Tropical_Prosopis_Species_A_Field_Guide
11. Reina L, Bennadji Z, Vinciguerra V, Ferreira F, Moyna G, Menendez P. Isolation and structural characterization of new piperidine alkaloids from *Prosopis affinis*. *Phytochem Lett.* 2015;14(2015):265-269. DOI: <http://dx.doi.org/10.1016/j.phytol.2015.10.022>
12. Pasiecznik NM, Felker P, Harris PJ, Harsh L, Cruz G, Tewari JC, Maldonado LJ. The *Prosopis juliflora*-*Prosopis pallida* complex: a monograph. Coventry: HDRA; 2001. URL: https://www.researchgate.net/publication/236982754_The_Prosopis_Juliflora-Prosopis_Pallida_Complex_A_Monograph
13. de Brito Damasceno GA, Ferrari M, Giordani RB. *Prosopis juliflora* (SW) DC, an invasive species at the Brazilian Caatinga: phytochemical, pharmacological, toxicological and

technological overview. *Phytochem Rev.* 2016;16(2):309-331. DOI: <http://dx.doi.org/10.1007/s11101-016-9476-y>

14. de Brito Damasceno GA, Souto AL, da Silva IB, Roque ADA, Ferrari M, Giordani RB. *Prosopis juliflora*: Phytochemical, Toxicological, and Allelochemicals. In: Merillon JM, Ramawat KG, editors. *Co-Evolution of Secondary Metabolites*. Springer Cham. 2018. DOI: https://doi.org/10.1007/978-3-319-76887-8_15-1

15. Persia A, Rinaldini E, Hapon MB, Gamarra Luques CD. Overview of genus *Prosopis* toxicity reports and its beneficial biomedical properties. *J Clin Toxicol.* 2016;6(5):1-7. DOI: <http://dx.doi.org/10.4172/2161-0495.1000326>

16. Patnaik P, Abbasi T, Abbasi SA. *Prosopis (Prosopis juliflora)*: blessing and bane. *Trop Ecol.* 2017;58(3):455-483. URL: https://www.researchgate.net/publication/322682224_Prosopis_Prosopis_juliflora_Blessing_and_bane

17. Preeti K, Avatar SR, Mala A. Pharmacology and therapeutic application of *Prosopis juliflora*: A review. *J Plant Sci.* 2015;3(4):234-240. URL: <https://article.sciencepublishinggroup.com/html/10.11648/j.jps.20150304.20.html>

18. Henciya S, Seturaman P, James AR, Tsai YH, Nikam R, Wu YC et al. Biopharmaceutical potentials of *Prosopis* spp. (Mimosaceae, Leguminosa). *J Food Drug Anal.* 2017;25(1):187-196. DOI: <https://doi.org/10.1016/j.jfda.2016.11.001>

19. da Silva VDA, da Silva AMM, e Silva JHC, Costa SL. Neurotoxicity of *Prosopis juliflora*: From natural poisoning to mechanism of action of Its piperidine alkaloids. *Neurotox Res* 2018;34(4):878-888. DOI: <https://doi.org/10.1007/s12640-017-9862-2>

20. Sharifi-Rad J, Kobarfard F, Ata A, Ayatollahi SA, Khosravi-Dehaghi N, Jugran AK et al. *Prosopis* plant chemical composition and pharmacological attributes: Targeting clinical studies from preclinical evidence. *Biomolecules* 2019;9(12):2-46. DOI: <https://doi.org/10.3390/biom9120777>

21. Samoylenko V, Dunbar DC, Jacob MR, Joshi VC, Ashfaq MK, Muhammad I. Two new alkylated piperidine alkaloids from western honey mesquite: *Prosopis glandulosa* Torr. var. *torreyana*. *Nat Prod Commun.* 2008;3(1):35-40. DOI: <https://doi.org/10.1055/s-2008-1075314>

22. Samoylenko V, Ashfaq MK, Jacob MR, Tekwani BL, Khan SI, Manly SP et al. Indolizidine, antiinfective and antiparasitic compounds from *Prosopis glandulosa* var. *glandulosa*. *J Nat Prod.* 2009;72(1):92-98. DOI: <https://doi.org/10.1021/np800653z>

23. Gomez AA, Mercado MI, Belizán MM, Ponessa G, Vattuone MA, Sampietro DA. In situ histochemical localization of alkaloids in leaves and pods of *Prosopis ruscifolia*. *Flora.* 2019^a. 256(2019a):1-6. DOI: <https://doi.org/10.1016/j.flora.2019.04.008>

24. Kolapo AL, Okunade MB, Adejumobi JA, Ogundiy MO. Phytochemical composition and antimicrobial activity of *Prosopis africana* against some selected oral pathogens. *World J Agric Sci.* 2009;5(1):90-93. URL: <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.415.4649&rep=rep1&type=pdf>

25. Palacios RA. Los mezquites mexicanos: biodiversidad y distribución geográfica. Bol Soc Argent Bot. 2006;41(1-2):99-121. URL: <http://hdl.handle.net/11336/30488>
26. Almaraz-Abarca N, da Graça Campos M, Avila-Reyes JA, Naranjo-Jimenez N, Corral JH, Gonzalez-Valdez LS. Antioxidant activity of polyphenolic extract of monofloral honeybee-collected pollen from mesquite (*Prosopis juliflora*, Leguminosae). J Food Compost Anal. 2007;20(2):119-124. DOI: <https://doi.org/10.1016/j.jfca.2006.08.001>
27. Singh A. Revegetation of coal-mine spoils using *Prosopis juliflora* in Singrauli coalfield is a harmful practice from an ecological viewpoint. Curr Sci. 2007;93(9):1204. URL: https://www.researchgate.net/publication/279178371_Revegetation_of_coal-mine_spoils_using_Prosopis_juliflora_in_Singrauli_coalfield_is_a_harmful_practice_from_an_ecological_viewpoint
28. Sirmah P, Muisu F, Mburu F, Dumarçay S, Gérardin P. Evaluation of *Prosopis juliflora* properties as an alternative to wood shortage in Kenya. Wood Resour Manag 2008;298(4):25-35. URL: http://bft.cirad.fr/cd/BFT_298_25-36.pdf
29. Lima HG, Gomes DC, Santos NS, Dias ER, Botura MB, Batatinha MJM et al. *Prosopis juliflora* Pods alkaloid-rich fraction: In vitro anthelmintic activity on goat gastrointestinal parasites and Its cytotoxicity on vero cells. Pharmacogn Mag. 2017;13(51): S684-S687. DOI: https://doi.org/10.4103/pm.pm_3_17
30. Valli S, Gokulshankar S, Mohanty BK, Ranjith MS, Ashutosh SR, Remya V. Anticryptococcal activity of alkaloid rich fraction of leaves of *Prosopis juliflora*-A future promising supplementary therapy for cryptococcosis and cryptococcal meningitis? Int J Pharm Pharm Sci. 2014;6(2):491-495. URL: https://www.researchgate.net/publication/288973539_Anticryptococcal_activity_of_alkaloid_rich_fraction_of_leaves_of_Prosopis_juliflora_-_A_future_promising_supplementary_therapy_for_cryptococcosis_and_cryptococcal_meningitis
31. Badri AM, Garbi MI, Gmaraldeem SM, Magzoub AA, Ibrahim IT, Saleh MS et al. Antioxidant activity and phytochemical screening of *Prosopis juliflora* leaves extract. Adv Med Plant Res. 2017;5(3):37-40. URL: http://www.netjournals.org/z_AMPR_17_020.html
32. Lakshmibai R, Amirtham D. Phytochemical analysis and antioxidant activity of *Prosopis juliflora* thorn extract. Malaya J Biosci. 2018;5(1):21-27. DOI: <http://dx.doi.org/10.13140/RG.2.2.35189.88806>
33. Lakshmibai R, Amirtham D, Radhika S. Preliminary phytochemical analysis and antioxidant activities of *Prosopis juliflora* and *Mimosa pudica* leaves. Int J Sci Eng Technol Res. 2015;4(30):5766-5770. URL: <http://ijsetr.com/uploads/514263IJSETR6174-1052.pdf>
34. Batista R, Santana CC, Azevedo-Santos AV, Suarez-Fontes AM, Ferraz JLDAA, Silva LAM et al. In vivo antimalarial extracts and constituents of *Prosopis juliflora* (Fabaceae). J Funct Food. 2018;44: 74-78. DOI: <https://doi.org/10.1016/j.jff.2018.02.032>
35. Singh S, Swapnil, Verma SK. Antibacterial properties of alkaloid rich fractions obtained from various parts of *Prosopis juliflora*. I J Pharma Sci Res. 2011;2(3):114-120. URL: <http://www.ijpsr.info/docs/IJPSR11-02-03-07.pdf>

36. Saadoun I, Challah LM, Alduhouri F. Antagonistic effect of the exotic plant *Prosopis juliflora* extract on different bacterial pathogens. Int J Curr Microbiol Appl Sci. 2014;3(7):865-873. URL: <https://www.ijcmas.com/vol-3-7/Ismail%20Saadoun,%20et%20al.pdf>
37. Alkaabi DS, Gasmelbari ME, Abumukhaimar NAH. Antimicrobial activity of United Arab Emirates indigenous medicinal plants *Prosopis cineraria*, *Prosopis juliflora* and *Acacia tortilis*. Hamdan Med J. 2020;13(2):110-114. DOI: https://doi.org/10.4103/HMJ.HMJ_82_19
38. Bazie S, Ayalew A, Woldetsadik K. Antifungal activity of some plant extracts against (*Colletotrichum musae*) the cause of postharvest banana anthracnose. J Plant Pathol Microbiol. 2014;5(2):1-4. DOI: <http://dx.doi.org/10.4172/2157-7471.1000226>
39. Nascimento MDP, Oliveira CRFD, Matos CHC, Badji CA. Effect of aqueous extract of *Prosopis juliflora* on the control of the mite *Tetranychus bastosi* in physic nut. Rev Caatinga. 2018;31(4):1054-1060. DOI: <https://doi.org/10.1590/1983-21252018v31n429rc>
40. Shah RH, Baloch MS, Khan AA, Ijaz M, Zubair M. Bioherbicide assessment of aqueous extracts of mesquite (*Prosopis juliflora*) on weeds control and growth, yield and quality of wheat. Planta Daninha. 2018;36:1-13. DOI: <https://doi.org/10.1590/S0100-83582018360100014>
41. Ibrahim M, Nadir M, Ali A, Ahmad VU, Rasheed M. Phytochemical analyses of *Prosopis juliflora* Swartz DC. Pak J Bot. 2013;45(6):2101-2104. URL: [https://www.pakbs.org/pjbot/PDFs/45\(6\)/38.pdf](https://www.pakbs.org/pjbot/PDFs/45(6)/38.pdf)
42. Silva AMM, Silva AR, Pinheiro AM, Freitas SRVB, Silva VDA, Souza CS et al. Alkaloids from *Prosopis juliflora* leaves induce glial activation, cytotoxicity and stimulate NO production. Toxicon. 2006;49:601-614. DOI: <https://doi.org/10.1016/j.toxicon.2006.07.037>
43. Yadav N, Rana AC. Pharmacological and pharmacognostical aspect of *Prosopis juliflora*: A review. World J Adv Res Rev. 2020;8(1):36-52. DOI: <https://doi.org/10.30574/wjarr.2020.8.1.0219>
44. Atawodi SE, Ogunbusola F. Evaluation of anti-trypanosomal properties of four extracts of leaves, stem and root barks of *Prosopis africana* in laboratory animals. Biokemistri. 2009;21(2):101-108. DOI: <https://doi.org/10.4314/biokem.v21i2.56478>
45. Abdou L, Karim S, Habou R, Mahamane A. Vegetative propagation trial of *Prosopis africana* (Guill. et Perr.) Taub. by air layering under Sudano-sahelian climate in the South-central Niger. J Bot. 2015;2015: 1-6. DOI: <https://doi.org/10.1155/2015/286582>
46. Houèthégnon T, Gbèmavo DSJC, Ouinsavi CAIN, Sokpon N. Structural Characterization of *Prosopis africana* populations (Guill., Perrott., and Rich.) Taub in Benin. Int J For Res 2015;2015: 1-9. DOI: <https://doi.org/10.1155/2015/101373>
47. Ezike AC, Akah PA, Okoli CO, Udegbumam S, Okwume N, Okeke C et al. Medicinal plants used in wound care: A study of *Prosopis africana* (Fabaceae) stem bark. Indian J Pharm Sci. 2010;72(3):334-339. DOI: <https://dx.doi.org/10.4103%2F0250-474X.70479>
48. Zarafi AB, Ayuba IS. Fungicidal activity of *Prosopis africana* and *Anogeissus leiocarpus* aqueous extracts against leaf blight of *Jatropha curcas* L. Arch Phytopathol Plant Protect. 2012;45(4):413-422. DOI: <https://doi.org/10.1080/03235408.2011.587972>

49. Elmezughi J, Shittu H, Clements C, Edrada-Ebel RA, Seidel V, Gray A. Bioactive natural compounds from *Prosopis africana* and *Abies nobili*. J Appl Pharm Sci. 2013;3(3):40-43. DOI: <https://doi.org/10.7324/JAPS.2013.30308>
50. López-Anchondo AN, López-de la Cruz D, Gutiérrez-Reyes E, Castañeda-Ramírez JC, De la Fuente-Salcido NM. Antifungal activity in vitro and in vivo of mesquite extract (*Prosopis glandulosa*) against phytopathogenic fungi. Indian J Microbiol. 2020;61(1):85-90. DOI: <https://doi.org/10.1007/s12088-020-00906-2>
51. Abdelmoteleb A, Valdez-Salas B, Ceceña-Duran C, Tzintzun-Camacho O, Gutiérrez-Miceli F, Grimaldo-Juarez O et al. Silver nanoparticles from *Prosopis glandulosa* and their potential application as biocontrol of *Acinetobacter calcoaceticus* and *Bacillus cereus*. Chemical Speciation & Bioavailability 2017;29(1):1-5. DOI: <https://doi.org/10.1080/09542299.2016.1252693>
52. Abdelmoteleb A, Gonzalez-Mendoza D, Valdez-Salas B, Grimaldo-Juarez O, Ceceña-Duran C. Inhibition of *Fusarium solani* in transgenic insect-resistant cotton plants treated with silver nanoparticles from *Prosopis glandulosa* and *Pluchea sericea*. Egypt J Biol Pest Control. 2018;28(4):1-5. DOI: <https://doi.org/10.1186/s41938-017-0005-0>
53. Salvat A, Antonacci L, Fortunato RH, Suárez EY, Godoy HM. Antimicrobial activity in methanolic extracts of several plant species from northern Argentina. Phytomedicine. 2004;11(2-3):230-234. DOI: <https://doi.org/10.1078/0944-7113-00327>
54. Gomez A, Sampietro DA, Mandova T, Grougnet R, Kritsanida M, Vattuone M. Compuestos antifúngicos en *Prosopis ruscifolia*: identificación y análisis de su utilidad en el control de especies toxigénicas de *Aspergillus*. Dominguezia 2018;34(2):37-45. Spanish. URL: <http://www.dominguezia.org/volumen/articulos/34205.pdf>
55. Gomez AA, Terán Baptista ZP, Mandova T, Barouti A, Kritsanida M, Grougnet R et al. Antifungal and antimycotoxigenic metabolites from native plants of northwest Argentina: isolation, identification and potential for control of *Aspergillus* species. Nat Prod Res. 2019b;34(22): 3299-3302. DOI: <https://doi.org/10.1080/14786419.2018.1560286>
56. Sirmah P, Dumarcay S, Masson E, Gerardin P. Unusual amount of (-)-mesquitol from the heartwood of *Prosopis juliflora*. Nat Prod Res. 2009;23(2):183-189. DOI: <http://dx.doi.org/10.1080/14786410801940968>
57. Sirmah P, Mburu F, Iyach K, Dumarcay S, Gerardin P. Potential antioxidant compounds from different parts of *Prosopis juliflora*. Journal of Tropical Forest Science. 2011;23(2):187-195. URL: <https://www.jstor.org/stable/23616919>
58. Gianinetto IB, Cabrera JL, Juliani HR. Isolation of N-methylcassine from species of the genus *Prosopis*. J Nat Prod. 1980;43(5):632-633. URL: <https://pubs.acs.org/doi/abs/10.1021/np50011a017>
59. Singh S, Verma SK. Study of the distribution profile of piperidine alkaloids in various parts of *Prosopis juliflora* by the application of Direct Analysis in Real Time Mass Spectrometry (DART-MS). Nat Prod Bioprospect. 2012;2(5):206-209. DOI: <https://doi.org/10.1007/s13659-012-0069-1>
60. Sathiya M, Muthuchelian K. Investigation of Phytochemical profile and antibacterial potential of ethanolic leaf extract of *Prosopis juliflora* DC. Ethnobotanical Leaflets

2008;12:1240-1245.

URL:

<https://opensiuc.lib.siu.edu/cgi/viewcontent.cgi?article=1202&context=ebi>

61. Osano KO, Nyamai DW, Ogola PE, Ouko RO, Arika WM. Evaluation of in vivo toxicity of dichloromethane: methanolic leaf extracts of *Prosopis juliflora* in female Wistar albino rats. J Drug Metab Toxicol. 2016;7(1):1-11. DOI: <http://dx.doi.org/10.4172/2157-7609.1000200>

62. Bezuneh TT, Derressa HD, Duraisam R, Tura AM. Preliminary evaluation of anti-termite activity of *Prosopis juliflora* leaf extract against *Macrotermes* spp. (Isoptera: Termitidae). Cogent Environ Sci. 2019;5(1564170):1-7. DOI: <https://doi.org/10.1080/23311843.2018.1564170>

63. Luis AS, Karin JS, Schmeda-Hirschmann G, Griffith GA, Holt DJ, Jenkins PR. DNA binding alkaloids from *Prosopis alba*. Planta Med. 1999;65(1999):161-162. DOI: <https://doi.org/10.1055/s-2006-960454>

64. Matta D, Nanda H, Mahalingam G. Phytopharmaceutical potentials of *Prosopis lavigata*: A review. Asian J Pharm Clin Res. 2017;10(10):63-8. DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10i10.20316>

65. Ahmad VU, Basha A, Haque W. New alkaloids from *Prosopis juliflora* DC. Z Naturforsch B. 1978;33(3):347-348. DOI: <https://doi.org/10.1515/znb-1978-0322>

66. Ahmad VU, Sultana A, Qazi S. Alkaloids from the leaves of *Prosopis juliflora*. J Nat Prod. 1989;52(3):497-501. DOI: <https://doi.org/10.1021/np50063a005>

67. Dätwyler P, Ott-Longoni R, Schöpp E, Hesse M. Über Juliprosin, ein weiteres Alkaloid aus *Prosopis juliflora* a. dc. 180. Mitteilung über organische Naturstoffe. Helvetica Chimica Acta 1981;64(6):1959-1963. Deutch. DOI: <https://doi.org/10.1002/hlca.19810640629>

68. Nakano H, Nakajima E, Hiradate S, Fujii Y, Yamada K, Shigemori H et al. Growth inhibitory alkaloids from mesquite (*Prosopis juliflora* (Sw.) DC.) leaves. Phytochemistry 2004;65(5):587-591. DOI: <https://doi.org/10.1016/j.phytochem.2004.01.006>

69. Raghavendra MP, Satish S, Raveesha KA. Alkaloid extracts of *Prosopis juliflora* (Sw.) DC. (Mimosaceae) against *Alternaria alternata*. Journal of Biopesticides 2009;2(1):56-59. URL: <http://www.jbiopest.com/users/LW8/efiles/56-59.pdf>

70. Singh S, Verma SK. Synergistic effects of the alkaloids of *Prosopis juliflora*, causing multiple organ toxicity in mouse model. J Biol Act Prod Nat. 2016;6(4):323-336. DOI: <http://dx.doi.org/10.1080/22311866.2016.1240629>

71. Khuong-Huu Q, Ratle G, Monseur X, Goutarel R. Alcaloïdes Pipéridiniques II. Structures de la Prosopine et de la Prosopinine, Alcaloïdes du *Prosopis africana* (Guill. et Perr.) Taub.[1]. Bulletin des Societes Chimiques Belges 1972;81(1):425-441. French. DOI: <https://doi.org/10.1002/bscb.19720810141>

72. Cholich L, Pistán ME, Torres AM, Ortega HH, Gardner D, Bustillo S. Characterization and cytotoxic activity on glial cells of alkaloid-enriched extracts from pods of the plants *Prosopis flexuosa* and *Prosopis nigra* (Fabaceae). Rev Biol Trop. 2021;69(1):197-206. DOI: <http://dx.doi.org/10.15517/rbt.v69i1.43515>

73. dos Santos ET, Pereira MLA, Da Silva CFP, Souza-Neta LC, Geris R, Martins D et al. Antibacterial activity of the alkaloid-enriched extract from *Prosopis juliflora* pods and its influence on in vitro ruminal digestion. *Int J Mol Sci.* 2013;14(4):8496-8516. DOI: <https://dx.doi.org/10.3390%2Fijms14048496>
74. Jahromi MAF, Etemadfar H, Zebajad Z. Antimicrobial and antioxidant characteristics of volatile components and ethanolic fruit extract of *Prosopis farcta* (Bank & Soland.). *Trends Pharm Sci.* 2018;4(3):177-186. URL: https://tips.sums.ac.ir/article_42264.html
75. Yoon MY, Cha B, Kim JC. Recent trends in studies on botanical fungicides in agriculture. *Plant Pathol J.* 2013;29(1):1-9. DOI: <https://doi.org/10.5423/PPJ.RW.05.2012.0072>
76. Kishore GK, Pande S. Integrated management of late leaf spot and rust diseases of groundnut (*Arachis hypogaea* L.) with *Prosopis juliflora* leaf extract and chlorothalonil. *Int J Pest Manag.* 2005;51(4):325-332. DOI: <https://doi.org/10.1080/09670870500404567>
77. Deepa N, Nayaka SC, Girish K, Raghavendra MP. Synergistic effect of *Prosopis juliflora* extract and chemical fungicides against seed borne toxigenic fungi. *Int J Adv Life Sci.* 2013;6(4):312-317. URL: <https://www.cabdirect.org/cabdirect/abstract/20143030211>
78. Ikram N, Dawar S. Application of *Prosopis juliflora* (Sw.) DC. extracts in the management of root infecting fungi of cowpea and mungbean. *Int J Biol Biotechnol.* 2014;11(4):581-587. URL: <http://www.ijbbku.com>
79. Ehteshamul-Haque S, Sultana V, Ara J, Perveen S, Ghaffar A. Use of *Prosopis* spp., in the control of root infecting fungi on okra. *Pak J Bot.* 1996;28(1):109-113. URL: <https://eurekamag.com/research/003/005/003005596.php>
80. Dean R, Van Kan JA, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD et al. The Top 10 fungal pathogens in molecular plant pathology. *Mol Plant Pathol* 13(4):414-430. DOI: <https://doi.org/10.1111/j.1364-3703.2011.00783.x>
81. Giblin FR, Tan YP, Mitchell R, Coates LM, Irwin JAG, Shivas RG. *Colletotrichum* species associated with pre-and post-harvest diseases of avocado and mango in eastern Australia. *Australas Plant Pathol* 2018;47(3):269-276. <https://doi.org/10.1007/s13313-018-0553-0>
82. Griffie PJ. Resistance to benomyl and related fungicides in *Colletotrichum musae*. *Transactions of the British Mycological Society* 1973;60(3):433-439. [https://doi.org/10.1016/S0007-1536\(73\)80028-2](https://doi.org/10.1016/S0007-1536(73)80028-2)
83. Spalding DH. Resistance of mango pathogens to fungicides used to control postharvest diseases. *Plant Disease* 1982;66(12):1185-1186. DOI: <https://doi.org/10.1094/PD-66-1185>
84. Gutiérrez-Alonso JG, Gutiérrez-Alonso O, Nieto-Angel D, Teliz-Ortiz D, Zavaleta-Mejia E, Delgadillo-Sanchez F et al. Evaluación de Resistencia a Imazalil, Prochloraz y Azoxystrobin en Aislamientos de *Colletotrichum gloeosporioides* (Penz.) Penz. y Sacc. y Control de la Antracnosis del Mango. *Rev Mex Fitopatol.* 2003;21(3):379-383. Spanish. URL: <https://www.redalyc.org/pdf/612/61221321.pdf>
85. Deressa T, Lemessa F, Wakjira M. Antifungal activity of some invasive alien plant leaf extracts against mango (*Mangifera indica*) anthracnose caused by *Colletotrichum gloeosporioides*. *Int J Pest Manag.* 2015;61(2):99-105. DOI: <https://doi.org/10.1080/09670874.2015.1016135>

86. Ashfaq MK, Abdel-Bakky MS, Tahir Maqbool M, Samoylenko V, Abdur Rahman A, Muhammad I. Efficacy of prosopilosidine from *Prosopis glandulosa* var. *glandulosa* against *Cryptococcus neoformans* infection in a murine model. *Molecules* 2018;23(7):1674. DOI: <https://doi.org/10.3390/molecules23071674>
87. Miri A, Khatami M, Ebrahimi O, Sarani M. Cytotoxic and antifungal studies of biosynthesized zinc oxide nanoparticles using extract of *Prosopis farcta* fruit. *Green Chem Lett Rev.* 2020;13(1):27-33. DOI: <https://doi.org/10.1080/17518253.2020.1717005>
88. Thakur R, Singh R, Saxena P, Mani A. Evaluation of antibacterial activity of *Prosopis juliflora* (SW.) DC. leaves. *Afr J Tradit Complement Altern Med.* 2014;11(3):182-188. DOI: <http://dx.doi.org/10.4314/ajtcam.v11i3.26>
89. Odhiambo RS, Patrick KG, Helen KL, Gathu NC, Kimani NF, Waithaka WR et al. Antibacterial activity of ethanolic extracts of *Prosopis juliflora* against gram negative bacteria. *Eur J Exp Biol.* 2015;5(11):43-46. URL: https://www.researchgate.net/publication/317527711_Antibacterial_activity_of_ethanolic_ext_racts_of_Prosopis_juliflora_against_gram_negative_bacteria
90. Satish S, Raveesha KA, Janardhana GR. Antibacterial activity of plant extracts on phytopathogenic *Xanthomonas campestris* pathovars. *Letters of Applied Microbiology* 1999;28(2): 145-147. DOI: <https://doi.org/10.1046/j.1365-2672.1999.00479.x>
91. Ahmed NE, Osman NA, Elhussein SA, Mohamed W, Adam YS. Antibacterial activity of *Prosopis juliflora* against tomato bacterial leaf spot disease agent *Xanthomonas campestris* pv *vesicatoria*. Saygili H et al. editors *Acta Horticulturae* 808. ISHS 2009. Proceedings of the II International Symposium on Tomato Diseases. Kusadasi Turkey. 2009. DOI: <https://doi.org/10.17660/ActaHortic.2009.808.53>
92. Sheick MS, Malik AR, Meghavanshi MK, Mahmood I. Studies on some plant extracts for their antimicrobial potential against certain pathogenic microorganisms. *Am J of Plant Sci* 2012;3(2):209-213. DOI: <https://doi.org/10.4236/ajps.2012.32025>
93. Vasquez BL. Resistant to most insecticides. Walker TJ editors. *Book of Insects Records* University of Florida. Florida, Gainesville. 1995. Available: https://entnemdept.ufl.edu/walker/ufbir/chapters/chapter_15.shtml#:~:text=Species%20that%20have%20developed%20resistance,Forgash%201984%2C%20Georghiou%201986
94. Rajappan K, Ushamalini C, Subramanian N, Narasimhan V, Kareem AA. Effect of botanicals on the population dynamics of *Nephotettix virescens*, rice tungro disease incidence and yield of rice. *Phytoparasitica* 2000;28(2):109-113. DOI: <https://doi.org/10.1007/BF02981739>
95. Rawat S, Srivastava M. Evaluation of efficacy of formulations of plant *Prosopis juliflora* against *Callosobruchus chinensis* Linn. *Glob J Sci Front Res.* 2012;12(4):1-4. URL: https://globaljournals.org/GJSFR_Volume12/4.Evaluation%20of%20Efficacy%20of%20Formulations.pdf
96. Senthilkumar N, Gurusubramanian G, Murugan K. Joint Insecticidal Action of *Bacillus thuringiensis* kurstaki with *Annona squamosa* L. and *Prosopis juliflora* Swartz DC on *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae). *Proc Zool Soc.* 2014;67(1):8-17. DOI: <https://doi.org/10.1007/s12595-013-0059-2>

97. Costa RMC, Freire A. Efeito alelopático de extratos aquosos de *Prosopis juliflora* (SW.) DC na emergência e no crescimento inicial de plântulas de *Mimosa tenuiflora* (WILLD.) Nativa 2018;6(2):139-146. Portuguese. DOI: <https://doi.org/10.31413/nativa.v6i2.4768>

99. Al-Humaid AI, Warrag MOA. Allelopathic effects of mesquite (*Prosopis juliflora*) foliage on seed germination and seedling growth of bermudagrass (*Cynodon dactylon*). J Arid Environ. 1998;38(2):237-243. DOI: <https://doi.org/10.1006/jare.1997.0312>

UNDER PEER REVIEW