

Catastrophism and uniformitarianism in Decision Making of Meghalayan Age in East India.

Abstract

Penultimate Epoch is the Holocene had three-way broad subdivisions (late 11.7KYBP (Year before Present), middle 8.325KYBP, and early 4.6KYBP) based on numerous imprints of geomorphologic, meteorological, ice-core, speleothem, and vegetation proxies in nature. Micro-scale ratification of the Holocene epoch, was Greenlandian (11.7KYBP-8.2KYBP), Northgrippian (8.2 to 4.2KYBP), and Meghalayan stage from (4.2KYBP to 50YBP), considering year 2000AD as threshold (b2k) based on DYE-3, GRIP, NGRIP1, NGRIP2 and GICC05, the Greenland Ice Core Chronology 2005 records. The Meghalayan age started with 200years mega drought that shattered primeval agriculture based civilization globally. The stalagmite from various rocks, archeological findings, pollens, δO_{18} isotopic studies and anomalies in Indian summer Monsoon imprints emphasize collapse of well-set Northgrippian Human civilizations in Mesopotamia, Indus and Ganga river plain (Indian subcontinent), the Akkadian Empire (NE Syria). The stratigraphic records offer evidences of climate changes, RSLR, geomorphologic, vegetative cover, Hydrologic processes. The stalagmite proxies, shift of ISM, erratic ITCZ annual drive, left proxies in Stalagmites, in the Mawmluh cave could end to the Holocene epoch classification. The Meghalayan age left exhibits of the upper Holocene that sprinted with severe prolonged droughts due to unstable erratic climate, multiple apocalyptic hazards, and extreme events due to sun-earth geometry, like shifts in Indian summer monsoon, Intertropical Convergence Zone (ITCZ), westerly disturbances, and El Nino Southerly Oscillation (ENSO) activities along eastern part of Himalayas. The ratification and retrofitting of the geological time scale during Holocene well structured and biostratigraphy of eastern India tailor-made.

Keywords: Climate Changes, Holocene, Indian Summer Monsoon, Meghalayan, stalagmites, Eastern India,

1.0 Introduction:

The earth is about 4.54billion years and went through various transformations synchronous with geo-chronometric age as established from radiometric dating of fragments from the Canyon

Diablo iron meteorite. In 2018, newest accepted geologic time scale as per the International Chronostratigraphic Chart International Union of Geological Sciences (IUGS) the Holocene epoch has been subdivided as Greenlandian (11.7KYBP before the year 2000 (b2k)), Northgrippian (8.3 KYBP b2k), Meghalayan (4.2KYBP before b2k), and later succeeded by the Anthropocene epoch, yet not included in Chronostratigraphic Chart 2018. In GSSP run, the 1st marker is bio-stratigraphy whereas the secondary marker considered as the physico-chemical for future GSSP, (Lucas 2018^[1]; Davydov 2020^[2]; Head 2021^[3]). The stratigraphic timescale used for the Greenland ice cores Northgrippian (NGRIP) (I), at 75.10°N; 42.32°W) and NGRIP II (75.10°N; 42.32°W), GRIP, and DYE-3 (Dye 3 in 1971 to 372 m) until 2005. Currently the Centre for Ice and Climate adopted a new timescale in 2005 as the Greenland Ice Core Chronology (GICC05). The NGRIP, and GRIP ice cores, cover the period 7.9–14.8 KYBP (the Bølling, Allerød, the Younger Dryas, and the early Holocene), (Rasmussen et al. 2007^[4]; Walker et al., 2018^[5]). The Anthropocene has coined with substantial consumption carbon energy that have thwarted the natural earth system, (Mishra 2017^[6]; Martin et al., 2021^[7]; Bojie et al., 2022^[8]).

Catastrophism gives a picture of the changes in the geo-hydro-bio sphere on the earth crust with time. The uniformitarianism depicts the continuous and regular changes that occur to the earth crust. Present study is the unveiling of the catastrophic sequences in a uniform manner to explain the dark period from 4200YBP to 50YBP under base period 2000.

Table1: The revised classification of the Holocene by GSSP (ICS)

| EON/ERA /Period | Epoch | Age | Years | As per markers | Physiognomies |
|--|--|----------------------------------|--|---|---|
| Phanero-zoic /Cenozoic /Quaternary (b2k : Before 2000AD) | Anthop-ocene | Yet to be named | 1950-till date yet to be stamped | Demography and human overrule Geo-bio-hydro-environment | Maximum energy use, Human stress on earth; drying of Aral Sea |
| | Holocene (As per DYE-3, NGRIP, or GICC-5 | Meghalayan; Upper/ Late Holocene | 4.2kyrb2k; less shown GICC05 time scale; | Bio-stratigraphy/ physicochemical as 2ndsry marker with Human impact; ($\delta^{18}O$); | Cave Mawmluh; The Lower Eocene Laka-dong dolomite (Calcium rocks) & Therria Sandstone |
| | | Northgrippian; Middle /Mid | 8.2 kyrb2k ; NGRIP 1 ice core. | Bio-stratigraphy as 1 st hand marker climatic cooling | Central Greenland (75.10°N; 2.32°W) It is the ice core at a depth |

| | | | | | |
|--|--|--------------------------------------|-------------------------|---|---|
| | | Holocene | | tailed by rising temp ^{re} . during the Early Holocene | of 1228.67m, near the 8.2 ka event |
| | | Greenlandian; Lower/ early Holocene; | 11.7KYBP b2k; & NGRIP 2 | Change of the evaporation at rain/snow source after Greenland Stadial1/Younger Dryas, together with changes in dust load,& CC | The central Green-land ice sheet; subseries /sub-epoch is located at 1492.45 m in the NGRIP2 ice core |

The Meghalayan started 4.2KYBP, shaped by continuous 200years **mega-drought**/ very high flood phenomenon that shattered well-developed civilization in Asia such as Mesopotamia(for prolonged salinization in Syria), Yangtze Valley, Nubia (reduced river flow), Indus valley (Mahenzodaro and Harrapan), Ganges flood plains (Mahajanapadas), Egypt, Greece (Anatolia), Israel, Palestine, and Turkey, cultures across the globe where tropical monsoon went weak. The changes were due to shift of monsoon pattern, ITCZ, Huge Azores High and the western disturbance after a good SW monsoon period in mid-Holocene (Northgrippian) period, (Kidder et al., 2012^[9]; Kathayat et al., 2017^[10]; Jacobson et al., 2021^[11]). Later during mid Meghalayan **period**, many Vedic civilizations on Ganga River Valley (Guje Kingdom, Sahiya), due to shift of ISM and ICTZ.

1.1 Reasons for study:

The world has encountered a series of climatic extreme events and various types of apocalyptic disasters in last two decades from 21st Century. The cataclysm of all types of waves of disasters like biological (Pandemics), Geophysical (Tsunamis); Earthquake (Haiti); Volcanic activities (Hunga Tonga) and many others passed over India. The conception of those are not instantaneous but a continuous change in sun-earth geometry. The geo, bio and hydrological changes are the results of geospatial complexities of climate changes inherited from paleo climatic proxies. From ice core explorations conducted Greenland the first two ages of Holocene epoch (**Greenlandian and Northgrippian**) prefixed from ice core studies such as Greenlandian and Northgrippian. The penultimate compartment and its geological establishment was only possible after exploration of the stalagmites from the Calcareous Rock fabric Mawmluh cave.

Least correlation made between later part of Holocene with climate, geology, limnology and lithology of the area surrounding the stalagmites under study. In addition, the ice core studies of Greenland exploration is silent about the post Holocene from 4300YBP. Present study is an attempt to study the available paleo-climatic proxies to phase the Meghalayan age from 4.2KYBP to 70KYBP, and link with the climatic vicissitudes that has occurred during the Anthropocene epoch.

2.0 Methodology

Meghalayan word hailed from the demarcated rock mosaic analyzed by Mike Walker (ICS, 2018) located in the stalagmite in one of the longest and deepest Cave Mawmluh entrance (25°15'44"N; 91°42'54"E) at 1290m high stalagmites proxies in the Meghalaya state, east India. The stalagmites in Meghalaya studied. The results of various oxygen isotopic investigations inferred differential numbers of neutrons that recognized dry paleo-climate with weak paleo-monsoon in the transit period between 4300 to 4100 Years before present, YBP, Vyawahare 2018^[12]. The rock slice and the mosaics formed links to major geological activities like splitting of continents, swings in climate, and can be associated with appearance of various types of the biome with geological time scale, (Amos J 2018^[13]). Because layers of stalagmites/stalactites are shaped due to deposit of calcium opulent water infiltrating into the cave surface. The tephrochronology study of layers signifies cycles of deposit of calcareous materials (volcanic ashes) dated by isotopic methods. The Meghalayan age accepted officially by International commission of stratigraphy (ICS) on July 13, 2018 (Walker, 2012^[14]).

The Archives which signifies about an geographic stages are fluvial, marine and windblown sediments, lake and peat from lacustrine areas, glacial moraines and proglacial deposits, ice cores, corals, speleothems, tree rings stalagmite and tufa's. The Proxies that also help in ascertaining the geomorphic changes with geological time scale are sediment size/distribution, mineralogy changes, Pollens, Magnetic properties of minerals. The fossil imprints, isotopic composition of organic matter, Oxygen, Nitrogen, carbon and hydrogen, carbon vs Nitrogen, Formation of gypsum under extreme climate etc. add to it., Harris et al 2008^[15]; Singhvi et al., 2008^[16]; Laskar et al, 2021^[17])

Various methods to ascertain geospatial age

$$i. \quad A = \text{Age of a phenomenon} = \frac{A_p - A_f}{A_u} \quad \text{eqn 1}$$

Where A = age to be determined; A_p = present status of activity; A_f = time taken for reaching final status of activity and; A_u = unit rate of activity (Singhvi et al., 2008)

- ii. Geo-chronometry method to find the age by dating method where the depth of the stratigraphic layer, more is the relative age under sub-jurisdiction of degenerative process and the amount of organic residue left.
- iii. Carbon dating method: Carbon has three isotopes i.e. carbon-12 ($^{12}\text{C}_6$), carbon-13 ($^{13}\text{C}_6$), and carbon-14 ($^{14}\text{C}_6$). Carbon-12 and carbon 13 are stable whereas carbon-14 is unstable and undergoes radioactive decay and transforms stable nitrogen ($^{14}\text{N}_7$) which take a long period depending upon half-life period. Radiocarbon dating used to measure ages of things that were geologically recently formed (initial av. half-life 5730 for each half-life and used for age range of application of 70000years). The carbon dating is used against organic materials such as bones, wood, charcoal, shells (Peppe et al., 2013^[18]).

The radiocarbon dating equation of $^{14}\text{C}_6$ to $^{14}\text{N}_7$ is:



$$N_t = N_0 e^{-kt} \quad \text{eqn 3}$$

$$\text{And} \quad k = \frac{\ln 2}{t_{0.5}} \quad \text{eqn 4}$$

Where N_0 = Number of atoms of the isotope in the original sample (when time $t = 0$), N_t = Number of residue atoms on elapse of time (t). k = the rate constant for the radioactive decay

The other physical dating methods used are K-Ar dating for potassium/glass minerals, Luminescence (10 to 10^6 YBP) for Quartz, feldspars, carbonates etc., Fission tracking (1,000 to billion YBP) and many others. The chemical dating methods in geological correlated with existence of similar fossils called “Biostratigraphy” or “Tephrochronology” where the Sediment/ashes strata like Foraminifer, Ammonite occur in identical age. The Annual Layer Counting Methods, i.e. dendrochronology or lake sediments or speleothems analyzed to find the age of the proxies.

Present study includes study of all the available methods and proxies to construct the Holocene epoch that induced its shift to the Anthropocene Epoch perturbed the geo-bio-hydro sphere.

3.0 Ice core exploration Green land:

Haas et al., 1998^[19] identified six compartments in Holocene epoch 9.6 to 9.2KYBP, 8.6 to 8.15, 7.55 to 6.9; 6.6 to 6.2; 5.35 to 4.9; 4.6 to 4.4; 3.5 to 3.2 and 2.6 to 2.35KYBP as radiocarbon years considering the climates of sites at Alps and Swiss Plateau. The 21st century Greenland ice core chronology depicted from the DYE-3, GRIP, and NGRIP (I, and II) ice cores, identified as the Greenland Ice Core Chronology 2005 (GICC05). The ice core tell less about the late ages of the Holocene epoch. The Global Boundary Stratotype Section and Point (GSSP), as per the NGRIP2 Greenland ice core at a depth of 1492.45 m. dated back to 11.7KYBP b2k (before 2000 CE) event (Greenlandian age). The NGRIP1 ice core, at a depth of 1228.67m signifies 8.2KYBP b2k event. The late Holocene setting at the 4.2KBYP b2k event (Meghalayan age) could not be explained by the ice core or by GICC-5 chronology. The speleothem from (Stalagmite KM-24) in Mawmluh Cave, Meghalaya, could be able to help the geologist for the 4.2KYBP event (Rasmussen et. al., 2006^[20]; and Walker 2018^[5])





Fig 1: Greenland Ice core records (DYE-3, GRIP, NGRIP (I, and II) and GICC-5; less presenting late Holocene (Meghalayan age) (Source modified [Rasmussen et. al., 2006^{\[20\]}](#); [Walker 2012^{\[14\]}](#))

3.1 Speleothem Stalactites and Stalagmites Eastern India

Speleothem, stalactites and stalagmites in Caves in India (the CaCO_3 - H_2O) paleo-thermometer) provide important proxies of Holocene paleo-climate and paleo-monsoon, as not subjected to erosion, diagenesis. They are calcite and aragonite materialization in speleothems of different hills and their cave systems and terrestrial **deposits in Meghalaya**. The geological formation has 200km long x 30km wide lime stone band plunged into range of hills of West Garo, West Khasi Hills, eastern parts of Khasi and Jaintia Hills (**Fig -1**). The Meghalayan Adventures Association (MAA) has explored cave passage of length 320km and about a milieu of entrances to these caves at a height of 1000m to 1100m and some uncharted. The bio-speleology search has explored some caves (krems in local name) like Um-Lawan, Mawkhyrdop, and Mawmluh, Wah Ser, Krang Maw, Siju, Pyrda, Umthloo, Kotsali, and Wah Tylliang etc in Meghalaya. Mawmluh is 8th stretch of cave of span 7194m with about $\approx 4000\text{m}$ is undefended for caving by Meghalayan adventure association (MAA). The stalagmite in the Himalayas (Chulerasim), Bhimbetka caves (Bhopal), Mawsmal (Shillong), Kailash, & Kotumsar caves (Basstar, Chhattisgarh), Belum caves (AP at 3229m), Gupteswar and Khandagiri caves (Odisha), Mahakal caves WB, Dungewari caves (Bihar), and Borra Caves etc. are the exhibits and proxies that depicts about Meghalayan age which are yet to be studied.




Fig 2: The Index map and mult-proxy of the Meghalayan age stalagmite, East Khasi Hills, Mawmluh cave, (KM-24), (source modified: [Kalpana et al., 2021^{\[21\]}](#); [Dildi et al 2021^{\[22\]}](#))

The petrogenic study of calcareous stalagmite, at the Mawmluh cave entry, East Khasi Hills on river Lum Lawbah, in 2012 revealed a ratification of geological age of the late Holocene epoch as the Meghalayan stage from 4.2KYBP b2k. Walker and Rasmussen are the geologist analyzed the rock fabric and could solve the after myth of the 4.2kybp events bio-stratigraphically. The before present (BP) **refers to the** year 1950 i.e. start of Anthropocene whereas b2k of the ice core considered before the year 2000. Other targeted stalagmites explored are Mawkhydrop, Mawsmal, Rupnath, Mondilcole etc., [Harrish et al., 2008^{\[15\]}](#).

3.2 Archeological Findings India

Many archeological findings against fall of civilizations, such as Mesopotamia, Akkadian Empire, and BET Dwarka submerged Island; Durgadevi (WB) Sindh civilizations (Harrapa and Mohenjo-Daro) are contemporary to early Meghalayan stage. The archeological excavations, rock records, speleothems, radiocarbon studies, dendrochronology about the Holocene rainfall incidences and climatic variability affecting agriculture, lake/lagoon morphology, ([Weiss et al., 1993^{\[23\]}](#), [Cullen et al., 2000^{\[24\]}](#), [Kotlia et al., 2015^{\[25\]}](#), [Kong et al., 2017^{\[26\]}](#), [Weiss, H., 2017^{\[27\]}](#), [Quamar et al. 2021^{\[28\]}](#)).

4.0 Past reviews

The human settlement with agriculture based life probably started around early Holocene time. The history of habitation in India as hunter-gatherers dates back to 30KYBP as per haplo groups DNA tests. That proved Indians are West Eurasian origin, particularly in Hindu upper cast. The existence of homosapiens found in various excavations and stalagmites even up to 65KYBP ([Singh et al., 1999^{\[29\]}](#), [Bamsad et al., 2001^{\[30\]}](#), [Banerji et al., 2017^{\[31\]}](#), [Clarkson et al., 2020^{\[32\]}](#), [Callaway 2021^{\[33\]}](#)). The climate change (CC), in short or spatial scale in ISM linked with agricultural yield, socio-political economy, prevalence of biological disasters, and geo-bio-hydro sphere in Indian subcontinent. The climate is the cooling engine of the earth's planetary system, [Berkel. H 2012, Chiotis 2018^{\[34\]}](#).

The trade winds, El Niño and the Southern Oscillation (ENSO), irregular oscillation of sea surface temperatures, the asymmetrical oscillation of SST (sea surface temperatures) called positive Indian Ocean dipole (PIOD) etc. are the associated meteorological phenomenon associated with the climate change of India. (Wang et al., 2005^[35], Borzenkova et al., 2015^[36], Mishra S. P. et al 2020^[37]). Late Holocene massive drought/famine is rock record neither a myth nor a Hyper object proved by rock records anointed as 4.2KYBP-old event or Meghalayan emergence by the International Commission on Stratigraphy (ICS) in 2018, after Walkar, 2012^[14]

The multi-proxy data, such as pollen records, Radio carbon dating, paleo-temperature, lake sedimentation, sediment transfer, ecological collections reveals about earth science changes. The Meghalayan stage of late Holocene climate has been compartmentalized with median spacing of 400yeas with control point being 3000YBP, both in Mid latitude and equatorial regions, Sun et al., 2013^[38], Jenny et al., 2019^[39], Kaufman et al., 2020^[40], Hao et al., 2021^[41].

4.1 Paleontology proxies:

The Meghalayan cooling events in the Last Ice Age (LIA) has paleo-climatic and archaeological evidences from India. They indicate that the setting of the nature and societal changes have substantially influenced the abrupt climate change around 2.8ka BP (Park et al., 2019^[42]).




Fig 3: Holocene temp. anomalies during Meghalayan era (source rebuilt: Park et al. 2019^[42])

Pollen records (arboreal pollen percentage data) **reveal** the drying spells during the latter part of the Holocene epoch. The Late Holocene was compartmentalized four-time spans 4.2ka-3.7ka, 3.2 ka-2.8ka, 2.8ka to 2.4ka, and 2.4ka BP to present. The cooling episode was due to solar-insolation shift in atmospheric circulation. The mega droughts between 4.4ka to 4.0KYBP led to the collapse of ancient civilizations in East Asia. (Fairbridge et al., 1976^[43]; Park et al., 2018^[44]; Mishra et al, 2021^[45]), sparse pollen proxy data is available with some legendary and epic evidences of the late Holocene period in eastern India

The Kemp Mawmluh, of Meghalayan age is 7.15km long and rich in mineral deposits and adorned with numerous formations of stalactites and stalagmites, Chambers, rivers, moon milk, fishing pond, crawling part, vertical holes, etc. The stalagmites and the rock fabrics in the Mawmluh caves represents prominently the proxies of growths in stalagmites occurred during various geological time scale and prominently the Meghalayan age (4.2K to 0.07K YBP), (**Fig 3**) Walkar, 2012^[14]





Fig4: Photographs inside caves of Mawmluh; (source: <https://www.tripadvisor.in/Attraction>)

4.2 Multi-proxy flood records:

Evidences of sedimentary sequences in rhythmic cycles recorded in the various rivers like the Ganges, the Brahmaputra and the Mahanadi system in Meghalayan age. The Sagar island (21°37'21'' to 21°52'28''N lat. and 88°2'17'' to 88°10'25''E long.), (Gopinath et al., 2005^[46]) Mahisani island, mud point island at the Hugli estuary of the Ganga system. Similarly, the Mahanadi delta system has shown retrograding shoreline during Meghalayan age with huge sediment influx and flood flow with deltaic progradation and sea level adjustment. Major changes in delta building processes occurred during late Holocene up to 800YBP (Rao Sambhasiva et al 1978^[47], Mahalick 1996^[48], Mohanty et al., 2005^[49], Mishra et al., 2021^[50]). The largest Asian lagoon Chilika was a river estuary during Pleistocene. During Northgrippian age, maximum sea level rise and heavy ISM rainfall converted the large low marshy land to a gulf covering a vast area more than 1500km² (Venkatratnam 1970^[51], Sadakata and Kakani et al., 1993, Sethi and Mishra S P., 2020). The erratic ISM, intermittent very high floods and continuous long spell droughts during early Meghalayan age led to formation of Southern barrier islands and converted the gulf to a lagoon (Khandelwal et al., 2007^[54]). The distributaries in the Mahanadi river system have multiple ISM signatures and multi-proxy events in change of courses in the Meghalayan age. Few evidences are dominance of the Burdha, Old Kathajodi, Prachi and Ratnachira system and later weakening and emerging as Devi, Kuakhai, Kushabhadra, and Daya system and many-abandoned channels (Old Kathajodi, Alaka, Prachi, Sunamunhi and Malini etc) during Meghalayan era (Mishra et al, 2021^[50]).




Fig 5: Change in river courses in the Mahanadi delta; due to Hifloods Meghalayan age

4.3 Holocene Paleoclimate:

Terrestrial and marine climatic proxies recorded those that are sensitive climate changes. Three trend in paleo-climatic signatures established during Holocene. The rise in short-range cold phases called Greenlandian (11.0–8.0 KYBP); stable warm climate named Northgrippian (1.0–3.5 °C above during 8.0–4.5KYBP). The present titled as Meghalayan phase with declining temperature trend, and climatic uncertainties from 4.5KYBP to 71YBP. The proxies reflected from lake levels and vegetation. The Meghalayan stage include abrupt temperature settings like the Medieval Warm Period, the Little Ice Age and the Modern warm period, [Borzenkova et al., 2015^{\[55\]}](#). Geochemistry studies of Chachi and Luni along west coast of India reveal about two stages of time-based variabilities (4.6KYBP to 2.5KYBP) in vegetation, humidity, and environmental vicissitude and from 2.5KYBP to 70YBP, characterized by less rainfall and enough mesic vegetation ([Pillai et al., 2018^{\[56\]}](#)). Climate reconstructions during the Meghalayan age, classified as forcing of anthropogenic aerosol, sun earth geometry, volcanic eruptions, seismic, Green House gases (GHG) gasses and solar irradiance. The reconstruction of late Holocene (Meghalayan sage) in Northern Hemisphere (NH) the anomalies of surface air temperature (SAT), i.e. ice ages and worm periods. Models structured for SAT of paleo climate, reveal that GHG gases rise is responsible for the high SAT temperatures occurred at millennium frequency @0.36 times per 4000YBP for NH temperatures during late Holocene. The global

SAT plummeted by Little Ice Age, later rose continuously between AD 1450 and 1850, (Kobasi et al., 2013^[57], Mishra et al., 2020^[58], Scott 2021^[59]).

4.4 Inter Tropical convergence zone:

The tropical/subtropical landmass due to solar insolation during summer adds energy to the atmosphere. The energy try to push the inter tropical convergence zone (ITCZ) to lower energy zone shift towards north causing advection of moist and warm air masses from SW-ly direction. The motion of the ITCZ occur towards Himalayan range along with Tibetan Plateau block the advection of dry and cold air masses received from mid-latitude oceans as westerly circulation. The phenomenon called The Ventilation effect delimits movement of ITCZ further north. Similarly, during post monsoon, the westerly disturbances and subtropical highs push the ITCZ towards equator and finally push it to southern Hemisphere during February. The movement and positioning of ITCZ, indicate the intensity of ISM, creation of cyclonic disturbances in North Indian Ocean (NIO), including BoB and A.S. (Ramisch et al., 2016^[60]; Quamar et al., 2021^[61]) Fig 2 (b).

4.5 Azores High

The persistent and quasi-stationary atmosphere's circulation in N-Hemisphere, the Azores High or Bermuda High is the subtropical anticyclone in Atlantic Ocean positioned (25°N to 35°N) with deep tropical easterlies moving equatorward and mid-latitude westerlies propagating towards pole, that is the pivotal center in the global climate system. On study of inter annual ISM rainfall it is observed gradual shift from west to east and vice versa between NE Himalayas and western Himalayas being influenced by vigorous Bermuda High and triggered by Ross by wave. (Binkley 1987^[62]; Jadav 2021^[63]) Fig 2(a).

4.6 Indian Summer Monsoon:

The Asian monsoon intensification linked with the Tibetan Plateau uplift consequent upon the impact of collision of the Indian plate with Asian plate ≈ 50 MYBP (Royden et al. 2006^[64]) In India, the ISM (Indian Summer Monsoon) prevails in India during June to Sept of the year that give about (75% to 80%) of the annual rainfall of the area. Annual erraticism of ISM rainfall is sturdily predisposed by EL-Nino Southerly Oscillation (ENSO), Indian Ocean's N-S, gradient of

sea surface temperature, pIOD (positive Indian Ocean Dipole), boreal summer intra seasonal oscillation (BSISO), El Nino Modoki events and Madden-Julian oscillation (MJO). Present study focusses on the changes in ISM for dry period with mega drought events, that had brought two-century long severe drought and shattered many civilizations in Africa (North), the Middle East, and south Asia (**Fig 6**).




Fig 6: Shift of ITCZ from west to East Fig: 2(b) Position of ITCZ over Indian Subcontinent.

Source: <https://scied.ucar.edu/docs/why-monsoons-happen>

During Meghalayan age (before 4200YBP), the ISM rainfall was highest in Indus River valley so the Harappa civilization established and grew, later shift of the ISM to central India during early Meghalayan stage brought Sindh civilizations under rain scarce area causing the city to fall. The shift of monsoon from west to east and abnormal shift of ITCZ had changed the monsoon pattern in Indian sub-continent. The records of the isotopic shift reveals there was decrease in monsoon rainfall, [Berkelhammer et al., 2012^{\[65\]}](#). The uniform growth of logs of C4 plant rings during ≈ 4.7 to 2.0 KYBP signifies steady waning of ISM with maximum at ~ 2.0 KYBP being the weakest monsoon ([Kumar et al., 2019^{\[66\]}](#)). Forcing factors responsible for shift of ISM, climate variables, and unconventional swing ITCZ. The Little Ice Age, (LIA), India had prevalence of wet conditions in the north, west, and central states. The dry macroclimate occurred over majored part of peninsular India ([Banerji et al, 2020^{\[67\]}](#)).

4.7 Changes in RSLR:

There was subsidence due to glacio-isostatic adjustment (GIA) against global warming. The consequent mean Regional Sea Level Rise (RSLR) rates from ~ 6 ka to present were 1.4 mm yr⁻¹

inferred from geophysical model studies (Cronin et al., 2019^[68], Yan et al 2021^[69]). Kumar, et al. (2021^[71]) reconstructed transgression and regression of sea during Meghalayan age. The archaeological findings have established settlements during late Holocene in coastal eastern India (Erenda and Sirjua) but no settlement during Northgrippian, and Greenlandian period. Archeological discoveries of underwater monuments at Mahabalipuram, Vet-Dawrika, Ram Setu at Rameswarm etc




Fig 7: The Holocene RSLR changes with climate proxies (Source mod: Loveson et al 2015^[70])

Strand line movement hypothesized during 5.0KYBP but during 4.0KYBP regressive proceedings recorded in east coasts of India. Later transgressive events recoded between 2970±100YBP and c. 3400 YBP along the eastern coast and northern South China Sea (Kumar et al., 2021^[71]). The coastal transformations were recorded due to major transgressive event, Farooqui et al., 2000^[72] have reported sea-level regression following the major transgressive event at 3150 BCE to 2370 BCE which transformed the geomorphology of the Chilika, Kolleru and even Pulicat Lagoon in AP (Andhra Pradesh) (Jadav et al., 2018, Mishra S. P. 2020^[74]). A drop in regional sea level points during the Little Ice Age (LIA) (300-250YBP) (Fig 7). The archeological findings from various excavations have left their proxies at Gopalpur, Golabai, Delang (Hariraj Pur), Balasore (Durgadevi and Ranasahi) of Odisha indicates a distinct agro-based economy during Neolithic-Chalcolithic age (Mishra et al., 2021^[75]).

4.8 Westerly movement in Indian peninsula

During Holocene period, Ladakh had four prominent phases such as ~10,800–10,000YBP; ~8800–8600YBP; long phase of warm aridity climate at ~5200–2600YBP; ~1700–1500YBP and ~500 YBP) interfered by relatively warm climate. Along with ISM, the westerlies advancement had governed the hydro climate of northwestern parts of Bihar, WB, Jharkhand and Odisha India. The westerlies governed during Northgrippian while ISM dominated during Greenlandian age. During mid-Holocene (~3200YBP), Westerlies predominated the ISM. The lake and other records from the region in the years during Holocene—7200, 5200, and 2600 YBP although the westerlies have poured heavily in the Ladakh region latter,(Phartiyal et al., 2022^[76]).

5.0 Vegetation and Agriculture

5.1 Vegetation History Meghalayan age:

The pollen evidence suggests that between ca. 3000 and 2600 YBP, savannah vegetation occurred in the region having a comparative lesser monsoon rainfall. The forest expanded as an open-mixed tropical deciduous forest between 2600YBP and 2200YBP under a warm and moderately humid climate with increase in ISM rainfall.




Fig 8 : Rise in CO₂ in atmosphere during the Meghalayan Age; (Luthi, D., et al., 2008^[77])
 Later from 2200 - 2000 YBP, the open-mixed tropical deciduous forests in east India and adjoining Himalayas transformed the dense deciduous forests due to swing of the mean path of ISM with rise in monsoon rainfall. Afterwards the gradual warming and the rise of the ISM,

(2600YBP to 1850YBP), during the Roman Warm Period (RWP), with intensified human activities, decreased the land use pattern (**Fig 8**).

There was gradual intensification of the monsoon between ca. 2600 and 1800YBP, and an increase in the ISM until present warm period. **Presently**, dense mixed tropical deciduous forest has existed under a warm and relatively more humid climate indicating a further increase in monsoonal precipitation but reduced under anthropogenic stress (Bonnefille et al., 1999^[78]; Quamar et al., 2017^[61]; Kumar et al., 2019^[66]; Banerji et al., 2020^[67])

5.2 Mangrove vegetation:

Palynological studies **reveal that** salt tolerant Mangroves resistant to saline atmosphere were extending from Kanuru, and Machlipatnam coast until Myanmar coast including Chilika coast during mid Holocene (Khndelwal et al., 2008^[54], Farooqui et al., 2016^[72]). Researches but sporadic about formation of coastal indicators like those that barrier islands, mangrove, sand dune vegetation, mud flats, and Lagoons reported in literature. Mangroves; reducers of CC stresses, RSLR and drought; cover 0.7% of the global tropical forest area but provide lion's share of ecosystem services by sequestration, Lovelock et al., 2017^[79].

Reports available about RSL anomalies during Last Glacial Maximum (26 to 21KYBP), there were MSL rise in ≈ 130 – 120 m lower than present are the global Quaternary eustatic sea-level cycles. The Greenlandian age refers to sea level transgression in BoB coasts. That influenced geospatially the mangrove transgressive segments supplemented by ISM /EASM (East Asian summer monsoon) and variable sediment flow through major inflowing rivers like the Irrawaddy, the Brahmaputra, the Ganges, the Mahanadi systems and the largest Lagoon Chilika associated with various species of mangroves along their estuaries.

Initial phase of Meghalayan age (≈ 4.2 KYBP, and ≈ 2.6 KYBP), considered warm and humid climatic conditions. However, degrading trend in vegetative status reported. After 2.6KYBP, mangroves re-flourished along the BoB coasts until 2.3KYBP, consequent upon relative stability of sea level. The invasion of midland fresh water dry climate taxa, progressions and retrogressions of strandlines along East coast of India and West coast of Myanmar has huge mangrove spread. Present mangrove afforestation has been futile for selection species, RSLR and climatic traumas. The Mangroves plantation tried along Chilika coast last five years but the

growth is rarely luxuriant even if tried by CDA near Arakhakuda ([Pandey et al., 2014^{\[80\]}](#), [Feller et al., 2017^{\[81\]}](#), [Mondal et al., 2021^{\[82\]}](#))

6.0 Geological studies

6.1 Tectonic Subsidence:

The large deltas of the Mississippi, the Rhine, the Rhône, the Danube, the Nile, the Amazon, the Tigris-Euphrates, the Ganges, the Mahanadi, the Godavari and the Indus, were under tectonic subsidence. The major Tectonic Zones (TZ) are at Himalayas (East to west), Assam-Arakan, Baluchistan-Karakoram, Andaman-Nicobar and Unwavering Continental Regions (Zone V). The river basins of India, Myanmar and other countries surrounding Bay of Bengal are sedimentary. The East India shield margin comprised of vigorous tectonic activities in past along the Shillong plateau, Eastern Ghats, Chottanagpur plateau and the Southern Granulite Terrain (SGT). The Indian plate had easterly subduction between the Burma microplates as the core of the tectonic activities... The Indo-Burman ranges remains northerly and form the Naga Hills trending SW-NE. The result was the (IBCZ) Indo-Burman convergence zone associated with continuous tectonic activities and landslides undergoing subduction continuously along Shillong Plateau and Mikir Hills. The Tripura fold with associated thrust resulted the forehead basin along the Bengal Basin, [Biswas et al., 2014^{\[83\]}](#), [Kumar et al., 2015^{\[84\]}](#).

6.2 Transformations in deltas

During Meghalayan age, the deltas of the lower Ganges, the Mahanadi tri-delta, Subarnarekha delta along with the Golabai Sasan and Ramchandrapur within Chilika basin changed their morphologic, and eustatic sea level during the Meghalayan Holocene. Modifications in the coastal geomorphology observed in anastomosed drainage pattern, estuarine configurations, tidal inlets, along with sinking and shrinking of the lakes and lagoon, sedimentations, and marine transgression (change in strandline). Dating back from 4.2kYBP to 2.1kYBP (the Chalcolithic phase, Iron age), there is archeological evidences along deltas of the Mahanadi Odisha and the Chilika coast ([Kingwell et al., 2018^{\[85\]}](#), [Mishra S. P., 2018^{\[86\]}](#), [Hazara et al., 2020^{\[87\]}](#)).

7.0 Socio-political

7.1 End of Indian Civilizations:

In past, major civilizations (Harappa or Mahenzo-daro) built up along Indian subcontinents either in alluviums, flood plains or delta's of Sindh or Ganga rivers during pre and mid Holocene epoch. The collapse of those civilizations presumed due weak ISM, East Asian summer monsoon (EASM) and high solar insolation's during early part of post Holocene based on many climatic proxies and paleo-climatic records,(**Table 2**), [Cheng et al, 2021^{\[88\]}](#).

Table 2: Societal and technological growth of people during late Pleistocene in Indian subcontinent

| Period Name | Climate condition | Human popul ⁿ | Societal/ technology development | Sites as Microliths |
|------------------------------------|--|---------------------------------------|--|---|
| Paleolithic period (2MaYBP-10KYBP) | Extreme cold; Younger Dryas | <0.1mn Homosapiens; Stone age | Fire, Tools of stones, Ostrich Egg; Hunter gatherers; caves life, Saber-toothed tigers and cave lions | Bhimbetka(M.P),Hunsgi, Kurnool Caves, Narmada Valley (Hathnora, M.P), Kaladgi R., Mehargarh |
| Mesolithic Period; (10- 8.2 KYBP) | Major CC: Cold; glaciation | Green landian <1mn people | Agriculture, Cattle fostering); stable life; Mining, melting; casting; Dentistry | Brahmagiri (Mysore), Narmada, Vindya, Gujarat; Indus and Gangesvalley; |
| Neolithic Period (8.3-4.2 KYBP) | Warming started after the Ice age; Plague of Justinian (pandemic) | North-grippian (Megalithic) ≈1million | Primal Agro-based village, Ayurveda knowledge wheels cremation over burial; Flush toilet; Pepper in cooking; Kingdom & rullers; weighing scale | Burzahom, Gufkral (J&K), Mehrgarh Indus valley) ,Chirand (Bihar), Daojali Hading (Tripura), Koldihwa (UP), Mahagara (UP), Hallur & Paiyampalli (AP), Maski, Kodekal, Sangana Kaller, Utnur, Takkala Kota. |
| Chalcolithic Period (4.2-3.6 | Abrupt shift cooler to drier | Meghalayan; Bronze age; ≈6Mn people | Indo-Aryan Exodus Glass making; & Fall of Indus valley | Brahmagiri, Navada Toli (Narmada region), Mahishadal (W.Bengal), |

| | | | | |
|--|---|---|---|---|
| KYBP); Early Vedic period | climate | | culture, Ganges plain evolution; Plastic surgery | Chirand (Ganga region); Golabai & Harirajpur (Odisha) |
| Chalcolithic Period (3.6 -2.6 KYBP);Globally Medieval warm period) Megalith Culture; Jain & Buddh-ism; | ITCZ Shift Indus valley to Ganges Plain | Meghalayan; Iron-age India (Vedic- Upanisad);≈25Mn people | Influx Aryans; Boat making; Medicines; hand propelled wheel, Cemetery H culture ;Extraction copper Arravali Hills; Bronze alloy from Ambaji | Karim-Shahi (Gujurat); Sisupalgarh, Manik Patna Talapada, (Odisha); Atara njikhera (Etah), ,Sunet Punjab; Parsvanath,, Mahajanapads Ancient civilization; like China, Egypt & Greece |
| Classical era; 2.6KYBP- 800YBP (vedic age) Excellency in Iron works | Gupta Kingdom (300AD – 800AD): Classical Period | Meghalayan (Early medieval) | Science developed (Math, Astronomy & medicine); Kingdoms /Nations; wootzsteel (south India); Iron pillar Delhi. | Mauryan Period, Chola, Chera, Pandyas, Sakas, Kushanas; Gupta kingdoms plague, smallpox / measles (≈36mn lives) |
| Pre Christian era 1200YBP- 0300YBP | Moghul Period | Late Medieval India | Pyrotechnic works; Rise Islam/ Sufism; Textile, weaving & dying; Mansabdari/ Land Revenue System: Bandobasta; Babarnama (flora/ fauna); Ain-i-Akbari; crops, plants, fruits of central Asia; Europe | Many temples; Books writing; Gunpowder use in wars; Plague/black death, malaria, dysentery, flu, diphtheria, smallpox typhoid, leprosy, took 200mnlives; portraits paintings; Indo-Islamic architecture excellence. Tajmahal; Quetubminar etc |
| British era (1700 to 070YBP) | British rule | Pre-independence | Cosmetics/Perfumes; Dalton minimum | Great Bengal famine; Cholera; Plague; Yersinia |

| | | | | |
|--|--|--|--|--|
| | | | 1800-1820; New technologies; Good ISM; Lagoons squeezed; Industrial revolution, good rain, High rise population; less storms/disasters | pastis bact. H1N1 Pandemic; fire guns; Urbanization, |
|--|--|--|--|--|

Source: <https://www.clearias.com/indian-history-chronology/>; <https://humanjourney.us/ideas-that-shaped-our-modern-world-section/early-civilizations-harappa/>; <http://www.igntu.ac.in/eContent/IGNTU-eContent-374229503877-BA-AIHC-6-DrJanardhanaB-ScienceandTechnologyinAncientIndia-3.pdf>

8.0 Discussion:

The stalagmites in caves that restructure temperature anomalies from cave carbonates, the changes in the $\delta^{18}\text{O}$ values and the triple oxygen isotopic composition can predict their age (Laskar et al., 2021^[89]). The rapid climate changes (CC) have invited during the Meghalayan age has invited global warming with rise in SST and SAT, Ice sheet melting in glaciers, reduced polar ice sheet, MSLR/RSLR, Carbon sequestration, Extreme events, ocean acidification and many others (Nerem et al., 2018^[90]).

The hunter and gatherers of pre-Holocene started staying in river valleys with nucleated settlements with primary source of livelihood as agriculture (stable and favourable SW monsoon), and hunting as the secondary profession during late Holocene period (ca. 4.0–2.6KYBP).

Paleontological archives like Copal and resins along with historical evidences framed to construct biostratigraphy. Primal forests like Madagascar, East Africa, and Colombia resins, ambers studied and the declaration of Meghalayan age (4.2KYBP to 1950 AD) was questioned as a quick decision by the International Commission on Stratigraphy (ICS), (Solórzano et al., (2020)^[91]).

There were significant ISM anomalies during 4.2.K event for mentioned 200years (4.3 to 4.1KYBP) i.e. at the beginning of the Meghalayan age. The key players and driving forces can

be down trend in solar insolation, prevalence of cooling phase in the North in higher latitudes, erratic movement of the ITCZ from peninsular India to eastern part of Himalayas, Dominance of westerlies, prolonged La-Nina, weak ENSO, dominance of NIOD, and finally retrograding of Bay of Bengal with RSLR fall.

However [Dutta et al., \(2021\)^{\[92\]}](#) reported about a transition period of one millennium dry phase over Indian subcontinent between Northgrippian and Meghalayan age. Nevertheless, the 4.2KYBP drying event have dominance over NW-India whereas some parts of the Indian peninsula and eastern Himalayas received heavy rainfall, high floods, that helped in formation of Speleothem Stalactites and Stalagmites in the caves of Khasi Jaintia calcareous rock masses of Meghalaya . It was due to influence of long activity of the ITCZ in eastern foothills of Himalayas.

[Scropton et al., \(2020\)^{\[93\]}](#) reported that there were two prominent drought spell during pre-Meghalayan era. The first spell was abrupt 300-year long westerlies based with giving precipitation during winter months, (i.e. 4.26 and 3.97KYBP, associated with the 4.2KYBP event as per paleo hydro-climate studies.

[Rasmussen et al., 2007^{\[4\]}](#), and [Walker 2018^{\[5\]}](#) claimed the classification of tripartite Holocene age as Greenlandian, Northgrippian and Meghalayan based on ice core proxies and Speleothem fabrics in the last 20years has been ratified by Philip Gibbard of Cambridge University (Hindusthan Times on 1st Feb 2022) and incorporated by International Union of Geological Sciences (IUGS). The tagging of the stalagmite has now been tagged a Global Boundary Stratotype Section and Points (GSSP) but yet to be stamped.

9.0 Conclusion:

The multi-proxy records of archeological findings, ISM shifts, Holocene paleo climates, Pollen records, Speleothem findings [reveal](#) about sustainability of civilization is dependent on stable hydro-climate. The study of texture profiles for stalagmite best preserved documentation in constructing geological ages. The palynofacies at lime rocks are the calibrating tools of bio-stratigraphy. The research gap in designation of the stratigraphy towards fag end of the Holocene epoch was less explaining. The relative elevation due to, sediment flow, mineralogical, and geochemical features and paleontological proxies discovered in the Lakadong Limestone of

Triassic period of Mawmluh cave of Meghalaya, has made the ratification process simple along with ice core records of DYE-3, GRIP, NGRIP1, NGRIP2 and GICC05 at Greenland and officially stamped by The International Commission on Stratigraphy (ICS). From the multi-proxies, bio-stratigraphy and Radionuclides the established Greenlandian age from 11.7KYBP event, Northgrippian age from 8.2KYBP event and the Meghalayan age from 4.2KYBP event during the Holocene epoch.

Reference:

1. Lucas Spencer G., 2018, The GSSP Method of Chrono stratigraphy: A Critical Review. *Frontiers in Earth Science*, 6, DOI=10.3389/feart.2018.00191, <https://www.frontiersin.org/article/10.3389/feart.2018.00191>
2. Davydov, V. I., (2020). Shift in the paradigm for GSSP boundary definition. Elsevier, 1-76, <https://www.sciencedirect.com/science/article/pii/S1342937X20301866>; Manuscript_d69ddfeb1eeaaec6035f9eaf2c3d565
3. Head, M.J. Review of the Early–Middle Pleistocene boundary and Marine Isotope Stage 19. *Prog Earth Planet Sci* 8, 50 (2021). <https://doi.org/10.1186/s40645-021-00439-2>
4. Rasmussen, S.O., Vinther, B.M., Clausen, H.B., Andersen, K. K., (2007). Early Holocene climate oscillations recorded in three Greenland ice cores. *Quaternary Sc. Reviews*, 26, 1907-1914, doi:10.1016/j.quascirev.2007.06.015.
5. Walker, M., Head, M.J., Berkelhammer, M., Björck, S., Cheng, H., Cwynar L., Fisher, D., (2018). Formal ratification of the subdivision of the Holocene Series/Epoch (Quaternary System/Period): two new Global Boundary Stratotype Sections and Points (GSSPs) and three new stages/subseries, Episodes, <https://doi.org/10.18814/epiiugs/2018/018016>
6. Mishra S. P., 2017, The apocalyptic Anthropocene epoch and its management in India, *Int. Jour. Adv. Research*, Vol. 5(3), pp. 645-663; DOI: 10.21474/IJAR01/3555
7. Martin, M., Sendra, O., Bastos, A. Bauer, N. and others, (2021). Ten new insights in climate science 2021: A horizon scan, *Global Sustain.*, 4 (2021), p. E25
8. Bojie Fu, Michael E. Meadows, Wenwu Zhao, 2022. Geography in the Anthropocene: Transforming our world for sustainable development. *Geography and Sustainability*, 3(1), 1-6, <https://doi.org/10.1016/j.geosus.2021.12.004>, (<https://www.sciencedirect.com/science/article/pii/S2666683921000791>)
9. Kidder, T., Liu, H., Xu, Q., and Li, M. (2012). The Alluvial Geoarchaeology of the Sanyangzhuang Site on the Yellow River Floodplain, Henan Province, China. *Geoarchaeology* 27, 324–343. doi:10.1002/gea.21411

10. Kathayat, G., Cheng, H., Sinha, A., Yi, L., Li, X., Zhang, H., et al. (2017). The Indian Monsoon Variability and Civilization Changes in the Indian Subcontinent. *Sci. Adv.* 3 (12), e1701296. doi:10.1126/sciadv.1701296
11. Jacobson, M. J., Flohr, P., Gascoigne, A., Leng, M. J., Sadekov, A., Cheng, H., et al. (2021). Heterogeneous late Holocene climate in the Eastern Mediterranean—The Kocain Cave record from SW Turkey. *Geophysical Research Letters*, 48, e2021GL094733. <https://doi.org/10.1029/2021GL094733>
12. Vyawahare M., (2018). 'Meghalayan Age' makes the state a part of geologic history. *Hindusthan Times*, New Delhi, Jul 18, 2018 09:00 PM IST
13. Amos Jonathan, (2018), Welcome to the Meghalayan Age - a new phase in history. *Climate change*, BBC News, 18 July 2018
14. Walker M.C.J., M. Berkelhammer, S. Björk, et al., (2012). Formal subdivision of the Holocene Series/Epoch: a Discussion Paper by a Working Group of INTIMATE and the Sub commission on Quaternary Stratigraphy (International Commission on Stratigraphy), *J. of Quaternary Science*, 27(7): 649-659.
15. Harries, D.B., Ware, F.J., Fischer, C.W., Biswas, J., Kharpran B.D., (2008). A Review of the bio speleology of Meghalaya, India. *J. of Cave and Karst Studies*, 70(3), 163–176.
16. Singhvi, A. K., Kale V.S., (2008). *Paleoclimate Studies in India: Last Ice Age to the Present*. IGBP-WCRP-SCOPE-Report Series-4, 1-41, Publisher; Indian National Science Academy, Bahadur Shah Zafar Marg, New Delhi-110002
17. Laskar, A. H., Bohra, A., 2021. Impact of Indian Summer Monsoon Change on Ancient Indian Civilizations during the Holocene. *Front. Earth Sci.*, 06 September 2021 | <https://doi.org/10.3389/feart.2021.709455>
18. Peppe, D. J. & Deino, A. L. (2013) Dating Rocks and Fossils Using Geologic Methods. *Nature Education Knowledge* 4(10):1
19. Haas JN, Richoz I, Tinner W, Wick L. Synchronous Holocene climatic oscillations recorded on the Swiss Plateau and at timberline in the Alps. *The Holocene*. 1998; 8(3):301-309. doi:10.1191/095968398675491173
20. Rasmussen, S. O., K. K. Andersen, A. M. Svensson, et. al. (2006), A new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res.*, 111, D06102, doi:10.1029/2005JD006079.
21. Kalpana, MS, Routh, J., Fietz, S., Lone, MA. Mangini A, (2021). Distribution and Paleo-environmental application of fatty acids in speleothem deposits from Krem Mawmluh, Northeast India. *Frontiers in Earth Sci.*, 1-16, <https://doi.org/10.3389/feart.2021.687376>
22. Dildi, D., Pandey N., (2021). Petrographic analysis of Krem (cave) Mawmluh stalagmite from Meghalaya, northeast India, *J. of Earth System Sc.* 130(4):18, DOI: 10.1007/s12040-021-01697-w

23. Weiss H., Courty MA, Wetterstrom W., Guichard F., Senior L., Meadow R., Curnow A. (1993). The genesis and collapse of third millennium North Mesopotamian civilization. *Science*. 20; 261(5124):995-1004. Doi: 10.1126/science.261.5124.995. PMID: 17739617
24. Cullen, H. M., deMenocal P. B., Hemming, S., Hemming, G., et. al., (2000). Climate change and the collapse of the Akkadian empire: Evidence from the deep sea. *Geology*; 28, 4, 379–382;
25. Kotlia, B. S., Singh, A. K., Joshi, L. M., Dhaila, B. S., Kolita et al., (2015), : Precipitation variability in the Indian Central Himalaya during last ca. 4000 years inferred from a speleothem record: Impact of Indian Summer Monsoon (ISM) and Westerlies, *Quatern. Int.*, 371, 244–253, <https://doi.org/10.1016/j.quaint.2014.10.066>, 2015.
26. Kong, W., Swenson, L. M., and Chiang, J. C. H.: Seasonal Transitions and the Westerly Jet in the Holocene East Asian Summer Monsoon, *J. Climate*, 30, 3343–3365, <https://doi.org/10.1175/JCLI-D-16-0087.1>, 2017.
27. Weiss, H., 2017, 4.2 ka BP Megadrought and the Akkadian Collapse. In: Weiss, H. (Ed.), *Megadrought and Collapse. From Early Agriculture to Angkor*. Oxford University Press, Oxford, pp. 93–160
28. Quamar MF, Kar R, Thakur B., (2021).Vegetation response to the Indian Summer Monsoon (ISM) variability during the Late-Holocene from the central Indian core monsoon zone. *The Holocene*. 31(7):1197-1211. doi:10.1177/09596836211003191
29. Singh, I. B., Sharma, S., Sharma, M., Srivastava, P., and Rajagopalan, G. (1999). Evidence of Human Occupation and Climate of 30 Ka in the Alluvium of Southern Ganga Plain. *Curr. Sci.* 76 (7), 1022–1026.
30. Bamshad, M., Kivisild, T., Watkins, W. S., Dixon, M. E., Ricker, C. E., Rao, B. B., Naidu, J. Met. Al., Jorde, L. B. (2001). Genetic evidence on the origins of Indian caste populations. *Genome research*, 11(6), 994–1004. <https://doi.org/10.1101/gr-1733rr>
31. Banerji, U., Bhusan, R., and Jull, A. J. T. (2017). Mid-late Holocene Monsoonal Records from the Partially Active Mudflats of Diu Island, Southern Saurashtra, Gujarat, Western India. *Quat. Int.* 443, 200–201. doi:10.1016/j.quaint.2016.09.060
32. Clarkson, C., Harris, C., Li, B. et al. Human occupation of northern India spans the Toba super-eruption ~74,000 years ago. *Nat Commun* 11, 961 (2020). <https://doi.org/10.1038/s41467-020-14668-4>
33. Callaway E., 2021. Oldest DNA from a Homo sapiens reveals surprisingly recent Neanderthal ancestry. *Nature*, Apr, 592 (7854):339. doi: 10.1038/d41586-021-00916-0.
34. Chiotis Eustathios, (2018). *Climate Changes in the Holocene, Impacts and Human Adaptation*. [edited by] Eustathios Chiotis. Description: Boca Raton: CRC Press, Taylor & Francis Group, 2018.

35. Wang, YJ., Cheng, H., Edwards, RL., He, YQ., Kong, XG., An, ZS., et al., (2005). The Holocene Asian monsoon: Links to solar changes and North Atlantic climate. 308 (5723) 854-857, DOI: 10.1126/science.1106296
36. Borzenkova I. Zorita, E., Borisova, O., Kalniņa L., et. al. (2015). Climate Change During the Holocene (Past 12,000 Years). In: The BACC II Author Team (eds) Second Assessment of Climate Change for the Baltic Sea Basin. Regional Climate Studies. Springer, Cham. https://doi.org/10.1007/978-3-319-16006-1_2
37. Mishra S. P., Ojha, Ananta Charan, 2020, Analysis and Prediction of Upsurge in Cyclogenesis over the Arabian Sea Fabric, Journal of Xidian University, ISSN 1001-2400pp- 1275-1286Volume 14(5), 2020, <https://doi.org/10.37896/jxu14.5/140>
38. Sun, A.; Feng, Z., (2013). Holocene climatic reconstructions from the fossil pollen record at Qigai Nuur in the southern Mongolian Plateau, Holocene, 23, 1391–1402.
39. Jenny, J.P Koirala, S., Eaves, I. G., et al., (2019), Human and climate global-scale imprint on sediment transfer during the Holocene. Proceedings of the National Academy of Sci., 116 (46) 22972-22976; DOI: 10.1073/pnas.1908179116
40. Kaufman D, McKay N, Routson C, Erb M, Davis B, Heiri O, Jaccard S, et. al., (2020). A global database of Holocene paleo-temperature records. Sci Data. 2020 Apr 14; 7(1):115. doi: 10.1038/s41597-020-0445-3. Erratum in: Sci Data. 2020 June-Aug):271.
41. Hao, Q.; Yang, S.; Song, Z.; Wang, Z.; Yu, C.; Wang, H. Vegetation Determines Lake Sediment Carbon Accumulation during Holocene in the Forest–Steppe Ecotone in Northern China. Forests 2021, 12, 696. <https://doi.org/10.3390/f12060696>
42. Park, J., Park, J., Yi, S. et al. (2019). Abrupt Holocene climate shifts in coastal East Asia, including the 8.2 ka, 4.2 ka, and 2.8 ka BP events, and societal responses on the Korean peninsula. Sci Rep 9, 10806. <https://doi.org/10.1038/s41598-019-47264-8>
43. Fairbridge, R. (1976). Effects of Holocene Climatic Change on some Tropical Geomorphic Processes. Quaternary Research, 6(4), 529-556. doi:10.1016/0033-5894(76)90025-9
44. Park, J. et al. (2018). The 8.2 ka cooling event in coastal East Asia: High-resolution pollen evidence from southwestern Korea. Scientific Reports 8, 12423 (2018).
45. Mishra Siba Prasad, Seth K. C., 2021, The imprints of Holocene climate and environmental changes in the South Mahanadi Delta and the Chilika lagoon, Odisha, India—An overview, In book: Holocene Climate Change and Environment, ELSIVIER, 457-482, <https://doi.org/10.1016/B978-0-323-90085-0.00015-2>
46. Goliath G., Sera Lathan, P., 2005, Rapid erosion of the coast of Sager island, West Bengal – India. Envy. Earth Sci., 48(8):1058-1067, DOI: 10.1007/s00254-005-0044-9
47. Rao, S., Rao Konkani M., N., Vaidyanadhan, R. (1978): Morphology and evolution of Mahanadi and Brahmani Baitarani Deltas. Proceedings of Symposium on Morphology and Evolution of Landforms, Dept. Geology, Delhi University, India, pp. 241-248.

48. Mahalik N.K., Das C. and Wataru Maejim A, "Geomorphology and evolution of the Mahanadi delta, India", Journal of Geoscience, Osaka City University, Vol-39(6) pp- 111-122, 1996.
49. Mohanti M. Swain M. R., (2005), Mahanadi river delta, east coast of India : an overview on evolution and dynamic processes. India Water Portal,1-9
50. Mishra, S. P., Nanda R. N., Mishra S., Sethi K. C.,2021, Anthropocene Physiography and Morphology of Chilika; India. Annual Research & Review in Biology 36(2): 71-95, 2021; Article no.ARRB.66525 ISSN: 2347-565X, NLM ID: 101632869 (NAAS)
51. Venkatrathnam, K., "Formation of the Barrier spit and other sand ridges near Chilika Lake on the east coast of India, Marine Geology", Vol.9, Issue 2, P 101-106 , (Sept 1970), Elsivier.
52. Rao Nageswara K, Takayasu Katsumi, Sadakata N, Hemamalini B., "Holocene evolution of deltas on the east coast deltas of India, present status and future prospects", Kay R, editor, Deltas of the World. New York: American Society of Civil Engineers, 1 – 15, 1993.
53. Sethi B. K., Mishra S. P., Sethi K/ K., Barik K. K., Deltaic Expansions of the Mahanadi Tri-delta and the Chilika Lagoon: Geospatial Approach; Current Journal of Applied Science and Technology 39(34):46-65;(2020); DOI: 10.9734/CJAST/2020/v39i3431036
54. Khandelwal, A., Mohanty, M., Rodriguez F. G., Scharf W. S., "Vegetation history and sea level variation during the last 13,500 years inferred from a pollen record at Chilika Lake, Orissa, India", Vegthist Archaeobot Vol.17, 335-344 Springer verlag-07, 2007
55. Borzenkova I., Zorita, E., Borisova, O., Kalniņa, L., et al. (2015) Climate Change During the Holocene (Past 12,000 Years). In: The BACC II Author Team (eds) 2nd Assessment of Climate Change for the Baltic Sea Basin. Regional Climate Studies. 25-49, Springer, Cham. https://doi.org/10.1007/978-3-319-16006-1_2
56. Pillai, AAS, Anoop, A, Prasad, V., (4 more) et al. (2018). Multi-proxy evidence for an arid shift in the climate and vegetation of the Banni grasslands of western India during the mid-to late-Holocene. Holocene, 28 (7),1057-1070, doi.org/10.1177/0959683618761540
57. Kobashi, T., Goto-Azuma, K., Box, J.E., GAO, C. C., Nakaegawa, T. (2013). Causes of Greenland temperature variability over the past 4000 yr: implications for northern hemispheric temperature changes. Clim. of the Past, 9(5), 2299-2317. doi: 10.5194/cp-9-2299-2013
58. Mishra S. P., Mishra S. and Siddique M. D., 2020; The Anthropocene Dialogues on Climate Change to Human Health of Homosapiens in India; Current J. of Applied Sci. and Tech., 39(24): 13-30, 2020; DOI: 10.9734/CJAST/2020/v39i2430869
59. Scott M., (2021). What is the hottest Earth has been "lately"? NOAA, Climate. Government, Updated Sept. 7, 2021

60. Ramisch, A., Lockot, G., Haberzettl, T., Haberzettl, T., Hartmann, K., et al. (2016). A persistent northern boundary of Indian Summer Monsoon precipitation over Central Asia during the Holocene. *Sci Rep* 6, 25791 <https://doi.org/10.1038/srep25791>
61. Quamar Md. Firoze (2021) Monsoonal climatic reconstruction from Central India during the last ca. 3600 cal yr: signatures of global climatic events, based on lacustrine sediment pollen records, *Palynology*, DOI: 10.1080/01916122.2021.1930605
62. Binkley M. (1987) Azores (Bermuda) high. In: *Climatology. Encyclopedia of Earth Science*. Springer, Boston, MA. https://doi.org/10.1007/0-387-30749-4_24
63. Yadav, R.K., (2021) Relationship between Azores High and Indian summer monsoon. *npj Clim Atmos Sci* 4, 26, <https://doi.org/10.1038/s41612-021-00180-z>
64. Royden, L.H., Burchfiel, B.C., Hilst, R.V.D., Whipple, K.X., et al., (2006). Uplift and evolution of the eastern Tibetan plateau. Presented in the Geological Society of America Philadelphia, Annual Meeting 2006, session 109-1
65. Berkelhammer, M.B., Sinha, A., Stott, L., Cheng, H., Pausata, F.S.R., Yoshimura, K. (2012). An abrupt shift in the Indian Monsoon 4000 years ago. *Geophysical Monographs Series*, v.198, pp.75–87.
66. Kumar K, Agrawal S, Sharma A, Pandey S. (2019). Indian summer monsoon variability and vegetation changes in the core monsoon zone, India, during the Holocene: A multi-proxy study. *The Holocene*. 29 (1):110-119. doi:10.1177/0959683618804641
67. Banerji US, Arulbalaji P, Padmalal D. Holocene climate variability and Indian Summer Monsoon: An overview. *The Holocene*. 2020; 30(5):744-773. doi:10.1177/0959683619895577
68. Cronin TM, Clevenger MK, Tibert NE, et al. (2019). Holocene sea-level variability from Chesapeake Bay Tidal Marshes, USA. *The Holocene*. 29 (11):1679-1693. doi:10.1177/0959683619862028
69. Yan T, Yu K, Wang R, Liu W, Jiang L, (2021). Records of sea-level high-stand over the Meghalayan age/late Holocene from uranium-series ages of beach rock in Weizhou Island, northern South China Sea. *The Holocene*. 31(11-12):1745-1760. doi:10.1177/09596836211033215
70. Loveson, V.J., Nigam, R., 2019. Reconstruction of Late Pleistocene and Holocene Sea Level Curve for the East Coast of India. *J Geol Soc India* 93, 507–514 , doi:10.1007/s12594-019-1211-z
71. Kumar D. S, Gangopadhyay K, Ghosh A, Biswas, O., et al. (2021). Organic geochemical and palaeo botanical reconstruction of a late-Holocene archaeological settlement in coastal eastern India. *The Holocene*. 31(10):1511-1524. doi:10.1177/09596836211025970
72. Farooqui A., Bsip, R., Nautiyal C. M., (2016). Deltaic land subsidence and sea level fluctuations along the east coast of India since eight ka: A palynological study. *The Holocene* 26(9), DOI: 10.1177/0959683616640040

73. Yadav, R.K., Srinivas, R.K.G., Chowdary J.S., (2018), Atlantic Niño modulation of the Indian summer monsoon through Asian jet; *Clim. Atmos. Sci.*, 1, 10.1038/s41612-41018-40029-41615
74. Mishra S. P., July 2020; Human Evolution/Extinction up to Present Anthropocene: India; *Journal of Shanghai Jiaotong University*; JSJ.U-2222.14-F (1).pdf; ISSN:1007-1172; 16 (7); 115-133
75. Mishra S. P., Mishra S. and Siddique M. D., 2020; The Anthropocene Dialogues on Climate Change to Human Health of Homosapiens in India; *Current Journal of Applied Science and Technology* 39(24): 13-30, 2020;
76. Phartiyal B., Nag D., Joshi P., (2022). Holocene climatic record of Ladakh, Trans-Himalaya. Ed.: Kumaran N., Damodara P., *Holocene Climate Change and Environment*, Elsevier, 61-89, 2022, doi = {<https://doi.org/10.1016/B978-0-323-90085-0.00023-1>
77. Luthi, D., et al., 2008; Etheridge, D.M., et al. 2010; Vostok ice core data/J.R. Petit et al.; NOAA Mauna Loa CO2 record.
78. Bonnefille, R., Anupama, K., Barboni, D., Pascal, J., Prasad, S., Sutra, JP. (1999). Modern Pollen Spectra from Tropical South India and Sri Lanka: Altitudinal Distribution. *Journal of Biogeography*, 26(6), 1255–1280. <http://www.jstor.org/stable/2656066>
79. Lovelock, C. E., I. C. Feller, R. Reef, S. Hickey & M. C. Ball, 2017b. Mangrove dieback during fluctuating sea levels. *Scientific Reports* 7: 1680.
80. Pandey, S., Scharf, B.W., Mohanti M., 2014, Palynological studies on mangrove ecosystem of the Chilka Lagoon, east coast of India during the last 4165 yrs. BP. *Elsivier, Quarterly Int.*, 325, 126-135, DOI: 10.1016/j.quaint.2013.09.001
81. Feller, I.C., Friess, D.A., Krauss, K.W. et al. The state of the world's mangroves in the 21st century under climate change. *Hydrobiologia* 803, 1–12 (2017). <https://doi.org/10.1007/s10750-017-3331-z>
82. Mondal, B., Sahaa, A. K., Roy, A., (2021). Spatio-temporal pattern of change in mangrove populations along the coastal West Bengal, India. *Environmental Challenges*, 5, 100306
83. Biswas S. K., (2014). Active tectonic zones of Indian plate and intra-plate dynamics. Dept. of Atomic Energy, Atomic minerals directorate for exploration and research; *Exploration and Research for Atomic Minerals* 24:25-37
84. Kumar, A., Mitra, S., and Suresh, G. (2015), Seismotectonics of the eastern Himalayan and indo-burman plate boundary systems, *Tectonics*, 34, 2279– 2295, doi:10.1002/2015TC003979.
85. Kingwell-B., E., Harvey, Karoune N. E., Mohanty, R.K., Fuller, D.Q., (2018). Archaeobotanical Investigations into Golabai Sasan and Gopalpur, Two Neolithic-Chalcolithic Settlements of Odisha. *Ancient Asia*, 9, 5. DOI: <http://doi.org/10.5334/aa.164>

86. Mishra S. P., Mishra S. K., 2018, The Cataclysm of Geo-Bio-Climate in Short-Lived Holocene and in Anthropocene epochs: A Critical Review, International Journal of Science and Research (IJSR) Vol. 7(9), PP-1445 – 1462, DOI: 10.21275/ART20191537
87. Hazra S., Das S., Ghosh A., Raju P.V., Patel A. (2020). The Mahanadi Delta: A Rapidly Developing Delta in India. In: Nicholls R., Adger W., Hutton C., Hanson S. (Eds) Deltas in the Anthropocene. Palgrave Macmi., Cham. https://doi.org/10.1007/978-3-030-23517-8_3
88. Cheng, J., Wu, H., Liu, Z. et al. Vegetation feedback causes delayed ecosystem response to East Asian Summer Monsoon Rainfall during the Holocene. Nat Commun 12, 1843 (2021). <https://doi.org/10.1038/s41467-021-22087-2>
89. Laskar, A. H., Bohra A., (2021). Impact of Indian Summer Monsoon Change on Ancient Indian Civilizations During the Holocene. Earth Sci., <https://doi.org/10.3389/feart.2021.709455>
90. Nerem, R. S., Beckley, B. D., Fasullo J. T., Hamlington, B. D., Masters D., Mitchum G. T., (2018). "Climate-change–driven accelerated sea-level rise detected in the altimeter era." PNAS, 2018 DOI: 10.1073/pnas.1717312115
91. Solórzano-Kraemer, M.M., Delclòs, X., Engel, M.S. et al., (2020). A revised definition for copal and its significance for palaeontological and Anthropocene biodiversity-loss studies. Sci Rep 10, 19904 <https://doi.org/10.1038/s41598-020-76808-6>
92. Dutt, S., Gupta, A. K., , Devrani, R., Yadav R. R., Singh, R. K., (2021), Regional disparity in summer monsoon precipitation in the Indian subcontinent during North-grippian to Meghalayan transition. Current Science, 120 (9), 1449-1458
93. Scroxton, N., Burns, S. J., McGee, D., Godfrey, L. R., Ranivoharimanana, L., and Faina, P.: Circum-Indian ocean hydro-climate at the mid to late Holocene transition: The Double Drought hypothesis and consequences for the Harappan, Clim. Past Discuss. [Preprint], <https://doi.org/10.5194/cp-2020-138>, 2020.