

Effect of zinc and iron on growth and productivity of summer mungbean

Abstract

Micronutrients like zinc and iron play an important role in growth and development of mungbean. The scarcity of these micronutrients also has an impact on seed quality of mungbean. A field experiment on optimization of nutrient management practices through soil and foliar application in mungbean was conducted during summer season 2021. The treatments included control, 0.5% ZnSO₄ at flower initiation (FI), 0.5% ZnSO₄ at FI & pod initiation (PI), 0.5% FeSO₄ at FI, 0.5% FeSO₄ at FI & PI, 0.5 % ZnSO₄ & 0.5% FeSO₄ at FI, 25 kg ha⁻¹ ZnSO₄ soil application, 50 kg ha⁻¹ FeSO₄ soil application. Plant height (29.85 cm), number of branches per plant (8.43) and number of nodules per plant (16) were improved significantly by foliar spraying of 0.5% ZnSO₄ and 0.5% FeSO₄ at FI. Significantly maximum no. of pods per plant, seeds per pod, seed yield (521 kg ha⁻¹) and stover yield (1307 kg ha⁻¹), net returns (Rs. 8020/ha) and benefit cost ratio (1.28) were recorded under 0.5% ZnSO₄ and 0.5% FeSO₄ spray at Flower initiation stage.

Keywords: Mungbean, foliar spray, zinc, iron, growth, yield and yield attributes, economics.

Introduction

Mungbean (*Vigna radiata* L.) is an important pulse crop grown during summer season. It is commonly called as “Moong Dal” in India. Moong dal is a widely used dal that serves both culinary and medicinal uses. It is well known for its high protein (22-24%), fibre and iron content. It also fixes atmospheric nitrogen and improves soil fertility by adding 20-25 N kg ha⁻¹. The yield of summer mungbean is more than that of *kharif* crop because of the abundance sunlight, controlled moisture conditions provided by irrigation, less pest and disease infestation.

Micronutrients like zinc and iron play an important role in growth and development of mungbean. The scarcity of these micronutrients also has an impact on seed quality of mungbean.

Zinc is an essential vitamin which is required by mungbean plant involved in auxin production, dehydrogenase enzyme activation, ribosomal fractions stabilization Babar, H. et al. [1]. The plant's photosynthetic activity is severely hampered by a shift in chloroplast pigments caused by Zn shortage. Application of zinc increased the seedling growth, morphological and yield parameters, grain yield and grain zinc concentration of mungbean Haider, M.U. et al. [2]. Iron is found in several important proteins, including nitrogenase, leghaemoglobin and ferredoxin. It is involved in chlorophyll and thylakoid production as well as chloroplast formation. In legumes, iron deficiency inhibited nodule formation and development. As a result, iron deficiency in legumes is also a clear indicator of nitrogen inadequacy. The application of FeSO_4 plays an important role in chlorophyll synthesis and plant growth regulator Jin, Z. et al. [3]. Foliar application of iron solutions in many crops, including legumes, is one of the most commonly used methods for correcting iron deficiency.

Considering the view of above facts, the field investigation was conducted to determine the influence of zinc and iron on growth and productivity of summer mungbean.

Materials and Methods

The experiment was conducted at the field of Regional Research and Technology Transfer Sub Station (RRTTSS), Kirei, Sundargarh under North Western Plateau Zone of Odisha, during summer season of 2021. The mungbean cultivar used in experiment was 'IPM 02-14'. The experiment was carried out in Randomized Block Design (RBD) with three replications. The soil of the experimental field was sandy loam in texture, acidic in reaction (pH 5.7), low in available nitrogen ($145.41 \text{ kg ha}^{-1}$), medium in available phosphorus (35.13 kg ha^{-1}), low in available potassium (71.68 kg ha^{-1}), low in zinc and iron content of 0.53 and 4.23 ppm, respectively. In this investigation the treatments included T_1 control (RDF), T_2 (0.5% ZnSO_4 at flower initiation), T_3 (0.5% ZnSO_4 at flower initiation & pod initiation), T_4 (0.5% FeSO_4 at flower initiation), T_5 (0.5% FeSO_4 at flower initiation & pod initiation), T_6 (0.5% ZnSO_4 and 0.5% FeSO_4 at flower initiation), T_7 (25 kg ha^{-1} ZnSO_4 through soil application) and T_8 (50 kg ha^{-1} FeSO_4 through soil application). A uniform dose of NPK 20-40-20 kg ha^{-1} and FYM 5 tha^{-1} were applied as basal to all the treatments.

The mungbean crop was sown on 9th February, 2021 in furrows at a spacing of 30×10 cm. Before sowing, seed @ 20 kg ha⁻¹ was treated with Vitavax power @ 2 g kg⁻¹ of seed. Zinc sulphate and iron sulphate were applied 0.5% foliar application at flower initiation and pod initiation separately and both the stages. Hoeing and hand weeding were done at 20 and 35 days after sowing. Irrigation was given prior to the preparation of the field and subsequent irrigations were given after every 15 days from the date of sowing. The crop was harvested on 21st April, 2021. Important observations were recorded at the appropriate time and economics were estimated using market price.

Results and Discussion

Growth parameters: It was observed from the results that different nutrient management practices had significant effect on growth parameters of mungbean in terms of plant height (29.85 cm), number of branches per plant (8.43) and number of nodules per plant (16) with foliar application of 0.5% ZnSO₄ and 0.5% FeSO₄ at flower initiation stage. It might be due to positive response of zinc in the production of IAA, sugar translocation and response of Fe as a cofactor of many enzymes help in various physiological processes especially affects photosynthesis and chlorophyll synthesis, cell division, water and nutrients absorption, finally help in improving various plant growth parameters. Corroborative results were given by Sitaram, T. et al. [4], Meena, K.K. et al. [5], Ali, B. et al. [6], Khan and Prakash [7], Lakshmi, E.J. et al. [8], Saini, A.K. [9], Suhathiya and Ravichandran [10] and Masih, A. et al. [11].

Yield attributes and yield: Yield attributes like pod length, 1000-seeds weight showed non-significant effect on different treatments of zinc and iron. Yield attributes viz. pods per plant (15.63) and seeds per pod (10.50) were observed significantly higher in foliar spray of zinc and iron sulphate i.e. 0.5% ZnSO₄ and 0.5% FeSO₄ at flower initiation stage. These results attributed to stimulatory effect of zinc and iron on most of the physiological and metabolic processes and nitrogen metabolism, synthesis of chlorophyll, plant growth regulator, improves photosynthesis and assimilates transportation to sink resulted enhanced seed yield of mungbean. The foliar application of zinc sulphate and iron sulphate at flower initiation and bud initiation stage are responsible for efficient translocation of photosynthate from source to sink, this causes' higher number of pod formation and higher test weight. These results are in accordance with Awlad,

H.M. et al. [12] and Teixeira, I.R. et al. [13], Kumawat, R.N. et al. [14], Meena, K.K. et al. [5] and Soni and Kushwaha [15].

Seed yield (521 kg ha^{-1}) and stover yield (1307 kg ha^{-1}) was found to be higher with application of 0.5% ZnSO_4 and 0.5% FeSO_4 at flower initiation stage, that might be due to significant effect on yield attributing characters like number of pod bearing branches per plant, pods per plant and 1000-seeds weight. Similarly higher yield attributes and yield were noticed with the combined foliar spray of micro nutrients with zinc attributed to optimum availability of nutrients for luxurious crop growth and efficient partitioning of assimilates from source to sink. Higher stover yield of mungbean might be due to direct influence of zinc on auxin production which in turn enhanced the elongation processes of plant development Masih, A. et al. [11], Prasanna, K.L. et al. [16] and Choudhary, P. et al. [17].

Economics: The highest cost of cultivation ($\text{Rs.}30095 \text{ ha}^{-1}$) was observed in 50 kg ha^{-1} FeSO_4 soil application. This was because of higher quantity of applied FeSO_4 as basal compared to foliar feeding treatments. Gross returns ($\text{Rs.}36470 \text{ ha}^{-1}$) and net returns ($\text{Rs.}8020 \text{ ha}^{-1}$) and benefit cost ratio (1.28) was recorded highest under treatment 0.5% ZnSO_4 and 0.5% FeSO_4 at flower initiation stage. Foliar application of 0.5% ZnSO_4 at flower initiation and 0.5% ZnSO_4 at flower initiation and pod initiation earned more net returns by $\text{Rs. } 7265 \text{ ha}^{-1}$ and $\text{Rs. } 6940 \text{ ha}^{-1}$ than control, however, all foliar fertilized treatment exhibited greater net returns over control. This increase in net return was due to significantly higher value of seed and stover yield as compared to cost of cultivation of corresponding treatment. The results were supported by Soni and Kushwaha [15], Sharma, A. et al. [18] and Sajid, M. et al. [19].

Conclusion

From the results of above study, it was concluded that foliar spray with 0.5% ZnSO_4 and 0.5% FeSO_4 at flower initiation stage in summer mungbean produced maximum seed (521 kg ha^{-1}) and stover yield (1307 kg ha^{-1}) with maximum net return of $\text{Rs. } 8020 \text{ ha}^{-1}$ and benefit cost ratio (1.28) because at this level we recorded maximum plant growth and yield of mungbean as compared to control. So this nutrient management practice can be recommended to farmers for getting higher productivity and profitability of summer mungbean.

Table 1 Effect of zinc and iron on growth attributes of summer mungbean

| Treatment | Plant height (cm) at harvest | No. of branches/plant at harvest | No. of nodules/plant at 45 DAS |
|--|------------------------------|----------------------------------|--------------------------------|
| T₁ - Control (RDF: 20-40-20 kg/ha and FYM 5 tha⁻¹) | 24.17 | 4.43 | 9.67 |
| T₂ - T₁+ 0.5% ZnSO₄ FI | 28.53 | 8.10 | 14.37 |
| T₃ - T₁+ 0.5% ZnSO₄ FI & PI | 28.90 | 8.27 | 15.13 |
| T₄ - T₁+ 0.5% FeSO₄ FI | 27.23 | 6.87 | 12.03 |
| T₅ - T₁+ 0.5% FeSO₄ FI & PI | 28.23 | 7.77 | 12.87 |
| T₆ - T₁+ 0.5% ZnSO₄ and 0.5% FeSO₄ FI | 29.85 | 8.43 | 16.00 |
| T₇ - T₁+ 25 kg/ha ZnSO₄ soil application | 26.53 | 6.73 | 11.87 |
| T₈ - T₁+ 50 kg/ha FeSO₄ soil application | 25.60 | 6.07 | 11.10 |
| SE (m)± | 0.41 | 0.20 | 0.33 |
| CD (P=0.05) | 1.22 | 0.60 | 0.98 |

*RDF: Recommended dose of fertilizer, *FI: Flower initiation, *PI: Pod initiation, *SEm: Standard error of the mean

Table 2 Effect of zinc and iron on yield attributes and yield of summer mungbean

| Treatment | No. of pods/plant | Pod length (cm) | No. of seeds /pod | 1000-seed weight (g) | Seed yield (kg/ha) | Stover yield (kg/ha) |
|--|-------------------|-----------------|-------------------|----------------------|--------------------|----------------------|
| T₁ - Control (RDF: 20-40-20 kg/ha and FYM 5 tha⁻¹) | 9.87 | 6.17 | 8.00 | 29.80 | 402.3 | 1074.7 |
| T₂ - T₁+ 0.5% ZnSO₄ FI | 14.77 | 6.60 | 10.10 | 30.73 | 502.3 | 1274.7 |
| T₃ - T₁+ 0.5% ZnSO₄ FI & PI | 15.03 | 7.03 | 10.30 | 31.40 | 511.3 | 1296.0 |
| T₄ - T₁+ 0.5% FeSO₄ FI | 12.97 | 6.57 | 9.33 | 29.50 | 456.0 | 1178.7 |
| T₅ - T₁+ 0.5% FeSO₄ FI & PI | 13.77 | 6.63 | 10.00 | 30.13 | 450.3 | 1169.0 |
| T₆ - T₁+ 0.5% ZnSO₄ and 0.5% FeSO₄ FI | 15.63 | 7.03 | 10.50 | 31.27 | 521.0 | 1307.0 |
| T₇ - T₁+ 25 kg/ha ZnSO₄ soil | 12.90 | 6.50 | 9.20 | 30.27 | 446.7 | 1195.0 |

| application | | | | | | |
|--|-------|------|------|-------|-------|--------|
| T₈ - T₁+ 50 kg/ha FeSO₄ soil application | 12.40 | 6.20 | 8.33 | 28.93 | 439.0 | 1202.7 |
| SE (m)± | 0.38 | 0.21 | 0.17 | 0.57 | 9.63 | 12.20 |
| CD (P=0.05) | 1.12 | N/S | 0.52 | N/S | 28.60 | 36.25 |

*RDF: Recommended dose of fertilizer, *FI: Flower initiation, *PI: Pod initiation, *SEm: Standard error of the mean

Table 3 Effect of application of zinc and iron on production economics of summer mungbean

| Treatment | Cost of cultivation (Rs./ha) | Gross return (Rs./ha) | Net return (Rs./ha) | B:C ratio |
|--|------------------------------|-----------------------|---------------------|-----------|
| T₁ - Control (RDF: 20-40-20 kg/ha and FYM 5 tha⁻¹) | 27595 | 28140 | 545 | 1.02 |
| T₂ - T₁+ 0.5% ZnSO₄ FI | 28200 | 35140 | 6940 | 1.25 |
| T₃ - T₁+ 0.5% ZnSO₄ FI & PI | 28505 | 35770 | 7265 | 1.25 |
| T₄ - T₁+ 0.5% FeSO₄ FI | 28145 | 31920 | 3775 | 1.13 |
| T₅ - T₁+ 0.5% FeSO₄ FI & PI | 28445 | 31500 | 3055 | 1.11 |
| T₆ - T₁+ 0.5% ZnSO₄ and 0.5% FeSO₄ FI | 28450 | 36470 | 8020 | 1.28 |
| T₇ - T₁+ 25 kg/ha ZnSO₄ soil application | 29645 | 31290 | 1645 | 1.06 |
| T₈ - T₁+ 50 kg/ha FeSO₄ soil application | 30095 | 30730 | 635 | 1.02 |
| SE (m)± | 8.12 | 623.80 | 618.03 | 0.02 |
| CD (P=0.05) | 24.88 | 1910.44 | 1892.78 | 0.06 |

*RDF: Recommended dose of fertilizer, *FI: Flower initiation, *PI: Pod initiation, *SEm: Standard error of the mean, *Rs: Rupees

References

1. Babar H, Yusop MK. and Saleem M. Role of zinc in plant nutrition- a review. American Journal of Experimental Agriculture. 2013;3(2):374-391.
2. Haider MU, Hussin M, Farooq M and Nawaz A. Soil application of zinc improves the growth, yield and grain zinc biofertilization of mungbean. Soil and Environment. 2018;37(2):123-128.

3. Jin Z, Minyan W, Lianghuan W and Jiangguo W. Impacts of foliar iron and boron application on iron biofertilization and nutritional quality of rice grain, *Journal of Plant Nutrition*. 2008; 31(9): 1599-1611.
4. Sitaram T, Sharma SK and Reager ML. Growth attributes and nutrient uptake of green gram as influenced by vermicompost and zinc in arid western Rajasthan. *Advanced Research Journal of Crop Improvement*. 2013;4(1):65-69.
5. Meena KK, Meena RS and Kumawat SM. Effect of sulphur and iron fertilization on yield attributes, yield and nutrient uptake of mungbean (*Vigna radiata*). *Indian Journal of Agricultural Sciences*. 2013;83(4): 472-6.
6. Ali B, Ali A, Tahi M, and Ali S. Growth, seed yield and quality of mungbean as influenced by foliar application of iron sulfate. *Pakistan Journal of Life and Social Sciences*. 2014;12(1):20-25.
7. Khan K and Prakash V. Effect of rhizobial inoculation on growth, yield, nutrient and economics of summer urdbean (*Vigna mungo* L.) in relation to zinc and molybdenum. *Journal of Food Legumes*. 2014;27(3):261-263.
8. Lakshmi EJ, Rameshbabu PV, Reddy GP, Umamaheswari P and Reddy APK. Effect of foliar application of secondary nutrients and zinc on growth and yield of blackgram. *International Journal of Chemistry Study*. 2017;5(6):944-947.
9. Saini AK. Effect of iron and sulphur fertilization on growth and yield of greengram [*Vigna radiata* L.]. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(4): 1358-1361.
10. Suhathiya K and Ravichandran M. Effect of foliar application of DAP and micronutrients on crop growth attributes and yield of rice fallow blackgram (*Vigna mungo* L.). *International Journal of Advanced Research Biological Science*. 2018;5(4):72-76.
11. Masih A, Dawson J and Singh RE. Effect of levels of phosphorus and zinc on growth and yield of greengram (*Vigna radiata* L.). *International Journal of Current Microbiological Applied Science*. 2020;9(10): 3106-3112.
12. Awlad HM, Chowdhury MA, and Tahukde NM. Effect of sulphur and zinc on nodulation, dry matter, yield and nutrient content of soybean. *Pakistan Journal of Biological Sciences*. 2003;6:461-466.
13. Teixeira IR, Boren AB and Araujo GAA. Manganese and zinc leaf application on common bean grown on a cerrado soil. *Science of Agriculture*. 2004;61:77-81.

14. Kumawat RN, Rathore PS, Nathawat NS and Mahatma M. Effect of sulfur and iron on enzymatic activity and chlorophyll content of mungbean. *Journal of Plant Nutrition*. 2006; 29:1451-1467.
15. Soni J and Kushwaha HS. Effect of foliar spray of zinc and iron on productivity of mungbean [*Vigna radiata* (L.) wilczek]. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(1):108-111.
16. Prasanna KL, Naidu SMM, Sumathi V and Nagamani C. Effect of nitrogen and zinc on growth, yield and economics of clusterbean [*Cyamopsis tetragonoloba* (L.)Taub]. *The Andhra Agricultural Journal*. 2013;60(2):260-263.
17. Choudhary P., Jhaharia A. and Kumar R. Influence of sulphur and zinc fertilization on yield, yield components and quality traits of soybean (*Glycine max* L.). *The Bioscan*. 2014;9(1):137-142.
18. Sharma A, Sharma RP, Katoch V and Sharma GD. Influence of vermicompost and split applied nitrogen on growth, yield, nutrient uptake and soil fertility in pole type frenchbean (*Phaseolus vulgaris* L.) in an acid alfisol. *Legume Research*. 2018;41(1):126-131.
19. Sajid M. Influence of zinc as soil and foliar application on growth and yield of okra (*Abelmoschus Esculentus* L.). *International Journal of Agricultural and Environmental Research*. 2016;2.2: 140-145.