

Role of nitrogen and spacing on growth and yield of summer Groundnut

(*Arachis hypogaea* L.)

Abstract: The effect of nitrogen and spacing on summer groundnut traits like growth and yield was studied through an experiment performed in 2021 KVK Farm of SHUATS. The experiment had 9 different treatments, replicated thrice in a Randomized control design. The treatments encompassed three different levels of nitrogen soil application and three different plant spacings. The outcome of the study revealed that in treatment (T₃) the grouping of 30 kg/ha nitrogen + 20 cm × 10 cm spacing resulted in maximum plant height (52.27cm) and in treatment (T₆) the usage of 30 kg/ha nitrogen + 25 cm × 10 cm layout resulted in a significant root nodules per plant (39.57) and dry weight (38.00 g). Treatment (T₂) showed a maximum crop growth rate (31.44 g/m²/day) in the combination of 25 kg/ha nitrogen + 20 cm × 10 cm arrangement and 20-40 DAS interval. Furthermore, treatment (T₆) had a maximum pods per plant (30.80), seed capitulate (2355.6 kg/ha), and haulm yield (3402.3 kg/ha) with 30 kg/ha nitrogen + 25 cm × 10 cm while treatment (T₁) obtained maximum harvest index (42.32%) with the use of 20 kg/ha nitrogen + 20 cm × 10 cm.

Key words: Groundnut, nitrogen, spacing, summer, yield, and yield attributes.

Introduction

Groundnut (*Arachis hypogaea* L.) is an herbaceous crop of Leguminosae family. The plant has central stem upright with many branches which are different from prostrate to almost erect based on the variety. Groundnut, also known as “The king of oilseeds”. In India, it is produced mostly as oilseed crop, covers an area of 40-50% and gives a production of 60 to 70 %. It is first major oil seed crop among all the other oil seed harvests. In India, it covers an area of 85 lakh hectares and produced 84 lakh tones from southern states (Andhra Pradesh, Tamil Nadu, Karnataka), from western and central parts (Gujarat, Maharashtra, Madhya Pradesh), from northern states (Uttar Pradesh, Rajasthan), north western part of the subcontinent i.e., Punjab and from eastern India (Orissa). However, the maximum yield and area of 84% is covered by southern and western regions.

Nitrogen is a significantly important element present in plants. It is included in chlorophyll, nucleotides, proteins, alkaloids, enzymes, hormones and vitamins (Marschner, 1995). It enhances rate of photosynthesis, helps with synthesis of metabolites and its transportation to the seeds. The effect of nitrogen fertiliser addition on soil natural matter status and soil significant characteristics is crucial for agricultural management and crop yield growth. N fertiliser also affects dry matter production, as well as N collection and distribution to various parts of yield plants for development, advancement, and other purposes (Khaliq and Cheema 2005). The deficiency of nitrogen causes restricted growth, and chlorotic leaves. The deficiency of N restricts protein and chlorophyll synthesis which results in premature flowering and limiting the growth cycle. The standing pattern of a plant is responsible for many important factors required for the crop production like light, water, nutrients, and weeds. Apart from this, crop canopy is another factor that helps in intercepting quantity of radiation and gives better yield. The distance between the row is important, as wider distance makes radiation unconsumed and lesser distance makes the crop plants packed and shaded.

However, thin spacing lower the production of the crop plants because of the competitiveness of the crop plants for nutrients and moisture (Kandel *et al.* 2013). Likewise, the row alignment too influences

photosynthetic competence and canopy temperature for a better light interception and greater photosynthetic efficiency, a sustained and uniform orientation as well as crop distribution is needed (Meena *et al.* 20–11). Proper application of spacing along with suitable nutrients will help in attaining maximum productivity within India. All these points are the basis of the present study which was laid out in Zaid 2021.

S.No	Treatment	Treatment Grouping
1.	T ₁	20 kg/ha Nitrogen + 20 cm × 10 cm
2.	T ₂	25 kg/ha Nitrogen + 20 cm × 10 cm
3.	T ₃	30 kg/ha Nitrogen + 20 cm × 10 cm
4.	T ₄	20 kg/ha Nitrogen + 25 cm × 10 cm
5.	T ₅	25 kg/ha Nitrogen + 25 cm × 10 cm
6.	T ₆	30 kg/ha Nitrogen + 25 cm × 10 cm
7.	T ₇	20 kg/ha Nitrogen + 30 cm × 10 cm
8.	T ₈	25 kg/ha Nitrogen + 30 cm × 10 cm
9.	T ₉	30 kg/ha Nitrogen + 30 cm × 10 cm

Materials and methods

The present study was guided and carried out at the KVK farm, SHAUTS, Prayagraj (U.P.) during the Zaid 2021 which is situated at 25° 40'42" north latitude, 81° 85'56" east longitude, and 98 metres above mean sea level (MSL). The investigational field was composed of neutral and deep soil from central Gangetic alluvium. The soil texture was sandy loam and nearly nonaligned in soil effect with pH 7.3. It had little organic carbon (0.57%), accessible N (230kg/ha), obtainable P (32.10kg/ha) and available K (235kg/ha). The combination included three different level of N applications and spacings. The experiment was having 9 varied treatments imitated thrice and carried in randomized block design : T₁ : N 20kg/ha + 20cm × 10cm, T₂ : N 25kg/ha + 20cm × 10cm, T₃ : N 30kg/ha + 20cm × 10cm, T₄ : N 20kg/ha + 25cm × 10cm, T₅ : N 25kg/ha + 25cm × 10cm, T₆ : N 30kg/ha + 25cm × 10cm, T₇ : N 20kg/ha + 30cm × 10cm, T₈ : N 25kg/ha + 30cm × 10cm, T₉ : N 30kg/ha + 30cm × 10cm. the fertilizers used for the experiment were carbamide, ssp, and mop. The crop planted was Kadiri-9 on 10th April 2021. and harvested on 22nd July 2021 (104 DAS).

Results and Discussion

Growth parameters

The results suggest, treatments significantly effects progress and groundnut harvest. Table 1 shows an increase in height of the groundnut. Increase in plant height with increase in nitrogen level might be due to proper nutrition availability and plant water status which resulted in increase in vegetative growth of the plants reported by Hasan *et al.* (2019). Application of 30 kg/ha nitrogen + 20 cm × 10 cm spacing in Treatment (T₃) give maximum height (52.27cm) which is statistically at par to treatment (T₂) height (51.83cm) with usage of 25 kg/ha nitrogen + 20 cm × 10 cm spacing. Similar outcomes were recorded by other researchers Meena *et al.* (2011) and Parameshwara reddy *et al.* (2019). Significantly higher dry weight (38.00 g/plant) and root nodules/plant (39.57) were observed in treatment (T₆) having 30 kg/ha nitrogen + 25 cm × 10 cm spacing which is accurately concurrent with the appliance of 25 kg/ha nitrogen + 25 cm × 10 cm (37.93 g/plant) and 20 kg/ha nitrogen + 25 cm × 10 cm (37.09 g/plant). Treatment (T₆) reports higher root nodules with the usage of 30 kg/ha nitrogen + 25 cm × 10 cm layout which is accurately significantly with the application of 25 kg/ha nitrogen + 25 cm × 10 cm (38.73). Treatment (T₂) have significantly higher crop growth rate (31.44 g/m²/day) from 20-40 DAS interval with the application of 25 kg/ha nitrogen + 25 cm × 10 cm, which is accurately at par with the application of (T₁) 20 kg/ha nitrogen + 20 cm × 10 cm (29.79 g/m²/day), (T₃) 30 kg/ha nitrogen + 20 cm × 10 cm (30.83 g/m²/day), (T₄) 20 kg/ha nitrogen + 25 cm × 10 cm (28.93 g/m²/day) and (T₆) 30 kg/ha nitrogen + 25 cm × 10 cm (28.77 g/m²/day). This might be due to lesser competition amongst plants for space, light and nutrition as well and competition increased simultaneously at closer geometry as evident from data of nutrient concentration in planta. Comparable outcome were revealed by Sree *et al.* (2020).

Yield parameters

The data reported in table 2 depicts maximum pods per plant (30.80) in treatment (T₆) with the usage of 30 kg/ha nitrogen + 25 cm × 10 cm layout. These results were statistically in parallel with the 25 kg/ha nitrogen + 25 cm × 10 cm (30.53) in treatment (T₅) and 20 kg/ha nitrogen + 25 cm × 10 cm (30.56) treatment (T₄). However, no significant difference was observed within the treatment for seed index (100 seed). Treatment (T₆) had a higher seed yield (2355.6 kg/ha) and haulm yield (3402.3 kg/ha) with the application of 30 kg/ha nitrogen + 25 cm × 10 cm spacing. Treatment (T₅) 25 kg/ha nitrogen + 25 cm × 10 cm and (T₄) 20 kg/ha nitrogen + 25 cm × 10 cm were recorded statistically comparable with treatment (T₆) 30 kg/ha nitrogen + 25 cm × 10 cm in both seed yield and haulm yield. All these results report a positive effect of nitrogen on various parameters involved in growth of the crop plant. Parallel findings were described by Sree *et al.* (2020). The maximum harvest index (42.32%) was detected in treatment (T₁) 20 kg/ha nitrogen + 20 cm × 10 cm, which is statistically comparable with application of 20 kg/ha nitrogen + 30 cm × 10 cm (41.89). This might be attributed to early expansion of seed produce in highly dense plant by enhancing growth factors, once the reproductive phase starts, the leads to harvestable crops while other climatic conditions are optimal. Comparable results are stated by Gawas *et al.* (2020). The optimum planting rate of Groundnut diminishes intra plant competition through juvenility, increases the produce of the plant, ground cover, and light seizure, and leads to maximum dry matter and yield (Zhao *et al.*, 2017). This might be accredited to additional resources at the optimal plant density initiated more leaf the optimum plant density initiated more plant height, number of root nodules, rate of crop growth, quantity of pods per plant.

Conclusion

According to the findings of the study, treatment (T₆) produced the highest seed yield (2355.62 kg/ha) and haulm yield (3402.3 kg/ha) as a result of the application of 30 kg/ha nitrogen and 25 cm x 10 cm spacing. The results of the investigation are dependent on a single season; therefore, further investigation is required for confirmation.

Acknowledgment

I acknowledge **Dr. RAJESH SINGH**, my advisor for helping me in preparing and improving this manuscript. Further, I am thankful to the faculty members of the Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh (U.P), India, for their support during this research

References

- Gawas, D., Mane, A.V., Burondkar, M.M., Kanase, T., Kasture, M.C. and Dalavi, V.V. 2020. Yield performance of groundnut (*Arachis hypogaea* L.) genotypes under varying plant densities and seasons. *International Journal of Chemical Studies* **8**(6): 2383-2387.
- Gomez, K.A. and Gomez, A.A. 1984. Statistical Procedures for Agricultural Research. 2nd Ed., Wiley and Sons, Inc. New York, USA.
- Hasan, M., Josh, L. K., and Alam, M. D. 2019. Influence of nitrogen levels on oil seed crops: A review. *Indian Journal of Agricultural Sciences* **4**:29-36.
- Kandel, T.P., Pandey, P.B., Basnet, B.K., Bhatta, M.R., Sah, K.S. and Thapa, R.B. 2013. Effect of row spacing and direction of sowing on yield and yield attributing characters of wheat cultivated in Western Chitwan, Nepal. *Agricultural Sciences* **4**(7): 309-316.
- Kumar S. N. 2009. Effect of plant density and weed management practices on production potential of groundnut (*Arachis hypogaea* L.). *Indian Journal of Agricultural Research* **43**(1):57-60.
- Kumar, S.L., Radder, B.M., Malligawad, L.H. and Manasa, V. 2014. Effect of Nitrogen and Phosphorus levels and ratios on yield and nutrient uptake by Groundnut in Northern Transition Zone of Karnataka. *The Bioscan* **9**(4): 1561-1564.
- Khaliq, A. and Cheema, Z. A. (2005). Influence of irrigation and nitrogen management on some agronomic traits and yield of hybrid sunflower. *International Journal of Agricultural Botany*, **7**: 915-919
- Marschner, H. 1995. Mineral Nutrition of Higher Plants. Academic Press, New York, NY.
- Meena, B.P., Kumawat, S.M. and Yadav, R. S. 2011. Effect of planting geometry and nitrogen management on groundnut (*Arachis hypogaea* L.) in loamy sand soil of Rajasthan. *Indian Journal of Agricultural Sciences* **81**(1): 86-88.
- Parameshwarareddy, R., Angadi, S.S. and Yenagi, B.S. 2019. Influence of plant spacing and fertilizer levels on growth and yield of summer groundnut. *Journal of Pharmacognosy and Phytochemistry* **8**(3): 3745-3748.
- Sree, S.M., Sudhakar, P., Mahesh, V.U., Prathima, T. and Krishna, G.T. 2020. Physiological responses of variable growth habit groundnut (*Arachis hypogaea* L.) genotypes at different planting densities. *International Journal of Current Microbiology and Applied Sciences* **9**(6): 3717-3725.
- Zhao, C., Shao, C., Yang, Z., Wang, Y., Zhang, X. and Wang, M. 2017. Effects of planting density on pod development and yield of peanuts under the pattern of precision planted peanuts. *Legume Research* **40**(5): 901-905.

Table 1. Nitrogen and spacing influence on growth parameters of groundnut.

Treatment	Plant height (cm) at harvest	Dry weight (g/plant) at harvest	Root nodules at harvest	CGR (g/m ² /day) 20-40 DAS	RGR (g/g/day) 20-40 DAS
T ₁	50.80	34.09	33.20	29.79	0.099
T ₂	51.83	33.63	32.93	31.44	0.104
T ₃	52.27	34.75	34.60	30.83	0.098
T ₄	49.10	37.09	38.17	28.93	0.103
T ₅	49.20	37.97	38.73	28.33	0.106
T ₆	50.03	38.00	39.57	28.77	0.101
T ₇	47.37	34.15	35.33	21.84	0.102
T ₈	47.03	34.35	35.57	22.62	0.110
T ₉	48.00	35.36	37.03	22.58	0.103
SEm (±)	0.32	0.45	0.35	0.97	0.001
CD (5%)	0.97	1.34	1.06	2.91	-

Table 2. Nitrogen and spacing effect on yield and yield parameters of groundnut.

Treatment	Number of pods/plants	Seed index (g)	Seed yield (kg/ha)	Haulm yield (kg/ha)	Shelling percentage (%)	Harvest Index (%)
T ₁	24.33	36.66	2050.3	2748.2	67.10	42.32
T ₂	25.36	37.26	2131.0	3011.1	67.00	41.25
T ₃	25.56	37.10	2164.0	3095.2	67.36	41.14
T ₄	30.56	37.63	2328.1	3373.6	68.93	40.82
T ₅	30.53	37.03	2353.6	3380.3	68.73	41.04
T ₆	30.80	37.50	2355.6	3402.3	68.11	40.91
T ₇	28.33	37.26	2087.3	3172.3	67.93	41.89
T ₈	28.60	37.93	2118.0	3225.4	67.66	41.17
T ₉	28.76	37.43	2121.2	3351.6	68.78	40.60
SEm (±)	0.23	0.58	20.03	13.81	0.48	0.23
CD (5%)	0.70	-	60.04	41.4	-	0.68

Figure 1: Graph relates about plant height




Figure 2: Graph relates about plant dry weight




Figure 3: Graph relates about number of root nodules




Figure 4: Graph relates about CGR




Figure 5: Graph relates about RGR

