

**INFLUENCE OF WEATHER ON GROUNDNUT GENOTYPES UNDER  
EXTENDED SOWING WINDOW DURING KHARIF SEASON**

**ABSTRACT**

A field experiment was conducted at Main Agricultural Research Station, University of Agricultural Sciences, Dharwad during the year 2018. The experiment was laid out in split plot design consist of four sowing window as main plots *i.e.*, June 1<sup>st</sup> fortnight, June 2<sup>nd</sup> fortnight, July 1<sup>st</sup> fortnight and July 2<sup>nd</sup> fortnight and four groundnut genotypes as sub plots *i.e.*, GPBD-4, G2-52, Dh-245 and JL-1085 with three replications. The groundnut crop sown during first fortnight of June recorded significantly higher pod yield (3,551 kg ha<sup>-1</sup>) and haulm yield (4,477 kg ha<sup>-1</sup>) compared to second fortnight of July and was at par with first fortnight of July and second fortnight of June. Among groundnut genotypes, G2-52 recorded significantly higher pod yield (3,364 kg ha<sup>-1</sup>) and haulm yield (4,239 kg ha<sup>-1</sup>) over JL-1085 and was on far with Dh-245 and GPBD-4. The groundnut genotype, G2-52 sown during first fortnight of June recorded higher pod yield (3,841 kg ha<sup>-1</sup>) and haulm yield (4,840 kg ha<sup>-1</sup>) compared to JL-1085 sown during second fortnight of July.

**Key words:** *Arachis hypogaea, Climate, Date of sowing, Genotypes, growth, ground nut, yield, Sowing window, varieties, Weather etc.*

**1. INTRODUCTION**

Oil seeds are most important crops next to food grains interms of production in Indian agriculture. Groundnut may remain as an essential oilseed crop for the semi-arid regions if the projected demand for oils and fat needs to be met with sustainability. Groundnut constitute the important element of Indian diet. Groundnut kernel contains edible oil (48-50 %), protein (25 %), carbohydrates (20 %) and it is rich source of vitamins and minerals.

Ground nut is cultivated across the world in all tropical and sub-tropical countries and is commercially cultivated between 40° N and 40° S latitude. Securing global food supply in an increasingly volatile climate and rapidly growing population is and will remain to be one of the greatest challenges facing humanity in the 21st century [1,2&3]. Globally, the crop is cultivated in an area of 29.92 million hectares with a total production of 50.17 million metric tonnes of nut shell (pod). The average productivity is 1,680 kg per hectare [4]. According to the all India kharif crop coverage report, Government of India, as on 17th September 2021, groundnut was sown in around 49.14 lakh hectares as compared to last year (50.97 lakh ha). Among the states, Gujarat stood first in area coverage with 19.09 lakh ha followed by Rajasthan (7.76 lakh ha), Andhra Pradesh (6.27 lakh ha), Karnataka (4.75 lakh ha) and Madhya Pradesh (3.82 lakh ha)[5].

Yield of crop depend upon many climate factors like solar radiation, temperature, moisture and precipitation can be very important. The oilseed crops, particularly peanut is very much sensitive to climatic factors like radiation and temperature [6]. Groundnut can be grown under wide range of climatic condition however, it is growing best in temperature range between 22 °C and 37 °C. Yield variation can be depending upon the variability of rainfall *i.e.*, distribution and amount of rainfall. The effectiveness of rainfall in production of crop as influenced by the commencement of sowing rains as well as distribution and amount of rainfall in the course of the season [7].

Groundnut production and productivity is being reduced day by day in the country [8], particularly in Karnataka and cannot possible to meet the requirement of growing population, which may be due to non-availability of good quality seed of improved genotypes, improper sowing time and establishment methods etc. In dryland agriculture, farmers have limited choice for sowing time, however in irrigated situation sowing time is one of the crucial non-economic input affecting the crop yield [8]. Being a tropical crop, groundnut requires warm climate and can be grown in *kharif* season due to congenial climate. Therefore, proper sowing time of groundnut in *kharif* is important for enhancing yield. The improved groundnut genotypes increase 25-28 per cent of yield, while 30-32 per cent by improved management practices [9].

Crop management practices such as time of sowing and duration of genotype may also influence growth, yield and quality parameters of groundnut. The real impact of seed maturity is dependent on genotype, climatic conditions and interaction of genotype between climate. Out of several above factors responsible for its low productivity, proper time of sowing under good rainfall situation with high yielding genotypes is considered as a prime constraint. Hence, the present investigation was carried out on **Analyzing the yield of groundnut genotypes in relationship with climate under extended sowing window during *kharif* season** with the objective to study the effect of sowing dates on growth and yield of groundnut genotypes.

## 2. MATERIAL AND METHODS

The experiment was laid out in split plot design consist of four date of sowing as main plots *i.e.*, D<sub>1</sub>: First fortnight of June (14-06-2018), D<sub>2</sub>: Second fortnight of June (25-06-2018), D<sub>3</sub>: First fortnight of July (06-07-2018) and D<sub>4</sub>: Second fortnight of July (19-07-2018) and four groundnut genotypes as sub plots *i.e.*, G<sub>1</sub>: GPBD-4, G<sub>2</sub>: G2-52, G<sub>3</sub>: Dh-245 and G<sub>4</sub>: JL-1085 with three replications and total 16 treatment combination. Simple averages, percentage, differences are used and the data is expressed in form of graphs for easy understanding. The experiment was conducted in medium to deep black soils *i.e.*, *Vertisols*. The soil type of the experimental site was clay loam. The pH of the soil was neutral with 7.62. The soil was low in available nitrogen content (262 kg N ha<sup>-1</sup>) and medium in available phosphorous (33 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) and high in available potassium (389 kg K<sub>2</sub>O ha<sup>-1</sup>).

The all management practices are followed according to the UAS Dharwad package of practice. Seeds were weighed separately for each plot at the rate of 125 kg ha<sup>-1</sup>. Groundnut seeds were treated with (Carboxin 37.5% + Thiram 37.5%) *i.e.*, Vitavax power followed by *Rhizobium* and phosphate solubilizing bacteria. By following 30 cm×10 cm spacing (Row to row: 30 cm and plant to plant: 10 cm) the lines were opened with the help of marker and the seeds were hand dibbled and covered with soil. Basal application of farm yard manure @ 7.5 t ha<sup>-1</sup>, nitrogen @ 18 kg N ha<sup>-1</sup>, phosphorus @ 46 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and potassium @ 25 kg K<sub>2</sub>O ha<sup>-1</sup> were applied in the form of diammonium phosphate and muriate of potash along with this FeSO<sub>4</sub> and ZnSO<sub>4</sub> each @ 25 kg ha<sup>-1</sup> and gypsum at 500 kg ha<sup>-1</sup> during peg initiation stage. According to the date of sowing and duration of the genotypes when crop attains maturity *i.e.*, the inside of the shell turns dark

and the kernels reach maximum growth accompanied by good colouration of the seed coat, the crop was harvested.

Pods from the plot area were cleaned to remove the soil adhering to the pods, impurities and immature pods. The developed pods were sun dried completely and weighed. Pod yield per hectare was calculated and expressed in kilogram per hectare. After separating the pods from harvested groundnut plants, the remaining produce was sun dried and haulm yield per hectare was calculated and expressed in kilogram per hectare.

### **3. RESULTS AND DISCUSSION**

#### **3.1. EFFECT OF WEATHER PARAMETERS ON PERFORMANCE OF *KHARIF* GROUNDNUT**


The average rainfall recorded during the months of June to October (1950 to 2017) was 555.08 mm. However, the rainfall received during the crop growth period *i.e.*, June to October, 2018 was as 462.8 mm. During the experimental year 2018, a rainfall of 892.2 mm was received which was 172.6 mm higher than the normal (1950-2017). Out of 892.2 mm rainfall, 462.8 mm was received during the crop growth period for all date of sowing (June to October, 2018) (Table 1 and Fig. 1). The groundnut crop requires 400-500 mm rainfall, whereas different sowing window observed different amount and distribution of rainfall. First, second, third and four sowing window recorded 302, 288, 344 and 255 mm rainfall, respectively (Table 2 and Fig. 2).

The crop sown during first fortnight of June recorded 302 mm rainfall in 34 rainy days, which was well distributed *i.e.*, 71 mm (10 days) between 0-30 days, 106.4 mm (15 days) between 30-60 days, 56.6 mm (5 days) between 60-90 days and 68.4 mm (4 days) between 90 to harvest. Almost similar rainfall distribution was experienced during first fortnight of July and second fortnight of June.

**Table 1: Monthly mean meteorological data during crop growth period of groundnut (2018) and the average of past 68 years (1950-2017) at the Main Agricultural Research Station, UAS, Dharwad**

Month	Rainfall (mm)		Rainy days	Temperature (°C)				Relative humidity (%)	
	1950-2017	2018		Maximum		Minimum		1950-2017	2018
				1950-2017	2018	1950-2017	2018		
January	0.77	0.0	0	28.84	29.8	14.66	13.9	52.96	66.3
February	10.84	1.0	0	31.66	32.1	16.11	15.7	44.88	46.3
March	3.65	72.4	3	34.89	34.9	18.64	19.3	42.42	46.6
April	38.43	32.8	3	36.52	36.2	20.16	21.1	58.61	58.2
May	71.92	250.4	10	35.19	34.7	21.23	21.3	64.28	66.7
June	107.59	115.4	9	29.60	28.8	21.12	21.0	79.69	80.5
July	134.47	128.8	19	27.21	25.8	20.71	20.6	86.25	87.5
August	103.98	72.2	7	26.85	25.9	20.41	20.2	86.09	87.2
September	103.29	69.6	4	28.40	29.6	20.02	18.9	81.81	75.7
October	105.75	76.8	6	29.72	31.6	19.40	18.3	75.20	65.5
November	31.49	34.4	2	29.71	30.5	16.46	16.3	69.67	54.8
December	7.45	38.4	1	28.83	28.8	14.04	14.6	67.40	60.6
Total/mean	719.6*	892.2*	64*	30.6	30.7	18.6	18.4	67.4	66.3

**Note : \* mark represents the sum of all values**



**Figure 1: Monthly meteorological data during crop growth period of groundnut (2018) and the average of past 68 years (1950-2017) at the Main Agricultural Research Station, UAS, Dharwad**

**Table 2: Meteorological data during crop growth stages of groundnut for all four sowing window during *kharif* 2018 at the Main Agricultural Research Station, UAS, Dharwad**

Sowing window	Growth stage	Max. Temp. (°C)	Min. Temp. (°C)	RH Max. (%)	RH Min. (%)	Mean RH (%)	Rainfall (mm)	Rainy days
<b>D<sub>1</sub>: June 1<sup>st</sup> Fortnight</b>	<b>0-30</b>	27.42	20.60	88.20	77.33	82.77	71	10
	<b>31-60</b>	25.67	20.64	90.03	86.13	88.08	106.4	15
	<b>61-90</b>	26.52	19.58	89.03	79.23	84.13	56.6	5
	<b>91-Harvest</b>	30.57	19.10	81.40	65.45	73.43	68.4	4
	<b>Mean /Total</b>	<b>27.55</b>	<b>19.98</b>	<b>87.17</b>	<b>77.04</b>	<b>82.10</b>	<b>302.4*</b>	<b>34*</b>
<b>D<sub>2</sub>: June 2<sup>nd</sup> Fortnight</b>	<b>0-30</b>	26.21	20.69	89.30	83.30	86.30	118.8	18
	<b>31-60</b>	25.70	20.41	90.37	85.60	87.98	78.6	9
	<b>61-90</b>	28.53	19.03	85.23	70.07	77.65	39.4	2
	<b>91-Harvest</b>	31.66	19.61	80.35	63.15	71.75	51	5
	<b>Mean /Total</b>	<b>28.03</b>	<b>19.94</b>	<b>86.31</b>	<b>75.53</b>	<b>80.92</b>	<b>287.8*</b>	<b>34*</b>
<b>D<sub>3</sub>: July 1<sup>st</sup> Fortnight</b>	<b>0-30</b>	25.61	20.65	90.07	85.93	88.00	129.2	18
	<b>31-60</b>	25.90	20.12	89.87	84.80	87.33	75.2	8
	<b>61-90</b>	30.11	18.95	83.03	66.77	74.90	76.2	6
	<b>91-Harvest</b>	31.68	19.27	78.95	57.85	68.40	63	3
	<b>Mean /Total</b>	<b>28.33</b>	<b>19.75</b>	<b>85.48</b>	<b>73.84</b>	<b>79.66</b>	<b>343.6*</b>	<b>35*</b>
<b>D<sub>4</sub>: July 2<sup>nd</sup> Fortnight</b>	<b>0-30</b>	25.61	20.53	90.50	86.17	88.33	91.4	13
	<b>31-60</b>	27.63	19.20	86.80	73.20	80.00	25.8	2
	<b>61-90</b>	31.50	19.65	81.53	61.43	71.48	75.2	6
	<b>91-Harvest</b>	30.70	17.01	66.15	56.50	61.33	62.6	3
	<b>Mean /Total</b>	<b>28.86</b>	<b>19.01</b>	<b>81.26</b>	<b>69.33</b>	<b>75.29</b>	<b>255*</b>	<b>24*</b>

**Note : \* mark represents the sum of all values**




Figure 2: Meteorological data during crop growth stage of groundnut for all four sowing window during *kharif* 2018 at the Main Agricultural Research Station, UAS, Dharwad



Differences in the pod yield are mainly influenced by the variation in the distribution of rainfall. The higher pod yield was obtained when the crop was sown during first fortnight of June as the crop experienced good distribution of rainfall throughout the cropping period. The crop sown during first fortnight of July and second fortnight June received more rainfall at early stage of crop then, optimum distribution during rest of the crop period. Whereas, the crop sown during second fortnight of July experienced limited water supply resulting in moisture stress periods during the crop growth and development stages which has resulted in significantly lower pod yield. This sowing window received only 255 mm rainfall in 24 rainy days. Moisture stress between 30-60 days of the crop growth which affected the peak flowering, peg initiation and pod development.

The highest and the lowest mean maximum temperature were 36.2 °C (April) and 25.8 °C (July), respectively while the respective highest and the lowest minimum temperature were 21.3 °C (May) and 13.9 °C (January), respectively. During crop growth period the maximum and minimum temperature was observed equally in all sowing windows (Max. temp. 28.19 °C  $\pm$  0.40 and Min. temp. 19.67 °C  $\pm$  0.32). During the experimental year, the mean relative humidity ranged from 46.3 per cent in February to 87.5 per cent in July. The mean relative humidity decreased from first date of sowing (77.04 %) to fourth date of sowing (69.33 %). Whereas, 73.94  $\pm$  2.35 per cent deviation from the mean relative humidity was observed for all date of sowing (Table 1 and Fig. 2).

### **3.2. ACCUMULATED GROWING DEGREE DAYS AND ASSESSING THE MATURITY OF THE GROUNDNUT CROP**

Growth and development of groundnut are affected by different uncontrollable environmental conditions such as effect of different planting dates, growing degree days and agronomic traits of genotypes. The currently available methods used to predict groundnut maturity are based on hull colour determination and are somewhat labour-intensive and subject to the observer's ability to finely discriminate colour classes. In addition, an over-mature crop may lead to increased mechanical losses during the process of digging due to deteriorating peg strength with age. This mechanical loss, termed digging loss, is the retention of pods in the soil during the digging process due to separation of the pod from the stem attachment at the plant. Typical digging losses have been estimated to be 8% of the total yield but can reach 40 % at dates beyond optimal maturity [10&11]. Therefore, correctly assessing groundnut maturity prior





to digging is essential to the economic viability of groundnut production. Further, Ketring and Wheless (1989) found accumulated degree days to range between 1450 and 1670 at final harvest in case of groundnut [12]. In the present study, among the dates of sowing, growing degree days were found to be similar (Table 3). The range is from 1448 to 1485, indicating temperature condition prevailed during different sowing windows was uniform from sowing to maturity. Among, different genotypes under the study, higher growing degree days was observed with genotypes G2-52, GPBD-4 and Dh-245 as compared to JL-1085 indicating JL-1085 consumed less heat units to reach maturity. It may be attributed to the genotype JL-1085 being of relatively of shorter duration and had optimum maturity duration of 100 to 105 days as compared to other genotypes which are having optimum maturity of 105 to 110 days. These results show that overall groundnut plant development appears to be highly correlated with growing degree days of 1448 to 1485 under semi arid climate like Dharwad to determine the maturity of the crop. So in absence of the previous studies on this aspect in the study area, accumulated degree days between 1448 to 1485 could be used to judge the maturity of the groundnut crop.

### **3.3. SOIL MOISTURE CONTENT IN RELATION TO SOWING WINDOWS**

Soil moisture content from 0-30 cm depth at 10 days interval after sowing to harvest was recorded and presented in fig. 3. During crop period of first date of sowing (14.06.2018) highest soil moisture was observed in 24.07.2018, which coincides with the flowering to peg initiation stage (30-40 DAS) crop. Whereas, lowest moisture was recorded in 80-90 DAS *i.e.*, kernel development to maturity stage at 12.09.2018. In second date of sowing (25.06.2018) highest soil moisture was observed in 24.07.2018 which coincides with the emergence to flowering stage (20-30 DAS) crop. Whereas, lowest moisture was recorded in 70-80 DAS *i.e.*, pod formation to kernel development stage at 12.09.2018. During crop period of third date of sowing (06.07.2018) highest soil moisture was observed in 24.07.2018 which coincides with the germination to emergence stage (10-20 DAS) crop. Whereas, lowest moisture was recorded in 60-70 DAS *i.e.*, pod initiation to pod formation stage at 12.09.2018. In fourth date of sowing (19.07.2018) highest soil moisture was observed in 24.07.2018 which coincides with the sowing to germination stage (0-10 DAS) crop. Whereas, lowest moisture was recorded in 50-60 DAS *i.e.*, pegging stage to pod initiation stage at 12.09.2018 (Figure 3).

**Table: 3 Accumulated growing degree days under different date of sowing during *kharif*-2018**

<b>Accumulated GDD</b>					
<b>Date of sowing Genotypes</b>	<b>14<sup>th</sup> June</b>	<b>25<sup>th</sup> June</b>	<b>6<sup>th</sup> July</b>	<b>19<sup>th</sup> July</b>	<b>Mean</b>
<b>GPBD-4</b>	1503	1521	1530	1539	<b>1523</b>
<b>G2-52</b>	1503	1521	1530	1539	<b>1523</b>
<b>Dh-245</b>	1472	1490	1500	1509	<b>1493</b>
<b>JL-1085</b>	1427	1442	1458	1467	<b>1448</b>
<b>Mean</b>	<b>1476</b>	<b>1494</b>	<b>1504</b>	<b>1514</b>	



**Figure 3: Graphical representation of soil moisture content from 0 to 30 cm depth at 10 days interval after sowing to harvest under different dates of sowing during *kharif*.**

### **3.4. EFFECT OF SOWING WINDOW ON POD AND HAULM YIELD *KHARIF* GROUNDNUT**

Among the sowing windows, the crop sown during first fortnight of June recorded significantly higher pod yield ( $3,551 \text{ kg ha}^{-1}$ ) and haulm yield ( $4,477 \text{ kg ha}^{-1}$ ) and other yield attributes (Table 4) and was at par with the crop sown during first fortnight of July ( $3,372$  and  $4,222 \text{ kg ha}^{-1}$ , respectively) and second fortnight of June ( $3,163$  and  $4,056 \text{ kg ha}^{-1}$ , respectively). This might be due to better performance of yield attributes *viz.*, number of pods and pod weight. Similarly, higher leaf area at 60 DAS and at harvest and higher leaf area index, which resulted in higher leaf surface area for assimilation of photosynthates. The crop sown early during first fortnight June might have been exposed to better climatic condition resulting in higher photosynthetic rate and consequently increased leaf area. Higher dry matter production per plant at 60 DAS and at harvest was observed with early sown crop. Similar findings were observed by Guled (2013); Patel and Vaishnav (2003) in Gujarat and again proved by Sampath Kumar *et al.*, in Kadri, Andhra Pradesh (2020). [13,14&19].

The pod yield decreased by 29.30 per cent ( $2,509 \text{ kg ha}^{-1}$ ) with delayed sowing *i.e.*, second fortnight of July over the first fortnight of June (Table 4). It might be due to insufficient availability soil moisture at reproductive stages, like peak flowering, peg initiation and pod filling wherein, these are the critical stages of crop for moisture leading to stress reduced dry matter production and there by the translocation of photosynthates from leaves (source) to fruiting parts (sink). Hence there was lower number of pods and pod weight. Effect of rainfall had greater influence on vegetative growth of the crop under delayed sowing. Which ultimately reduced dry matter production, leaf area and leaf area index in second fortnight of July sowing window. Similar kind of results was observed by Sahu *et al.* (2004) and Thiyaagrajan *et al.* (2010) at Saurashtra region, Gujarat [7&15].

### **3.5. PERFORMANCE OF GROUNDNUT GENOTYPES UNDER DIFFERENT SOWING WINDOW**

It is necessary to grow the groundnut genotypes which can withstand weather aberrations by adapting to varied sowing window. Nageswar Rao (1992) revealed that, improved genotypes

contribute 25-28 per cent to the yield increase while improved management practices contribute 30-32 per cent [8]. Hence, it is an important factor to assess the performance of groundnut genotypes under varied sowing windows.

Among the different genotypes of groundnut, the genotype G2-52 recorded significantly superior pod yield (3,364 kg ha<sup>-1</sup>) and haulm yield (4,239 kg ha<sup>-1</sup>) over JL-1085 (2,868 kg ha<sup>-1</sup> & 3,639 kg ha<sup>-1</sup>, respectively) and was statistically on par with Dh-245 (3,209 kg ha<sup>-1</sup> & 4,044 kg ha<sup>-1</sup>, respectively) and GPBD-4 (3,154 kg ha<sup>-1</sup> & 3,975 kg ha<sup>-1</sup>, respectively). Similar findings were observed by Bharatha Lakshmi and Sambasiva Reddy, (2001) at Tirupati, Andhra Pradesh [12].

### **3.6. INTERACTION EFFECT OF SOWING WINDOW AND GENOTYPES ON YIELD OF *KHARIF* GROUNDNUT**

The interaction effect was found not significant between sowing window and genotypes for growth and yield parameters of groundnut. The groundnut genotype, G2-52 sown during first fortnight of June recorded higher pod yield (3,841 kg ha<sup>-1</sup>) and haulm yield (4,840 kg ha<sup>-1</sup>) compared to JL-1085 sown during second fortnight of July. This was also due to combination of genetic characters and also the optimum weather parameters during crop period.

Lower pod yield was recorded with second fortnight July with JL-1085 genotype (2,267 kg ha<sup>-1</sup>). Delay in sowing resulted in reduced pod yield (2868 kg ha<sup>-1</sup>) and haulm yield (3639 kg ha<sup>-1</sup>). This might be due to stress condition prevailed to the crop with delayed sowing due to lack of rainfall (255 mm) and rainy days (24). These results are in accordance with Kanade *et al.* (2016) at Pune and Mohite *et al.* (2017) at Kolhapur [13&14].

## **4. CONCLUSION**

Sowing of groundnut with normal onset monsoon *i.e.*, during first fortnight of June was highly productive (3,551 kg ha<sup>-1</sup>) and remunerative. But under delayed onset of monsoon, sowing could be extended up to second fortnight of July. The groundnut genotype, G2-52 was

superior and was on par with Dh-245 and GPBD-4. The groundnut genotype, G2-52 sown during first fortnight of June resulted in higher pod yield and haulm yield.

**Table 4: Effect of different sowing window on pod yield and kernel yield of groundnut genotypes**

	Pod yield (kg ha <sup>-1</sup> )					Haulm yield (kg ha <sup>-1</sup> )				
<b>Sowing window Genotype</b>	<b>14<sup>th</sup> June</b>	<b>25<sup>th</sup> June</b>	<b>6<sup>th</sup> July</b>	<b>19<sup>th</sup> July</b>	<b>Mean</b>	<b>14<sup>th</sup> June</b>	<b>25<sup>th</sup> June</b>	<b>6<sup>th</sup> July</b>	<b>19<sup>th</sup> July</b>	<b>Mean</b>
<b>GPBD-4</b>	3,517	3,215	3,421	2,464	<b>3,154</b>	4,432	4,115	4,276	3,079	<b>3,975</b>
<b>G2-52</b>	3,841	3,324	3,558	2,731	<b>3,364</b>	4,840	4,255	4,447	3,414	<b>4,239</b>
<b>Dh-245</b>	3,556	3,217	3,488	2,576	<b>3,209</b>	4,480	4,118	4,360	3,220	<b>4,044</b>
<b>JL-1085</b>	3,288	2,898	3,019	2,267	<b>2,868</b>	4,156	3,738	3,805	2,856	<b>3,639</b>
<b>Mean</b>	<b>3,551</b>	<b>3,163</b>	<b>3,372</b>	<b>2,509</b>		<b>4,477</b>	<b>4,056</b>	<b>4,222</b>	<b>3,142</b>	
<b>Source of variation</b>	<b>S.Em. ±</b>		<b>CD (P=0.05)</b>			<b>S.Em. ±</b>		<b>CD (P=0.05)</b>		
<b>Sowing window (D)</b>	122		422			153		530		
<b>Genotype (G)</b>	105		307			134		390		
<b>Two genotype means at same sowing window</b>	211		NS			267		NS		
<b>Two sowing window means at same or diff. genotype</b>	219		NS			277		NS		

NS: Non significant



## 5. REFERENCE

1. Fahad, S., Saud, S., Akhter, A., Bajwa, A.A., Hassan, S., Battaglia, M., 2021, Bio-based integrated pest management in rice: an agro-ecosystems friendly approach for agricultural sustainability. *J. Saudi Soc. Agric. Sci.*, 20: 94-102.
2. Fujimori, S., Hasegawa, T., Krey, V., Riahi, K., Bertram, C., Bodirsky, B.L., 2019, A multi-model assessment of food security implications of climate change mitigation. *Nat. Sustain.*, 2: 386-396
3. Harrison, M.T., Cullen, B.R., Mayberry, D.E., Cowie, A.L., Bilotto, F., Badgery, W.B., 2021, Carbon myopia: the urgent need for integrated social, economic and environmental action in the livestock sector. *Glob. Chang. Biol.*, 01:1-36
4. Anonymous. 2021. Peanut area, yield and production. United States Foreign Agricultural Services, editor. USDA
5. Anonymous, 2021, Groundnut Outlook - November 2021, Agricultural Market Intelligence Centre, PJTSAU. [www.agricoop.gov.in](http://www.agricoop.gov.in), pp. 1-3.
6. Banik, N. C., Nath, R. and Chakraborty, P. K., 2009, Effect of dates of sowing on growth and yield of groundnut crop. *J. Crop. Weed.*, 5(2): 59-62.
7. Sahu, D. D., Patra, B. K. and Patoliya, B. M., 2004, Effect of sowing time and rainfall distribution on groundnut yield. *IAN*, 24: 39-42
8. Sardana, V. and Kandhol, S. S., 2007, Productivity of semi-spreading and bunch type varieties of groundnut as influenced by sowing dates. *An Op. Acc. J.*, 5: 1-3.
9. Nageswara Rao., 1992, Some crop physiological approaches for groundnut improvement. *J. Oilseeds Res.*, 9 (2): 281-296.
10. Young, J. H., N. K. Person, J. O. Donald, and W. D. Mayfield., 1982, Harvesting, curing and energy utilization, pp. 458-485. In H. E. Pattee and C. T. Young (eds.). *Peanut Sci. Tech. Amer. Peanut Res. Educ. Society.*, Inc., Yoakum, TX, USA.
11. Lamb, M. C., Masters, M. H., Rowland, D., Sorensen, R. B., Zhu, H., Blankenship, P. D. and C. L. Butts., 2004, Impact of sprinkler irrigation amount and rotation on peanut yield. *Peanut Sci.*, 31: 108-113.
12. Ketring, D. L., and Wheless, T. G., 1989, Thermal time requirements for phenological development of peanut. *Agron. J.*, 81: 910-917.

13. Guled, P. M., 2013, Crop weather relationship studies in *kharif* groundnut (*Arachis hypogaea* L.) under middle Gujarat agroclimatic condition. *Ph. D. Thesis*, Anand Agricultural University. Gujarat.
14. Patel, J. S. and Vaishnav, P. R., 2003, Evaluation of different approaches to study the effect of rainfall on groundnut in dry farming area of Gujarat. *J. Agrometeorol.*, 5 (1): 76-83.
15. Thiyaagrajan, G., Ranghaswami, M. V., Rajakumar, D. and Kumaraperumal, R., 2010, Deficit irrigation effects on groundnut (*Arachis hypogaea* L.) with micro sprinklers. *Madras Agric. J.*, 97 (13): 40-42.
16. Bharatha Lakshmi, M. and Sambasiva Reddy, A., 2001, Effect of crop geometry on growth and yield of rainfed Spanish bunch groundnut varieties. *J. Oilseeds Res.*, 18(1): 134-135.
17. Kanade, S. G., Shaikh, A. A. and Jadhav, J. D., 2016, Effect of sowing dates in groundnut (*Arachis hypogaea* L.) on growth, yield attributing characters and yield. *Adv. Res. J. Crop Improv.*, 6 (1): 5-11.
18. Mohite, U. A., Mohite, A. B. and Jadhav, Y. R., 2017, Effect of sowing windows on growth and yield of groundnut varieties during *kharif* season. *Contemporary Res. India.*, 7 (2): 2231-2137.
19. Sampath Kumar D., Anitha D., Prathyusha D., 2020, Effect of dates of sowings on growth and productivity of different cultivars of groundnut (*Arachis hypogaea* L.). *J. Pharmacognosy and Phytochemistry*. 2020; 9(5):2211-2213.