

Effect of Bio Fertilizers on the Vegetative growth, yield and Economic Analysis of Strawberry (*Fragaria x ananassa* Duch.) Cv. Winter dawn in Vertical Farming System

ABSTRACT

A field experiment was carried out during 2021-2022 to see the Effect of Bio Fertilizers on the Vegetative growth, and Economic Analysis of Strawberry (*Fragaria x ananassa* Duch.) Cv. Winter dawn in Vertical Farming System with 10 treatments including control in combinations of organic and microbial sources of nutrients (Vermicompost, FYM, *Azotobacter*, *Azospirillum* and PSB) replicated thrice with 3 plants per replication in Randomized Block Design. Observations were recorded for vegetative growth, fruit yield, quality. In different combinations (biofertilizers and organic manure) the treatment T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) recorded highest plant height, plant spread, number of leaves and leaf area as compared to T₇ (Soil (50%) + FYM (50%) + PSB (2g) + *Azotobacter* (2g)). Plant treatment T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) registered earliest flowering and also highest number of flowers per plant. The maximum fruit weight, number of fruits per plant and yield were recorded with plants treated with T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) followed by T₇ (Soil (50%) + FYM (50%) + PSB (2g) + *Azotobacter* (2g)) at 90 DAP followed by T₈ (Soil (50%) + coir pith (50%) + *Azospirillum* (2g) + PSB (2g)) treatment. The maximum Benefit: Cost ratio (1: 3.39) was recorded in the T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) than T₇ (Soil (50%) + FYM (50%) + PSB (2g) + *Azotobacter* (2g)) due to its lower cost of production. The highest yield and best quality fruit were recorded in the combination of T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)).

Key words: strawberry, vertical farming, organic manure, biofertilizers.

INTRODUCTION

Strawberry (*Fragaria × ananassa* Duch.) belongs to the family Rosaceae. The cultivated strawberry (*Fragaria × ananassa* Duch.) was originated from the hybridization of two American species viz., *Fragaria chilioensis* Duch. and *Fragaria virginiana* Duch. All the cultivated varieties of strawberry are octaploid ($2n = 8x = 56$) in nature. It is herbaceous crop with prostrate growth habit, which behaves as an annual in sub-tropical region and perennial in temperate region. (Gallette and Bringhurst, 1990).

Strawberry is used as fresh fruit being rich in vitamin C (30-120 mg/100g fruit pulp) and ellagic acid, which has anti cancerous property. Fruits are attractive with distinct pleasant aroma and flavour, consumed as dessert and also have a special demand by the fruit processing units for the preparation of jams, ice cream, syrups etc. Vermicompost contains plant growth regulating materials, such as humic acids and plant growth regulators like auxins, gibberellins and cytokinin's; (Grappelli *et al.*, 1987), which are responsible for increased plant growth and yield of strawberry fruit crops. Biofertilizers are one of the best modern tools for agriculture and are used to improve the fertility and quality of the soil. It offers an economically attractive and ecologically sound route for augmenting nutrient supply that enables to plant growth and development of strawberry.

Keeping these facts in view, the present investigation was initiated to study. Low-enter agricultural gadget which is based at the input of natural substances preserve terrific promise now no longer simplest to limit the usage of artificial fertilizer, however additionally to enhance crop productiveness and to make sure surroundings sustainability towards nutrient mining and degradation of soil and water resources (Tilman *et al.*, 2002; Kravchenko *et al.*,

2017).

Among the natural amendment, farmyard manure (FYM) has been extensively utilized in agricultural fields and the composted shape of this manure is desired over clean manure to dispose of the hazard of N loss thru leaching and floor runoff, boom soil natural matter, suppress soil-borne pathogens and to mitigate greenhouse gas emissions (Darby *et al.*, 2004) Vermicompost is organic material that has been broken down in a mesophilic process by interactions between microorganisms and earthworms to produce fully stabilized organic soil amendments.

Vermicompost contains significant amounts of nutrients, a large valuable microbial population, and biologically active metabolites (gibberellins, cytokinins, auxins), as well as vitamins B, which can be used alone or in combination with other inorganic or organic fertilizers to improve crop in soils is influenced by soil physicochemical properties (organic matter, pH, temperature, soil depth, soil moisture) and microbiological properties (microbial interactions).

is a well-studied plant growth promoting bacteria due to its ability to colonise the roots of various plant species, the majority of which are agriculturally important (Cessán *et al.*, 2008).

Vertical farming is crop husbandry in which crops are planted in vertically managed layers in order to harness the untapped vertical area that is otherwise overlooked in almost all cultivation practises. Furthermore, approximately 80% of total arable land is currently underutilized globally. (Ellingsen & Despommier, 2008) Some urban planners and agricultural leaders have argued that cities must produce food internally in order to manage the demand-supply ratio and avoid falling food prices, harmful pollution, and inflation (Kodmany, 2018). Growing media/substrates are the materials in which the plant will grow and play an important role in productivity. They provide anchorage for plant roots, air spaces for respiration, and the ability to retain enough available water for plant growth. For growing plants, a variety of growth substrates are available. For growing plants, the growing medias are used solely or in combinations (Arancon N.Q and Edwards C.A, 2005). The types of media combinations vary depending on the plant species, as well as the grower and stage of growth, such as nursery raising or transplanting. Nowadays, various types of growing media are available on the market, such as coco-peat, coir, vermicompost, perlite, bagasse, sawdust, and so on, but some media performed well despite being insignificant in yield. (Rawahya *et al.*, 2009).

MATERIALS AND METHODS

The area of Prayagraj district comes under subtropical belt in the South East of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46°C – 48°C and seldom falls as low as 4°C – 5°C. The relative humidity ranged between 20–94 percent. The average rainfall in this area is around 1013.4 mm annually. However, occasional precipitation is also not uncommon during winter months. The experiment was conducted in the vertical system at the experimental site of Department of Horticulture, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Prayagraj during 2021–2022. Treatments namely T₀ CONTROL (sand, soil, coir pith), T₁ Sand (25%) + Soil (50%) + Vermicompost (25%) + *Azotobacter* (2g), T₂ Sand (25%) + Soil (50%) + Vermicompost (25%) + PSB (2g), T₃ Sand (25%) + Soil (50%) + Vermicompost (25%) + *Azospirillum* (2g), T₄ Soil (50%) + coir pith (25%) + FYM (25%) + *Azotobacter* (2g), T₅ Soil (50%) + coir pith (25%) + FYM (25%) + PSB (2g), T₆ Soil (50%) + coir pith (25%) + FYM (25%) + *Azospirillum* (2g), T₇ Soil (50%) + FYM (50%) + PSB (2g)

+ *Azotobacter* (2g), T₈ Soil (50%) + coir pith (50%) + *Azospirillum* (2g) + PSB (2g), T₉ Soil (50%) + Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g) were tested in randomized block design with three replications. The observations were recorded on three randomly selected plants from each treatment. Vegetative characters [Plant height (cm), Plant spread (cm), Number of leaves per plant, Leaf Area Index (cm²)], yield and economic attributes [Number of fruits per plant (90 days), Yield plant⁻¹(g) (90 days), Fruit diameter (cm) (90 days), Fruit length (cm) (90 days), TSS (°Brix) (90 days), Acidity (%) (90 days), Ascorbic Acid (mg/100g) (90 days), Net return (₹ /100²), Benefit: Cost ratio].

RESULTS AND DISCUSSION

VEGETATIVE CHARACTERS

It is clear from the (Table 1) & (Graph 1) that the maximum plant height (8.56 cm, 15.79 cm, 22.89 cm) was recorded in T₉ (Soil (50%) + Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)). The minimum plant height per plant (5 cm, 9.9 cm, 16.67 cm) was recorded in T₀ (CONTROL (sand, soil, coir pith)). Vermicompost is regarded as a more abundant source of plant nutrients (Arancon *et al.*, 2003 and 2004). *Azotobacter* is the most extensively studied free living nitrogen fixing bacteria, and it is known to synthesize biologically active growth promoting substances in addition to fixing atmospheric nitrogen (Yadav *et al.*, 2010) and are further influenced by plant vegetative growth. PSB stimulated plant growth by promoting phosphate dissolution (Nowsheen *et al.*, 2006) as well as the biosynthesis of auxin (Sattar and Gaur, 1987) and IAA (Sattar and Gaur, 1987).

The maximum plant spread (28 cm) was recorded in T₉ (Soil (50%) + Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)). The minimum plant spread (23.07 cm) was recorded in T₀ (CONTROL (sand, soil, coir pith)). Vermicompost is regarded as a more abundant source of plant nutrients, growth regulators, enzymes, antifungal and antibacterial compounds (Arancon *et al.*, 2003 and 2004). PSB stimulated plant growth via phosphate dissolution (Nowsheen *et al.*, 2006) and the biosynthesis of auxin (Sattar and Gaur, 1987) and IAA (Sattar and Gaur, 1987). Vermicompost has previously been shown to have a positive effect on plant growth in strawberries (Arancon *et al.*, 2003 and 2004, Singh *et al.*, 2008, Nazir *et al.*, 2006, Yadav *et al.*, 2010).

The maximum number of leaves (27.56) was recorded in T₉ (Soil (50%) + Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) and the minimum number of leaves per plant (20.11) was recorded in T₀ (CONTROL (Sand, soil, coir pith)). The memoir fertilisers had a significant impact on the number of strawberry leaves per plant *Azotobacter* is the most hugely studied tone- ruling living nitrogen setting up bacteria, and in addition to its goods to dispose atmospheric nitrogen, it's grasped to synthesise biologically alive excrescence advancing quiddities likewise as IAA, GA, and others (Yadav *et al.*, 2010). The developments acquired are in attestation with the judgments of Yadav *et al.*, (2010) who reported that conjugated exercise of life conditions, vermicompost with inorganic conditions significantly compounded the number of leaves and flake demesne of strawberry and Nowsheen *et al.*, (2006) plant that the exercise of P.S.B.


The maximum leaf area (38.8 cm) was recorded in T₉ (Soil (50%) + Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)). The minimum leaf area (24.41 cm²) was recorded in T₀ (CONTROL (sand, soil, coir pith)). The mechanisms by which PSB stoked plant growth is through phosphate dissolution (Nowsheen *et al.*, 2006) and in the biosynthesis

of auxin (Sattar and Gaur, 1987) and IAA (Bareae *et al.*, 1976). Positive effect of vermicompost on plant growth has also been recorded by (Nazir *et al.*, 2006, Yadav *et al.*, 2010). Combined operation of memoir diseases, vermicompost with inorganic fertilisers significantly increased the number of leaves and splint area of strawberry (Aroncon *et al.*, 2003 & 2004, Singh *et al.*, 2008)

Table 1: Effect of biofertilizers on vegetative growth characters of strawberry under vertical system

Treatment	Treatment combination	Plant height (cm) (90 days)	Plant spread (cm) (90 days)	Number of leaves (90 days)	Leaf area (cm ²) (90 days)
T ₀	CONTROL (sand, soil, coir pith)	16.67	23.07	20.11	24.41
T ₁	Sand (25%) + Soil (50%) + VC (25%) + <i>Azoto.</i> (2g)	18.11	24.97	23.44	28.9
T ₂	Sand (25%) + Soil (50%) + VC (25%) + PSB (2g)	17	23.77	20.56	26.17
T ₃	Sand (25%) + Soil (50%) + VC (25%) + <i>Azosp.</i> (2g)	17.56	24.37	22.22	28
T ₄	Soil (50%) + coir pith (25%) + FYM (25%) + <i>Azoto.</i> (2g)	20.89	26.57	22.89	33.67
T ₅	Soil (50%) + coir pith (25%) + FYM (25%) + PSB (2g)	19.22	25.77	24	29.8
T ₆	Soil (50%) + coir pith (25%) + FYM (25%) + <i>Azosp.</i> (2g)	20.11	26.23	25.33	32.17
T ₇	Soil (50%) + FYM (50%) + PSB (2g) + <i>Azoto.</i> (2g)	22	27.33	26.89	36.5
T ₈	Soil (50%) + coir pith (50%) + <i>Azosp.</i> (2g) + PSB (2g)	21.56	26.9	26.22	34.41
T ₉	Soil (50%) + VC (50%) + <i>Azoto.</i> (2g) + <i>Azosp.</i> (2g)	22.89	28	27.56	38.8
F-test		S	NS	S	S
SE. d (+)		1.01	2.06	1.87	2.68
CD (5%)		2.12	4.33	3.93	5.7

VC: vermicompost, Azoto: *Azotobacter*, Azosp: *Azospirillum*, PSB: Phosphate solubilizing bacteria



Graph 1: Effect of biofertilizers on vegetative growth characters of strawberry

YIELD AND ECONOMIC ATTRIBUTES

Numerically in (Table 2) & graphically in (Graph 2) the maximum number of fruits plant⁻¹ (21) was recorded in T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) and the minimum number of fruits per plant (11.89) was recorded in T₀ (CONTROL (Sand, soil, coir pith)). It is clear from the table that there are significant differences among the treatments at 90 days after planting number of fruits plant⁻¹ may be due to increase in the number of leaves which worked as an effective photosynthesis structure and produced high quantum of carbohydrates in the plant system. Further number of flowers, which redounded advanced fruits per plant, under present study due to capability of vermicompost in producing growth hormone, enzymes, antifungal and antibacterial composites, which in turns enhanced marketable yield over other treatments. Analogous findings also reported by Wang (1996), Yadav *et al.*, (2010), Umar *et al.*, (2009), Singh *et al.*, (2008), Rana and Chandel (2003), Zargar *et al.*, (2008), Umar *et al.*, (2010) Dadashpour and Jouki (2012) and Verma and Rao (2013) in strawberry.

The maximum fruits yield plant⁻¹ (638.40 g) was recorded in T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) and the minimum number of fruits yield per plant (216.08 g) was recorded in T₀ (CONTROL (sand, soil, coir pith)). Increase in fruit yield may be due to increase in the number of leaves which worked as an effective photosynthesis structure and produced high quantum of carbohydrates in the plant system. Vermicompost contains plant growth regulating accoutrements, similar as humic acids and plant growth controllers like auxins, gibberellins and cytokinin (Grappelli *et al.*, 1987), Analogous findings also reported by Wang (1996), Yadav *et al.*, (2010), Umar *et al.*, (2009), Singh *et al.*, (2008), Rana and Chandel (2003), Zargar *et al.*, (2008), Umar *et al.*, (2010) Dadashpour and Jouki (2012) and Verma and Rao (2013) in strawberry.

The maximum fruit diameter (3.36 cm) was recorded in T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)). The minimum fruit diameter (2.47 cm) was observed in T₀ (CONTROL (sand, soil, coir pith)). Increase in fruit diameter

may be due to balanced clearness of macro and micronutrients and growth promoting hormones produced by different bio-fertilizers applied in different treatment combinations. This could be linked to superior fruit fillers as a result of more balanced nutrient intake, which could have led to improved metabolic conditioning in the plant, leading to a high protein and carbohydrate admixture (Singh *et al.*, 1970). Also, the different partitioning of photosynthesis towards the cesspool by *Azotobacter* inoculation increased the fruit size and fruit diameter (Rana and Chandel, 2003).

The maximum fruit length (4.46 cm) was recorded in T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) & the minimum fruit length (3.1 cm) was observed in T₀ (CONTROL (sand, soil, coir pith)). Fruit length growth can be because of balanced availability of macro and micronutrients and boom selling hormones produced with the aid of using unique bio-fertilizers carried out in unique remedy combinations. This can be attributed to higher fillings of culmination because of extra balanced uptake of vitamins which may also have caused higher metabolic sports withinside the plant in the long run main to excessive protein and carbohydrate synthesis (Singh *et al.*, 1970). Also, the unique partitioning of photosynthesis toward the sink with the aid of using *Azotobacter* inoculation multiplied the fruit duration (Rana and Chandel, 2003 and Tripathi *et al.*, 2010). Suthar (1993) confirmed that amongst diverse natural sources, vermicompost had the maximum critical position accompanied with the aid of using FYM.

The maximum (TSS 9.57°B) was recorded in T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) and the minimum (TSS 5.31°B) was observed in T₀ (CONTROL (sand, soil, coir pith)). Increased TSS and general sugars in berry with the utility of *Azotobacter* and vermicompost can be attributed because of the fast metabolic transformation of starch and pectin into soluble compounds and speedy translocation of sugars from leaves to the growing fruits. These findings trust the consequences of Singh and Singh, (2006); Haynes and Goh (1987); Singh *et al.*, (2008) in strawberry; Rathi and Bist (2004) in pear and Baksh *et al.*, (2008) in guava. Badr *et al.*, (2003) pronounced that sugar is produced through breaking the starch withinside the leaves and Anthocyanin will even increase with growing glucose ranges because of mobileular metabolism, ensuing withinside the accumulation of soluble sugars withinside the mobileular.

The maximum acidity 0.74% was observed in T₀ (CONTROL (sand, soil, coir pith)) and minimum acidity 0.59% was recorded in Soil T₉ (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g). The similar findings also reported by (Singh *et al.*, 2008) who noticed that the fruit harvested plant receiving vermicompost were TSS and ascorbic acid increases, acidity decreased and color more attractive.

The maximum ascorbic acid (71mg/100g) was observed in T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) and minimum ascorbic acid (25.7mg/100g) was recorded in T₀ (CONTROL (sand, soil, coir pith)). Increased ascorbic acid at better tiers of nitrogen would possibly have resulted in view that absorption of nitrogen can be exerted regulatory function as a vital and all through ripening of culmination the carbohydrate reserves of the roots and stem are drawn upon closely through culmination which would possibly have resulted into better ascorbic acid in culmination. Increased ascorbic acid in culmination is the settlement with the findings of EI-Hamid *et al.*, (2006) who mentioned that the utility of P.S.B. on strawberry resulted growth in ascorbic acid and Singh *et al.*, (2008) who mentioned that the fruit harvested from plant receiving vermicompost have been farmer, TSS and ascorbic acid increases, acidity reduced and satiation greater attractive.

The greatest net return was obtained 2,29,050 Rs/100m² in T₉ (Soil (50%) +Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) and cost benefit ratio was found (3.34) in T₉ (FYM+ Vermicompost+ PSB+ *Azotobacter*). The minimum cost benefit ratio was recorded (1.16) in T₀ (CONTROL (sand, soil, coir pith)). Many workers also reported the higher net return and cost benefit ratio with the application of organic fertilizers and bio fertilizers viz. Yadav *et al.*, (2010), Verma and Rao, (2012) in strawberry.




Fig 1: Strawberry plants under vertical farming system

Table2: Effect of biofertilizers on yield and quality attributes of strawberry under vertical system

Treatment	Treatment combination	Number of fruits per plant (90 days)	Yield plant ¹ (gram) (90 days)	Fruit diameter (cm) (90 days)	Fruit length (cm) (90 days)	TSS (° Brix) (90 days)	Acidity (%) (90 days)	Ascorbic Acid (mg/100g) (90 days)	Net return (₹ /100 ²)	Benefit: Cost ratio
T ₀	CONTROL (sand, soil, coir pith)	11.89	216.08	2.47	3.1	6.03	0.74	25.7	15,700	1.16
T ₁	Sand (25%) + Soil (50%) + VC (25%) + <i>Azoto</i> . (2g)	15.56	333.82	2.8	3.57	7.03	0.71	38.7	74,350	1.78
T ₂	Sand (25%) + Soil (50%) + VC (25%) + PSB (2g)	13.56	268.17	2.57	3.27	6.36	0.74	35.3	40,550	1.41
T ₃	Sand (25%) + Soil (50%) + VC (25%) + <i>Azosp.</i> (2g)	14.33	294.56	2.74	3.43	6.77	0.69	53.3	54,300	1.57
T ₄	Soil (50%) + coir pith (25%) + FYM (25%) + <i>Azoto</i> . (2g)	18.33	457.72	3.23	3.97	8.44	0.64	50.7	1,37,170	2.41
T ₅	Soil (50%) + coir pith (25%) + FYM (25%) + PSB (2g)	16.89	383.04	2.97	3.7	7.6	0.66	46.3	98,970	2.01
T ₆	Soil (50%) + coir pith (25%) + FYM (25%) + <i>Azosp.</i> (2g)	17.56	419.79	3.03	3.84	8.06	0.65	42.3	1,17,920	2.21
T ₇	Soil (50%) + FYM (50%) + PSB (2g) + <i>Azoto</i> . (2g)	19.89	554.33	3.47	4.33	9.1	0.60	62	1,86,740	2.93
T ₈	Soil (50%) + coir pith (50%) + <i>Azosp.</i> (2g) + PSB (2g)	19	511.30	3.33	4.03	8.6	0.61	54.3	1,64,350	2.69
T ₉	Soil (50%) + VC (50%) + <i>Azoto</i> . (2g) + <i>Azosp.</i> (2g)	21	638.40	49.56	4.46	9.57	0.58	71	2,29,050	3.34
F-test		S	S	S	S	S	S	S		
SE. d (+)		1.77	54.45	1.74	0.23	0.36	0.04	2.93		
CD (5%)		3.71	114.4	3.66	0.47	0.76	0.09	6.16		

VC: vermicompost, Azoto: *Azotobacter*, Azosp: *Azospirillum*, PSB: Phosphate solubilizing bacteria



Graph 2: Effect of biofertilizers on yield and quality attributes of strawberry

CONCLUSION

On the basis of present investigation, it is concluded that the treatment T₉ (Soil (50%) + Vermicompost (50%) + *Azotobacter* (2g) + *Azospirillum* (2g)) was found the best in terms of vegetative growth (Plant height (cm), Plant spread (cm), Number of leaves, Leaf area)(cm²) and yield and quality parameters (Number of fruits per plant Yield plant⁻¹(gram) Fruit diameter (cm), Fruit length (cm), TSS (°Brix), Acidity (%), Ascorbic Acid (mg/100g)Net return (₹ /100²), Benefit: Cost ratio).The maximum gross return (Rs. 3,26,800), net return (Rs. 2,29,050) and Benefit cost ratio (1: 3.34) was also observed in treatment T₉ followed by (2.93) in T₇ (Soil (50%) + FYM (50%) + PSB (2g) + *Azotobacter* (2g)) under vertical farming system.

Some urban planners and agricultural leaders have argued that cities must produce food internally in order to manage the demand-supply ratio and avoid falling food prices. From the above research work it is also resulted that how vertical farming can play a beneficial role in urban farming. Not only it will utilize the land requirement in urban areas but also helpful in doubling the income through the vertical farming. It can be easily set up on the home's wall and roofs by reducing the space.

REFERENCES:

- Al-Kodmany, K. (2018). *The Vertical City: A Sustainable Development Model*; WIT Press: Southampton, UK
- Arancon, N.Q. and Edwards, C.A. (2005) Effects of Vermicomposts on Plant Growth. Soil Ecology Laboratory, The Ohio State University, Columbus.
- Arancon, N.Q., Edwards, C.A. and Bierman, P. (2006). Influences of Vermicomposts on Field Strawberries: Effects on Soil Microbial and Chemical Properties. *Bioresource Technology*, **97**, 831-840.
- Atiyeh R., S. Lee, C. Edwards, N. Arancon, J. Metzger (2002). The influence of humic acids derived from earthworm-processed organic wastes on plant growth. *Bioresource technology* **84** (1): 7-14.
- Badr, A.C., Genet, P., Dunand, F.V., Toussaint, M.L., Epron, D.A., & Badot, P.M. (2003). Effect of Copper on growth in Cucumber plants and its relationships with carbohydrate accumulation and change in ion contents. *Journal Plant Science*. **166**: 1213 – 1218.
- Bambal, A.S., Verma, R.M., Panchbhai, D.M., Mahorkar, V.K. and Khankhane, R.N., (1998) Effect of biofertilizers and nitrogen levels on growth and yield of cauliflower (*Brassica oleracea* var. Botrytis). *Orissa Journal Horticulture* **26**(2): 14-17.
- Bareae, J.M., Navorvo, E., and Montonya, E., (1976). Production of plant growth regulators by rhizosphere phosphate solubilizing bacteria. *J. Appl. Bacteriol.* **40**: 129-134
- Cassán, F., V. Sgroy, D. Perrig, O. (2008). Masciarelli and V. Luna. Phytohormone production by *Azospirillum* spp. physiological and technological aspects of plant growth promotion. In *Azospirillum* spp.: cellphysiology, plant interactions and agronomic research in Argentina; 61-86.
- Dadashpour, A. and Jouki, M. (2012). Impact of Integrated Organic nutrient handling on fruit yields and quality of strawberry cv. Kurdistan in Iran. *J. Orntl. & Hort. Plants*, **2**(4): 251-256.
- Darby, H. M., Stone, A. G., and Dick, R. (2004). Compost and manure mediated impacts on soilborne pathogens and soil quality. *Soil Sci. Soc. Am. J.* **70**, 347–358. doi: 10.2136/sssaj2004.0265
- Ellingsen E, Despommier D (2008). The Vertical Farm-The origin of a 21st century Architectural Typology. *CTBUH Journal* 2008;**3**:26-34.
- Evanylo, G., Sherony, C., Spargo, J., Starner, D., Brosius, M., and Haering, K. (2008). Soil and water environmental effects of fertilizer-, manure-, and compost-based fertility

- practices in an organic vegetable cropping system. *Agric. Ecosyst. Environ.* 127, 50–58. doi: 10.1016/j.agee.2008.02.014
- Gajbhiye, R.P., Sharma, R.R. and Tewari, R.N. (2003) Effect of bio-fertilizers on growth and yield parameters of tomato. *Indian Journal Horticulture* **60**(4): 368-371.
- Galletta GJ, Maas JL (1990) Strawberry genetics. *HortScience* **25**:871-879
- K Grappelli, D.H., Galli, E. and Tomati, U. (1987) Earthworm casting effect on *Agaricus bisporus* fructification. *Agrochimica* **21**: 457- 462.
- Kim, S. Y., Pramanik, P., Bodelier, P. L. E., and Kim, P. J. (2014). Cattle manure enhances methanogens diversity and methane emissions compared to swine manure under rice paddy. *PLoS ONE* 9: e113593. doi: 10.1371/journal.pone.0113593
- Kravchenko, A. N., Snapp, S. S., and Robertson, G. P. (2017). Field-scale experiments reveal persistent yield gaps in low-input and organic cropping systems. *Proc. Natl. Acad. Sci. U.S.A.* 114, 926–931. doi: 10.1073/pnas.1612311114
- Kumar, A., Kumar, J. and Babu, R. (2007). Effect of inorganic and biofertilizers on growth, yield and quality of tomato. *Progressive Agriculture* **7**(1&2): 151-152.
- Nazir, N.; Singh, S.R. Aroosa, K.; Masarat, J. and Shabeena, M. (2006). Yield and growth of strawberry cultivar Senga Sengana as influenced by integrated organic nutrient management system. *Enl. & Eco.*, **24**. (3): 651-654. *Notes Bot. Hort. Agrobot. Cluj.*, **37** (1): 139-143.
- Ramasamy, P.K., K. Baskar and S. Ignacimuthu, (2011). Influence of vermicompost on kernel yield of Maize (*Zea mays* L.). *Elixir Agric.* **36**: 3119-3121.
- Rana, R.K. and Chandel, J.S. (2003). Effect of biofertilizers and nitrogen on growth, yield and fruit quality of strawberry. *Progressive Horticulture*, **35**(1): 25-30.
- Sattar, M.A. and Gaur, A.C. (1987). Production of auxins and gibberellins by phosphate dissolving microorganism. *Zentralbl Mikrobiol* **142** (5): 393395.
- Shukla, Y.R., Thakur, A.K. and Joshi, A. (2009) Effect of inorganic and biofertilizers on yield and horticultural traits of tomato. *Indian Journal Horticulture* **66**(2): 285-287.
- Singh, A. and Singh, J.N. (2009). Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. sweet charlie. *Indian Journal Horticulture* **66**(2): 220-224.
- Singh, Y.P., Dwivedi, R., and Dwivedi, S.V. (2008). Effect of biofertilizers and graded dose of nitrogen on growth and flower yield of calendula (*Callendula officinalis*). *Plant Archive* **8**(2): 957-958.
- Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., and Polasky, S. (2002). Agricultural sustainability and intensive production practices. *Nature* 418, 671–677. doi: 10.1038/nature01014

- Tripathi, V.K., Kumar, N., Shukla, H.S. and Mishra, A.N. (2010). Influence of *Azotobacter*, *Azospirillum* and PSB on growth, yield and quality of strawberry cv. chandler. *Abst: National Symposium on Conservation Hort.*, Dehradun, pp. 198-199.
- Umar, I. (2007). Studies on Integrated Nutrient Management in strawberry (*Fragaria × ananassa* Duch.) cultivar Chandler. PhD Thesis, SK University of Agricultural Sciences and Technology of Jammu, Udheywalla, Jammu.
- Umar, I., Wali, V.K., Rehman, M.V., Mir, M.M, Banday, S.A. and Bisati, I.A. (2010a.) Effect of subabul (*Leucaena leucocephala*), urea and biofertilizer application on growth, yield and quality of strawberry cv. Chandler. *Appl. Bio. Res.* **12**: 50- 54.
- Umar, I., Wali, V.K.; Kher, R. and Jamwal, M. (2009). Effect of FYM, Urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler.
- Wang, S.S., Patil, M.T. and Singh, B.R. (1998). Cultivar biofertilizer interaction study in strawberry. *Recent Horticulture* **4**: 43-49.
- Zargar, M.Y., Baba, Z.A. and Sofi, P.A. (2008). Effect of N, P and biofertilizers on yield and physico- chemical attributes of strawberry. *Agro Thesis* **6**(1): 3-8.