

**Efficacy of Newer selected insecticides, *Beaveria bassiana* , Neem oil against
Diamond back moth (*Plutella xylostella*)(L.) in Cabbage(*Brassica
oleracea var capitata*).**

A B S T R A C T

The trail was conducted at Crop research farm , Department of Entomology of Naini Agriculture institute, SHUATS, Prayagraj, (U.P) 2021-22 during rabi to study the efficacy of selected insecticides, *Beaveria bassiana* and Neem oil against Diamond back moth (*Plutella xylostella*) in Cabbage (*Brassica oleracea var capitata*) and the experiment was laid out in randomised block design with eight treatments and each was replicated thrice using a variety Green soccer 546. The treatments are Spinosad 45% SC, Indoxacarb 14.5%SC, Emamectin benzoate 5%SG, Chlorantraniliprole 18.5SC, Fipronil 5%SG, *Beaveria bassiana* , Neem oil 0.3% along with an untreated control. The data on Percent reduction were significantly superior over control but among all treatments, chlorantraniliprole 18.5% SC is best effective for diamond back moth had showing (80.35%) percent reduction followed by Spinosad 45% SC (77.06%), Indoxacarb 14.5SC (73.43%), Emamectin benzoate 5%SG (71.77%), Fipronil (68.26%), *Beaveria bassiana* (67.06%) and Neem oil was least effective showing (61.18%) reduction but superior over control. Among all the treatments, highest yield (314.9q/ha) and Cost-benefit ratio (1:7.59) was recorded in Chlorantraniliprole 18.5 %SC followed by Spinosad 45%SC with a yield of (273.73q/ha) and Cost-benefit ratio (1:6.77) as compared to control .

Keywords : *Beaveria bassiana*, Botanical ,Chlorantraniliprole , Cost-benefit ratio, *Plutella xylostella*

1. INTRODUCTION :

Cabbage is most popular winter vegetable and it belongs to family cruciferae and chromosome number $2n = 24$. Cabbage is rich source of vitamin C, Na, P, and dietary fibre. It is good for people who are suffering from diabetes. India is the largest producer of cabbage after china. India accounts for 8755000 tonnes of productivity in an area of 388000/ha. In India, Uttarpradesh accounts for production of 5.7 million tonnes in an area of 0.72 million ha .

Diamond back moth *Plutella xylostella* (L.) (Family : Plutellidae) is one of the most destructive pest causing yield loss of 52 % in Cabbage^[7] . Cabbage crop is attacked by 37 insect pests and out of that, diamond back moth is the important circumstance pest and it causing globally yield loss of approximately US\$1 billion dollar. Quality and production of cabbage is scale down due to this pest attack^[6]. The caterpillar fedoursly feed on cabbage leaves and make a tunnels underside of leaves and in severe infestation leaf was fully scrapped and leads to decline of yield.

In India, Diamond back moth was first observed in 1914 by Fletcher 1914. Diamond back moth was first documented in Europe region and after that it spreads to almost over the countries but their infestation varies from place to place ^[8].Diamond back moth inaugurate resistant to almost all group of insecticides. Due to the massive use and frequent of insecticides leading to environmental pollution, residual effect , decline in natural enemies population, Health hazards, insecticide resistance, so examing all these points, new molecules of insecticides are introduced to break down the chain of these parameters. especially, avermectins, pyrazoles, microbes, spinosyns, biopesticides playing a major role in managing the diamond back moth.

Diamond back moth has developed resistant to all types of insecticides so it is necessary select New molecules which are highly toxic to insect pests even at low dosages and these new molecules are satisfisying the ecofriendly qualities, easy availability in local region, less residual effect, non toxic to mammals, ecological safety.

By considering above points newer pest management plan was implemented comprising of insecticides/ biopesticide having different mode of action and different formulation were evaluated against major pest of cabbage. Therefore keeping above facts , recent advance research was undertaken “Efficacy of Newer Selected insecticides, *Beauveria bassiana*, Neem oil against diamond back moth *Plutella xylostella* (L.) in cabbage (*Brassica oleracea* var. *capitata*) and to Calcualate Cost Benefit ratio.

2. MATERIALS AND METHODS :

2.1. Experimental site and Location :

The Crop research farm was located on the right side of Rewa Road at 25° 22' 15.888" North Latitude and 81°51' 31.4712" East longitude and is about 98 m above mean sea level and situated 5km away on the right bank of Yamuna river.

2.2. Climate Condition :

The climate at Prayagraj is typical subtropical which prevails in the eastern part of Uttar Pradesh. The extremes of both summer and winter are experienced here. The maximum temperature was recorded during summer up to 47°C and the minimum temperature was recorded during winter up to 1.5°C. All necessary facilities for cultivation of crop were made available at research farm.

2.3. Experimental Details :

The experiment was laid out in a randomized block design with eight treatments replicated three times using variety Green soccer 546 in a plot size of (2m×2m) at a spacing of (60×45cm) with a recommended package of practices excluding plant protection. The Efficacy of Selected Insecticides, one biopesticide i.e. *Beauveria bassiana* (Almax) and 1 botanical i.e., Neem oil and five insecticides include Spinosad (Tracer), Indoxacarb (oxadizine), Emamectin benzoate (Rebel), Chlorantraniprole (coragen), Fipronil (Regent) were purchased from local pesticide traders. For comparison untreated check was included. Treatments application were started once the pest level cut across the ETL i.e., larvae per plant then the second application was given at 15 days interval respectively.

2.4. Assesment of pest population :

The insect larva was count from randomly selected plant in every plot and population per 5 plants was noted. After that mean of three replications was calculated for each treatment and the same was done with the untreated plot. The population of *Plutella xylostella* was recorded before 1day spraying and on 3rd day, 7th day and 14th day after insecticidal application. Then Percent reduction was calcualated by this formula :

Percent Reduction (%) formula :
$$\frac{T_a - T_b}{T_a} \times 100$$

Where ,

T_a = Number of pests individuals on particular day after application.

T_b = Number of pest individuals on particular day after application.

P = Percent reduction in population of pest.

2.5. Marketable head yield :

Healthy cabbage heads were harvested when they reached proper marketable size and weight from each treatment and it was evaluated for quintal per hectare and subjected to statistical analysis for variance.

2.6. Statistical Analysis :

The data recorded on larval population of Diamondback moth (*Plutella xylostella*) and yield of cabbage of each treatment based on replication wise were subjected to analysis of variance.

2.7. Cost-Benefit ratio :

Cost effectiveness of each treatment was evaluated based on net returns. Net return of each treatment was worked out by deducting total cost of the treatment from gross returns. Total cost of production included both cultivation as well as plant protection charges.

Table.1.Particulars of treatments used:

TREATMENT NO.	TREATMENT	DOSE	TRADE NAME	GROUP
T 1	Spinosad 45SC	0.5ml/L	Tracer	Spinosyn
T 2	Indoxacarb 14.5SC	0.25ml/L	Kareplus	Oxdiazine
T 3	Emamectin Benzoate 5% SG	0.45ml/L	Rebel	Avermectin
T 4	Chlorantraniliprole 18.5%SC	0.1ml/L	Coragen	Diamide
T 5	Fipronil 5% SC	25- 50gma.i./ha	Regent	Phenylpyrazole
T6	<i>Beauveria bassiana</i> (1X10 ⁸ CFU/gm)	5gm/L	Almax	Biopesticide
T 7	Neem oil 0.3%	3ml/L	Neem Aura	Botanical
T 8	Control	————	————	————

3. RESULTS :

3.1. Percent reduction of Diamond back moth after first spray :

The results in (Table 2) on mean percent reduction of Diamondback moth on 3rd, 7th, 14th day after first spray reporting that all the treatments were significantly superior over control. Around all the treatments, Chlorantraniliprole 18.5%SC recorded best effective for Diamondback moth (DBM) had showing maximum (70.23%) percent reduction followed by Spinosad 45%SC (67.38%) and Indoxacarb 14.5SC (63.51%) was observed par with Emamectin Benzoate 5% SG (61.68%)on percent reduction whereas , Fipronil 5% SC(57.72%), *Beauveria bassiana* (1X10⁸ CFU/gm) (56.58%) was found par with each other on percent reduction over control. Neem oil 0.3% (50.35%) was found least effective around all treatments on percent reduction but significantly superior over control.

3.2. Percent reduction of diamond back moth after second spray :

The data (Table 2) on mean percent reduction of Diamond back moth on 3rd, 7th, 14th day after second spray reporting that all the treatments were significantly superior over control. Around the treatments Chlorantraniliprole 18.5%SC recorded best effective for Diamondback moth had showing maximum (90.38%) percent reduction followed by with Spinosad 45% SC (86.74%) and Indoxacarb 14.5% SC (83.35%) was observed par with Emamectin Benzoate 5% SG (81.86%)on percent reduction whereas, Fipronil 5% SC (78.80%), *Beauveria bassiana* (1X10⁸ CFU/gm) (77.55%) was found par with each other on percent reduction over control, Neem oil 0.3% was found least effective among all treatments with minimum reduction of (72.01%) but significantly superior over control.

3.3. Percent reduction of diamond back moth after first and second spray :

The data (Table 2) on mean percent reduction of Diamond back moth after first and second spray reporting that all the treatments were significantly superior over control. Around the treatments Chlorantraniliprole 18.5%SC recorded best effective for Diamondback moth had showing maximum (80.35%) percent reduction followed by with Spinosad 45% SC (77.06%) and Indoxacarb 14.5% SC (73.43%) was observed par with Emamectin Benzoate 5% SG (71.77%)on percent reduction whereas, Fipronil 5% SC (68.26%), *Beauveria bassiana* (1X10⁸ CFU/gm) (67.06%) was found par with each other on percent reduction over control, Neem oil 0.3% was found least effective among all treatments with minimum reduction of (61.18%) but significantly superior over control.

3.4 Cost- benefit ratio :

All the treatments were resulted very effective and significantly superior over control. The data (Table 3) on cabbage head yield open up that Chlorantraniliprole 18.5%SC recorded highest yield (283.6q/ha) followed by Spinosad 45%SC (273.71q/ha) Indoxacarb 14.5%SC (233.48 q/ha), Emamectin benzoate 5% SG (221.72 q/ha), Fipronil 5% SC (211.27 q/ha), *Beauveria bassiana* (1X10⁸CFU/gm) (201.57 q/ha), Neem oil 0.3% (194.90 q/ha) when compared compared to control (165.15q/ha).

When the Cost benefit ratio analysis was carried out, an incredible results were obtained. Chlorantraniliprole 18.5%SC acquire highest CBR (1:7.59) followed by Spinosad 45%SC (1:6.77), Indoxacarb 14.5%SC(1:6.33), Emamectin Benzoate 5% SG (1:5.25), Fipronil 5% SC(1:5.78), *Beauveria bassiana* (1X10⁸ CFU/gm) (1:5.50), Neem oil 0.3% (1:5.1) as compared to control (1:4.32)

4. Discussion :

The present study entitled “Efficacy of selected Newer insecticides, *Beaveria bassiana* and Neem oil against diamond back moth(*Plutella xylostella*) in cabbage(*Brassica oleracea var capitata*) .

The data (Table2) on percent reduction of Diamondback moth on Cumulative mean of first and second spray reported that all the treatments were significantly superior over control. Around all the treatments used, Maximum reduction of diamond back moth was observed in Chlorantraniliprole 18.5%SC (80.35%) recorded best effective this results supported by **Dostara et al.,(2017)**^[4] followed by Spinosad 45%SC (77.06%) **Mandal et al., (2009)**^[14] and Indoxacarb 14.5%SC (73.43%) supported by **Jaishree banjaree (2017)**^[3], Emamectin Benzoate 5% SG (71.77%) **AD et al (2014)**^[2], Fipronil 5% SC(68.26%) supported **Deivendram et al., (2007)**^[5], *Beauveria bassiana* (1X10⁸ CFU/gm) (67.06%)supported by **Shelton et al., (1998)**^[17]. **Nikhil et al., (2020)**^[1] reported that Neem oil 0.3% was found least effective among all treatments with minimum percent reduction of (61.18%) but significantly superior over control .

The data (Table 3) on cabbage head yield open up that Chlorantraniliprole 18.5%SC recorded highest yield (283.6q/ha) finding supported by **Dostara et al.,(2017)**^[4] followed by Spinosad 45%SC (273.71q/ha)supported by **Sawant et al., (2018)**^[18], Indoxacarb 14.5%SC (233.48 q/ha) results supported by **Nikhita et al.,(2021)**^[7] Emamectin benzoate 5% SG (221.72 q/ha) **Prasad et al., (2006)**^[11], Fipronil 5% SC (211.27 q/ha) supported by **Deivendram et al., (2007)**^[5] *Beauveria bassiana* (1X10⁸CFU/gm) (201.57 q/ha) supported by **Shelton et al.,(1998)**^[17]. Neem

oil 0.3% (194.90 q/ha) results supported by **Devi *et al.*, (2017)**^[19] when compared compared to control (165.15q/ha).

The present above studies revealed that Chlorantraniliprole, Spinosad , Indoxacarb , was superior and they were highly effective for diamond back moth larval population whereas ,Emamectin benzoate, Fipronil, *Beauveria bassiana* was moderately effective against diamond back moth. **Deivendran *et al.*, (2007)**^[5] supported our finding that fipronil is moderately effective against Diamond back moth. **Yadav *et al.*, (2017)**^[20] supported our finding that indoxacab had better effective than neem but it is less efficient than spinosad. **Nikam *et al.*, (2014)**^[15] supported our finding that spinosad has better efficacy and highly toxic to DBM. **Lal and meena (2001)**^[13] supported our finding that chlorantraniliprole is an commanding molecule against diamond back moth. **Mandal *et al.*, (2009)**^[14] supported our finding that spinoas is superior molecule against diamond back moth .

Our results, justify that chlorantraniliprole is best effective insecticide for controlling diamond back moth showing maximum reduction supported by **Venkestwarulu (2011)**^[19] and **Jakhar *et al.*, (2019)**^[10] supported that Spinosad is effective for Diamondback moth had showing maximum reduction of larval population.

The highest yield and cost benefit ratio was recorded in Chlorantraniliprole (283.6q/ha) (1:7.59) respectively and this result supported by **Sawant *et al.*,(2018)**^[18] followed by Spinosad (273.71q/ha) (1:6.77) this result supported by **Purushotam *et al.*, (2020)**^[16].

Table.2. Efficacy of different insecticides, *Beauveria bassiana* and Neem oil against diamond back moth (*Plutella xylostella*) in Cabbage (*Brassica oleracea var capitata*)

PERCENT REDUCTION OF LARVAL POPULATION										
		FIRST SPRAY				SECOND SPRAY				Cumulative mean
	TREATMENTS	3DAS	7DAS	14DAS	MEAN	3DAS	7DAS	14DAS	MEAN	
T ₁	Spinoasd 45%SC	59.60 ^a	73.33 ^{ab}	69.23 ^{ab}	67.38 ^b	77.87 ^{ab}	90.10 ^b	92.26 ^b	86.74 ^b	77.06^b
T ₂	Indoxacarb 14.5%SC	55.77 ^b	69.52 ^{bc}	65.47 ^{bc}	63.51 ^c	74.15 ^{bc}	87.34 ^{bc}	88.56 ^c	83.35 ^c	73.43^c
T ₃	Emamectin benzoate 5%SG	53.62 ^b	67.61 ^{cd}	63.62 ^{cd}	61.68 ^d	72.31 ^{cd}	86.49 ^c	86.78 ^{cd}	81.86 ^c	71.77^d
T ₄	Chlorantraniliprole 18.5%SC	62.49 ^a	76.18 ^a	72.04 ^a	70.23 ^a	80.67 ^a	93.81 ^a	96.66 ^a	90.38 ^a	80.35^a
T ₅	Fipronil 5% SC	49.97 ^c	63.80 ^{de}	59.91 ^{de}	57.72 ^e	68.66 ^{de}	82.87 ^d	84.87 ^d	78.80 ^d	68.26^e
T ₆	Beaveria bassiana (1x 10 ⁸ CFU/gm)	48.03 ^c	62.85 ^e	58.96 ^e	56.58 ^f	66.83 ^e	81.92 ^d	83.92 ^d	77.55 ^d	67.06^e
T ₇	Neem oil 0.3%	42.29 ^d	56.37 ^f	52.40 ^f	50.35 ^g	61.19 ^f	75.45 ^e	79.40 ^e	72.01 ^e	61.18^f
T ₀	Control	-----	-----	-----	-----	-----	-----	-----	-----	-----
	Sem+ ₋	1.66	1.46	1.44	0.20	1.90	1.32	1.62	1.20	0.27
	CD (0.05)	3.62	3.20	3.14	0.45	4.08	2.88	3.54	1.53	1.38

Table.3. Economics of Cultivation :

Sr. No:	Treatment	Yield of q/ha	Market price q/(₹)	Total cost of yield (₹)	Common cost (₹)	Treatment cost (₹)	Net profit	Total cost	B:C Ratio
T1	Spinosad 45SC	273.71	750	2,05,283	26494	3800	174989	30,294	1:6.77
T2	Indoxacarb 14.5SC	233.48	750	1,75,110	26494	1145	147471	27,639	1:6.33
T3	Enamectin Benzoate 5% SG	221.72	750	166290	26494	5175	134621	31,669	1:5.25
T4	Chlorantraniliprole 18.5%SC	283.6	750	212700	26494	1495	186206	27,989	1:7.59
T5	Fipronil 5% SC	211.27	750	158453	26494	910	131049	27,404	1:5.78
T6	<i>Beauveria bassiana</i> (1X108 CFU/gm)	201.57	750	151178	26494	980	123704	27,474	1:5.50
T7	Neem oil 0.3%	194.9	750	146175	26494	2112	115256	28,606	1:5.10
T0	Control	152.15	750	123750	26494	-----	109369	26,494	1:4.32

CONCLUSION :

From the analysis of present study it was concluded that Chlorantraniliprole 18.5%SC (0.1ml/L) recorded best and proved best effective for Diamondback moth (DBM) among all treatments followed by Spinosad 45SC (0.5ml/L) ,Indoxacarb 14.5SC ,(0.25ml/L),Emamectin Benzoate 5% SG (0.45gm/L) ,Fipronil 5% SC (25-50gma.i./ha) in managing (*Plutella xylostella*) in cabbage (*Brassica oleracea*) on Percent reduction of larval population. Therefore, the biopesticide i.e. *Beauveria bassiana* (Almax) and botanical i.e., Neem oil may be useful in devising proper integrated pest management strategy against on Diamond Back Moth (*Plutella xylostella*).

Acknowledgement :

The authors are thankful to the Department of Entomology, Naini Agriculture institute, Sam Higginbottom university of agriculture and technology sciences, Prayagraj-211007, Uttarpradesh, India for providing us necessary facilities to undertaken the studies.

COMPETING INTERESTS :

Authors have declared that no competing interest exist .

REFERENCES :

1. **Auti, N. K. and Ashwani, K. (2020).**Comparative efficacy of certain chemicals with biopesticides against diamondback moth, *Plutella xylostella*(L.) in cabbage, *Brassica oleracea* (L.) *Journal of Entomology and Zoology Studies*, 8(6): 1350-1353.
2. **AD, G., Bhosle, B. B., Bokan, S. C., and Bhede, B. V. (2018).**Efficacy of newer insecticides against diamondback moth *Plutella xylostella* on cauliflower. *International Journal of Entomology Research* , 3(5): 45- 46
3. **Bajare, J. (2017).** Studies on insect pests of cabbage with special reference to seasonal incidence and management of diamondback moth.
4. **Dotasara, S. K., Agrawal, N., Singh, N., and Dinesh, S. (2017).** Efficacy of some new insecticides against diamond back moth *Plutella xylostella* (L.) on cauliflower. *International Journal of Current Microbiology and Applied Sciences*, 6(5):1958-1963.
5. **Deivendran, A., Yadav, G. S., and Rohilla, H. R. (2007).** Efficacy of some insecticides against *Plutella xylostella* (L.) on cauliflower. *Journal of Insect Science-Ludhiana*. 20(1):102.
6. **Dhanalakshmi, D. N., and Mallapur, C. P. (2008).** Evaluation of promising molecules against sucking pests of okra. *Annals of Plant Protection Sciences*, 16(1): 29-32.
7. **Gaddam, N. R., Srivastava, V. K., Tayde, A. R., and Tripathi, A. (2021).** Comparative efficacy of microbials and botanicals against diamondback moth *Plutella xylostella* (L.) on cabbage. *Journal of Entomology and Zoology Studies*, 9(1): 497-499.

8. **Grzywacz,D.,Rossbach,A.,Rauf, A.,Russel,D.A., Srinivasan,R.and Shelton, A.M.(2010)** Current control methods for diamond back moth and other brassica insect pests and the prospects for improved management with lepidopteran-resistant Bt vegetable brassicas in Asia and Africa. *Crop Protection* , 29: 68-79
9. **Huirem, .D. and Tayde A.R. (2017).** Comparative Efficacy of Bio-Agents and Botanicals on the Management of Diamondback Moth *Plutella xylostella* (L.) on Cabbage under Allahabad Agro climatic Conditions. *International Journal of Current Microbiology and Applied Sciences*, 6(7):711-716.
10. **Jakhar, M., Singh, S. K., and Choudhary, A. L. (2019).** Evaluate of bio-efficacy on different insecticides against diamondback moth on cauliflower. *Journal of Entomology and Zoology Studies*, 7(1): 1077-1081
11. **Kumar, P., and Devappa, V. (2006).** Bio-efficacy of Emamectin Benzoate 5% SG (Proclaim) against diamondback moth in cabbage. *Pestology*, 30(2): 23-25.
12. **Krishnamoorthy, A.(2004)** Biological control of diamondback moth *Plutella xylostella* (L.),an Indian scenario with reference to past and future strategies. *In Proceedings of the International Symposium Montpellier, France, CIRAD*, 204 - 11.
13. **Lal, O.P. and Meena, R.K.(2001).** Effects of certain insecticides against diamondback moth, *Plutella xylostella* (L.) on cabbage under field condition. *Pesticide Research Journal* 13(2): 242-246.
14. **Mandal, S. K., Randhir, K., Sudhir, D., and Vinod, K. (2009).** Field evaluation of some newer insecticides against the diamondback moth, *Plutella xylostella* (L.), on cauliflower. *Pest Management and Economic Zoology*, 17(1): 105-108.
15. **Nikam, T.A., Chandele, A.G., Gade, R.S. and Gaikwad, S.M.(2014)** Efficacy of chemical insecticides against diamond back moth, *Plutella xylostella* (L.) on cabbage under field condition. *Trends in Biosciences*, 7(12): 1196-1199.
16. **Sharma, P., Jangir, H., Yadav , D., and Jat , S.M. (2020)** Economics of Different Insecticidal Treatments against Diamond back Moth on Cabbage.

International Journal of Current Microbiology and Applied Sciences, 9(2):
2327-2334

17. **Shelton, A. M., Vandenberg, J. D., Ramos, M., and Wilsey, W. T. (1998).** Efficacy and persistence of *Beauveria bassiana* and other fungi for control of diamondback moth (Lepidoptera: Plutellidae) on cabbage seedlings. *Journal of Entomological Science*, 33(2): 142-151.
18. **Sawant, C. G., and Patil, C. S. (2018).** Bio-efficacy of newer insecticides against diamondback moth *Plutella xylostella* (L.) in cabbage at farmer's field. *International Journal of Current Microbiology and Applied Sciences*, 7(7): 2986-2998.
19. **Venkateswarlu, V., Sharma, R. K., and Sharma, K. (2011).** Evaluation of eco-friendly insecticides against major insect pests of cabbage. *Pesticide Research Journal*, 23(2): 172-180.
20. **Yadav, S., Raju, S., Yadav, M., Srivastava, A., and Dwivedi, P. (2017).** Bio-efficacy of new insecticidal molecules against the Diamondback moth *Plutella xylostella* (L.) on cauliflower. *Journal of Experimental Zoology*, 20(1): 465-469.

Author Details :

- 1) Tejasri Kommoji
Department of Entomology
Naini Agriculture Institute
Sam Higginbottom University of Agriculture and Technology Sciences,
Uttar Pradesh-211007 (India)
- 2) Dr. Anoorag R. Tayde
Assistant Professor
Department of Entomology
Naini Agriculture Institute
Sam Higginbottom University of Agriculture and Technology Sciences
Uttar Pradesh-211007 (India)

