

Effects of GA₃ and NAA on growth and yield of brinjal (*Solanum melongena* L.) cv.

Kashi Sandesh

ABSTRACT

A field experiment was conducted to find out the Effects of two concentrations of GA₃ and NAA and their combinations on growth and yield of brinjal (*Solanum melongena* L.) cv. Kashi Sandesh. The experiment was carried out at the Horticultural Research Farm, Department of Horticulture, Udai Pratap (Autonomous) College, Varanasi during the year 2020-21. The experiment was laid out in Randomized Block Design with three replications and nine treatments. Higher growth attributing characters viz. maximum plant height (71.14 cm), Number of leaves per plant (71.65), Number of branches (12.77), Days to 50% flowering (43.12) and yield & yield attributing characters viz. Number of fruit per plant (16.17), Fruit length (11.66 cm), Fruit weight (180.48 g), Fruit diameter (9.48 cm), Fruit yield (2.91 kg/plant), Fruit yield (29.22 kg/plot), Fruit yield (383.95 q/ha) were recorded under T₈- NAA + GA₃ (40 ppm + 50 ppm). Similarly, maximum Net return (421750.86 Rs./ha) and B:C ratio (2.73) were recorded under T₈- NAA + GA₃ (40 ppm + 50 ppm) followed by treatment T₇- and T₆. T₉- Control was produced poor performance among all the treatments.

Key Words-Brinjal, PGRs, Growth, Yield and Kashi Sandesh.

INTRODUCTION

One of the most prevalent, well-liked, and important vegetable crops growing in India and other areas of the world is brinjal (*Solanum melongena* L.). In India and other Asian nations including Bangladesh, Pakistan, and the Philippines, the brinjal is widely grown. China, Turkey, Japan, Egypt, Indonesia, Iraq, Italy, Syria, and Spain are major countries that produce brinjal. Indian-origin farmed brinjal has been grown for a very long period. It is one of the most widespread and well-liked vegetable crops in India, except where there are greater altitudes. India produces 83.47 lakh tonnes of brinjal over an area of 5.02 lakh ha, ranking second only to China as the world's top producer. Orissa, West Bengal, Bihar, and practically all other states cultivate brinjal to the greatest extent. The production of brinjal is governed not only by the

inherent genetic yield potential of the cultivars but it is greatly influenced by several environmental factors and cultivation practices.

Brinjal fruits are a reasonable supply of calcium, iron, and vitamins of the 'B' group (Kiran et al. 2010). Additionally prized for its therapeutic benefits, brinjal has been linked to the treatment of intestinal worms, rheumatoid arthritis, leucorrhoea, allergy-induced cough, and liver disease (Das and Barua; 2013). It offers nutritious fiber, vitamins, minerals, carbohydrates, and protein, just as other vegetables. Since brinjal is a warm-season crop, it needs a lengthy warm growth season. It is quite vulnerable to freezing. For effective production, a daily mean temperature of 13–21 °C is ideal. When the temperature drops below 17 °C, the crop's growth is negatively impacted. It can be effectively cultivated both during the rainy and summer seasons, and it can be grown up to 1200 meters above sea level. Plant growth regulators are recognized to have an impact on horticulture crops' increased yields and quality. Recently, the significance of PGRs in raising crop output has come to the attention of the entire world. GA₃ is an important growth regulator that may have many uses to modify the growth, yield and yield contributing characters of plant (Rafeekheret *al.* 2002). Plant growth regulators are used widely to improve plant performance. GA₃ is one of those growth regulators that have positive effect on plant growth through the effect on cell division and elongation (Batlanget *al.*, 2006). It recorded dipping of brinjal seedling roots in NAA at 0.1 or 0.2 ppm for 24 hours influenced growth and development (Vajaet *al.* 2017). The advantage of plant growth regulators like Gibberellic (GA₃), Indol Acetic Acid (IAA), Nephthaline Acetic Acid (NAA), 2-4- Diclorophenoxy Acetic Acid (2,4-D) can be taken to increase the yield of local variety of brinjal.

MATERIAL AND METHODS

The field experiment was carried out in the *Rabi* season of 2020–2021 at the research farm of Department of Horticulture at udaipratap (autonomous) college, Varanasi, Uttar Pradesh which is located Eastern region of the state between 25.3550° North latitude and 82.9753° East longitude. This farm has sufficient irrigation facilities accessible. The lowest temperature during the growing season is between 6 and 21.7⁰C, while the highest temperature during that time is between 17 and 35.1⁰C. During the growing period, relative humidity ranged from 24 to 94 percent. During the trial, average wind speeds ranged from 1.3 to 6.3 km hr⁻¹. During the testing period, the trail location got a total of 43.2 mm of rain in one wet day, providing favourable

conditions for crop development. The experiment consisted of two levels of gibberellic acid (GA_3), two levels of naphthalene acetic acid (NAA) and their combinations was arranged in randomized block design with three replications and nine treatments viz. (T₁) - NAA (20 ppm), (T₂) - NAA (40 ppm), (T₃)- GA_3 (25 ppm), (T₄) - GA_3 (50 ppm), (T₅) – NAA+ GA_3 (20 ppm + 25 ppm), (T₆) – NAA+ GA_3 (20 ppm + 50 ppm), (T₇) - NAA + GA_3 (40 ppm + 25 ppm), (T₈) - NAA + GA_3 (40 ppm + 50 ppm) and (T₉) - Control. Seedlings were uprooted from the nursery beds and transplanted in the main plots at a spacing of 70 cm row to row and 60 cm plant to plant. Transplanting was done in the evening hours immediately followed by irrigation for proper establishment of the seedlings. All the recommended package of practices was followed to raise a healthy crop.

The required weight of the PGRs was taken using electronic sensitive balance and solution was prepared by dissolving in 1 mg L^{-1} . The solution was poured into hand-held sprayer and was directly sprayed on the plants three times at 30, 45 and 60 days after transplanting. Spraying was performed early in the morning to avoid rapid drying of the spray solution, due to transpiration. Data were collected from randomly selected five plants in each row. The collected data includes Plant height (cm), No. of leaf/plant, No. of branches per plant, Days to 50% flowering, No. of fruit per plant, Fruit length(cm), Weight of fruits (g), Fruit diameter(cm), Fruit Yield(Kg/plant), Yield(kg/plot) and Yield(q/ha). Recorded data was analyzed using appropriate method of 'Analysis of Variance (ANOVA)' given by Gomez and Gomez (1984).

RESULTS AND DISCUSSIONS

Effect of treatments on growth attribute of brinjal

Effect of plant growth regulators on growth attribute of brinjalas showed in Table 1. Significantly tallest plant was found in T₈- NAA + GA_3 (40 ppm + 50 ppm) 41.92, 61.20 and 71.14 at 30, 45 and 60 DAT respectively followed by T₇ - NAA + GA_3 (40 ppm + 25 ppm) (36.18) and minimum was found in T₉ - Control (27.18). It might be due to gibberellin can promote the activity of xyloglucan endotransglycosylase (XET) which cause loosening of cell wall and increase cell permeability (Saptari and Dawi 2013). Similar result was also reported by Meena and Dhaka (2003). The no. of leaves per plant increased continuously from 30 to 60 DAT in all the treatments of Kashi Sandesh. At 30 DAT. Higher no. of leaves at 30, 45 and 60 DAT (44.35,

47.12 and 71.65 leaves respectively) were recorded under treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) followed by T₇ - NAA + GA₃ (40 ppm + 25 ppm) and T₆ - NAA + GA₃ (20 ppm + 50 ppm). However, minimum no. of leaves was recorded in treatment (T₉) - Control at 30, 45 and 60 DAT. followed by (68.59 and 67.38 leaves) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas, minimum no. of leaves (41.63 leaves) was recorded in T₉-Control. The green leaf is the site of all physiochemical reactions, and the rise in leaf number was influenced by growth regulators combinations as a result of increased protoplasm content in plants and faster metabolic processes. As a result of the extra nitrogen encouraging vegetative development, the number of green leaves increased. Similar result was also recorded by (Hemlata and Raza 2016) and (Khaleghiet al. 2021). Similarly, higher no. of branches at 30, 45 and 60 DAP (9.49, 10.07 and 12.77 plant⁻¹ respectively) was found with T₈ - NAA + GA₃ (40 ppm + 50 ppm) followed by T₇- GA₃ @ 50 ppm and T₆- NAA @ 40 ppm. Whereas, lowest no. of branches was recorded in treatment T₁- NAA (20 ppm) at 30 DAT and at 45 and 60 DAT minimum branches was recorded under T₉- Control. Such effect of PGRs combination on no. of branches was also reported earlier by (Kropi and Phonglosa 2020) and (Bagaleet al. (2022)). All the treatments were significantly influenced days taken to 50 per cent flowering. The minimum days to 50 per cent flowering (29.66) was obtained in the treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was found significantly superior over other treatments followed by (30.54) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm), whereas, maximum days to 50 per cent flowering (43.12) were noted in T₉- Control. Because of using the plant growth promoters in the right amount, early flowering comes in the plant (Kiranmayiet al. 2014).

Effect of treatments on yield and yield attribute of brinjal

Increment in growth attributing characters were ultimately reflected in yield attributing characters. Among the treatments, (T₈) - NAA + GA₃ (40 ppm + 50 ppm) recorded significantly higher number of fruits plant⁻¹ (16.17) followed (T₇) - NAA + GA₃ (40 ppm + 25 ppm) (15.20 fruits). While, significantly lowest number of fruits plant⁻¹ was recorded with T₁-NAA (20 ppm) (8.67). The maximum fruit length (11.66 cm) was recorded in the treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) followed by (11.29 and 10.25 cm) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas, minimum fruit length (5.45 cm) was recorded in T₉- Control. Similarly, maximum fresh weight (180.48 g) was recorded under

treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) followed by (177.99 and 176.50 g) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm). Whereas minimum fresh weight (163.59 g) was recorded under T₉- Control (Table 2). The maximum fruit diameter (9.48 cm) was recorded in the treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) followed by (8.49 and 7.43 cm) in T₇- NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm) and minimum fruit diameter (3.77 cm) was recorded under T₉- Control. Among all the treatments, (T₈) - NAA + GA₃ (40 ppm + 50 ppm) was recorded higher fruit yield plant⁻¹ (2.91 kg), fruit yield per plot (29.22 kg) and fruit yield quintal per hectare (383.95) followed (T₇) - NAA + GA₃ (40 ppm + 25 ppm). In other hand least fruit yield plant⁻¹, fruit yield per plot and yield quintal per hectare was observed in T₉- Control. Which are influenced by the growth cycle during vegetative stages and mirrored during productive phases. Different growth promoter combinations improved the values of several growth and yield contributing features. Similarly reported by several researchers *viz.*, (Khaleghiet *al.* 2021), (Patel *et al.* 2012), and (Ruidaset *al.* 2022).

Effect of treatments on yield and yield attribute of brinjal

It was revealed that the maximum net return of Rs. 422710.86 q/ha with Benefit:Cost ratio of 2.75 was recorded in T₈ (@ 40 ppm NAA + @ 50 ppm GA₃) followed by T₇-(@ 40 ppm NAA + 25 ppm GA₃) with net income of Rs. 416581.86 q/ha along with Benefit:Cost ratio of 2.71 and minimum net return of Rs.185598.25 q/ha with Benefit:Cost ratio of 1.49 was observed in control.). Net return and benefit cast ratio was more due to higher production fruit yield of brinjal (Table. 2). The effect of PGRs on economics of brinjal was also reported by(Athaneria *et al.*2011), (Veishnavet *al.*2012) and (Vandnaet *al.*2014).

Effect of treatments on fruit colour, fruit shape and fruit size of brinjal

As for fruit sample observed and given below in table 3 we can say that there are no any difference was found in fruit colour, Fruit shape and fruit size with the application of various doses of growth hormone, its combination and control. All the treatment including control was showed Purple colour, Round in shape and Medium size fruits. The similar result was recorded by Khaleghiet *al.*(2021) and Ruidaset *al.*(2022).

CONCLUSION

It can be concluded from the present investigation that PGRs combinations with (NAA - 40 ppm + GA₃ -25 ppm) was produce maximum growth and yield attributing characters at different crop stages. Similarly, highest fruit yield and Net return and benefit cast ratio were recorded with T₈ (@ 40 ppm NAA + @ 50 ppm GA₃).

UNDER PEER REVIEW

Table .1 Effect of PGR on growth attribute of brinjal

Treatment	Plant height (cm)			No. of leaf/plant			No. of branches/plant			Days to 50%flowering
	30DAT	45DAT	60DAT	30DAT	45DAT	60DAT	30DAT	45DAT	60DAT	
T₁	27.96	44.72	61.45	37.41	37.67	54.00	3.38	4.41	5.82	40.57
T₂	31.06	47.66	62.18	38.55	38.84	55.32	4.33	5.11	6.45	37.75
T₃	32.55	51.03	64.42	39.36	40.37	62.53	5.35	5.74	7.34	36.08
T₄	34.52	51.92	65.37	40.51	42.24	64.91	5.84	6.33	8.62	34.43
T₅	34.90	53.22	66.44	41.46	43.16	66.04	6.51	6.74	9.74	32.79
T₆	36.16	54.00	67.61	42.31	43.52	67.38	7.39	7.60	10.53	31.44
T₇	36.18	57.84	69.40	43.12	45.29	68.59	8.40	8.41	11.80	30.54
T₈	41.92	61.20	71.14	44.35	47.12	71.65	9.49	10.07	12.77	29.66
T₉	27.18	37.75	45.47	32.26	35.15	41.63	3.73	4.19	4.86	43.12
SE ±	0.55	0.54	0.28	0.27	0.20	0.46	0.20	0.15	0.20	0.42
CD(5%)	1.67	1.63	0.84	0.81	0.60	1.38	0.60	0.47	0.61	1.28

Table .2Effect of PGR on yield attribute, yield and economics of brinjal

Treatment	No. of fruit per plant	Fruit length(cm)	Weight of fruits (g)	Fruit diameter(cm)	Fruit Yield(Kg/plant)	Yield(kg /plot)	Yield(q /ha)	Net return(Rs./ha)	B:CR atio
T₁	8.67	5.56	166.51	3.94	1.44	21.90	356.61	381785.34	2.49
T₂	9.82	6.28	168.23	3.94	1.65	23.32	362.48	390577.82	2.55
T₃	11.13	7.40	170.65	4.72	1.89	24.14	366.97	397301.82	2.59
T₄	12.00	7.94	171.79	5.49	2.06	25.42	370.44	402470.82	2.62
T₅	12.93	8.88	174.34	6.50	2.25	25.84	373.57	406829.34	2.64
T₆	13.56	10.25	176.50	7.43	2.39	27.33	376.85	412073.34	2.68
T₇	15.20	11.29	177.99	8.49	2.71	28.13	379.84	416581.82	2.71
T₈	16.17	11.66	180.48	9.48	2.91	29.22	383.95	422710.86	2.75
T₉	11.63	5.45	163.59	3.77	1.90	19.91	206.69	185598.25	1.49
SE ±	0.18	0.28	1.18	0.12	0.02	0.18	0.85	-	-
CD(5%)	0.56	0.84	3.54	0.35	0.07	0.55	2.56	-	-

Table .3 Fruitcolour,fruitshapeandfruitsizeasinfluencedbydifferent treatments

Symbol	Treatment	Fruitcolour	Fruitshape	Fruitsize
T ₁	NAA(20ppm)	Purple	Round	Medium
T ₂	NAA(40 ppm)	Purple	Round	Medium
T ₃	GA ₃ (25ppm)	Purple	Round	Medium
T ₄	GA ₃ (50ppm)	Purple	Round	Medium
T ₅	NAA+ GA ₃ (20 ppm +25 ppm)	Purple	Round	Large
T ₆	NAA+ GA ₃ (20 ppm + 50 ppm)	Purple	Round	Large
T ₇	NAA+ GA ₃ (40 ppm +25 ppm)	Purple	Round	Large
T ₈	NAA+ GA ₃ (40 ppm + 50 ppm)	Purple	Round	Large
T ₉	Control	Purple	Round	Small

* NAA-Nephthaline Acetic Acid , GA₃- Gibberellic Acid and ppm- Part Per Million

References

- Athaneriya, M. K., Sengar, N. and Pandey, B. R. (2011). Influence of biofertilizer on growth and yield of chilli. *Vegetable Science*. **38**(1):101-103.
- Bagale, P., Pandey, S., Regmi, P., &Bhusal, S. (2022). Role of Plant Growth Regulator “Gibberellins” in Vegetable Production: An Overview. *International Journal of Horticultural Science and Technology*, **9**(3): 291-299.
- Batlang, V., Emongor, V.E. and Pule-Meulenburg, F. (2006). Effect of benzyladenine and gibberellic acid on yield and yield components of cucumber (*Cucumissativus* L. cv. ' tempo'). *J. Agron.*, **5**(3): 418-423.
- Das, M., &Barua, N. (2013). Pharmacological activities of *Solanummelongena* Linn.(Brinjal plant). *International Journal of Green Pharmacy (IJGP)*, **7**(4).

- Hemlata, P., & Raza, A. K. (2016). Effect of GA₃ and NAA on growth and yield of brinjal (*Solanum melongena* Lin.) cv. Pusa purple long. *Journal of Natural Resource and Development*, **11**(1): 32-35.
- Khaleghi, S., Baninasab, B., Mobli, M., & Ehtemam, M. H. (2021). Effect of plant growth regulators on two different types of eggplant flowers regarding style length and fruit setting. *Spanish Journal of Agricultural Research*, **19**(4): e0906-e0906.
- Kiran, J., Vyakaranahal, B. S., Raikar, S. D., Ravikumar, G. H., & Deshpande, V. K. (2010). Seed yield and quality of brinjal as influenced by crop nutrition. *Indian Journal of Agricultural Research*, **44**(1): 1-7.
- Kiranmayi, P., Jyothi, K. U., Kumari, K. U., Vani, V. S. and Sneetha, D. R. S. (2014). Effect of NAA, 4-CPA and boron on growth and yield of green chilli (*Capsicum annum* L.) var. Lam-353 in summer. *Agrotechnol.* **2**(4):216-222.
- Kropi, J., & Phonglosa, A. (2020). Response of different plant growth regulators on fruit yield of Brinjal. *International Journal of Agriculture, Environment and Biotechnology*, **13**(2): 129-131.
- Meena, S.S., and Dhaka, R.S. (2003). Economics of plant growth regulators in brinjal (*Solanum melongena* L.) under semiarid condition of Rajasthan. *Annals Agric. Res.*, **24** (2): 273-275.
- Patel, J. S., Sitapara, H. H., and Patel, K. A. (2012). Influence of plant growth regulators on growth, yield and quality of tomato and brinjal. *International Journal of Forestry and Crop Improvement*. **3**(2):116-118.
- Rafeekher, M., Nair, S. A., Sorte, P. N., Hatwal, G. P. and Chandan, P. M. (2000). Effect of growth regulators on growth and yield of summer cucumber. *Journal of Soils and Crops*. **12**(1):108-110.
- Ruidas, S., Karmakar, S., Purkait, A., Gangopadhyay, A., Saha, R., Mukherjee, K., & Hazra, D. K. (2022). Preparation, optimization, and testing of biostimulant formulations as stress management tools and foliar applications on brinjal and onion for growth and yield.

- Saptari, R.T., and Dewi, K., (2013). Effect of borax and gibberellic acid on the growth and development of red chilli (*Capsicum annuum* L. “gelora”). *The Third Basic Science International Conference*. (B41):1-3.
- Thompson, C. H, and Kelly, C. W. (1957). *Vegetable Crops*, McGraw – Hall Book Co. Inc, New York, p. 502.
- Vaja, A. D., Patel, J. B., Daki, R. N., & Chauhan, S. A. (2017). Effect of nitrogen and plant growth regulators on seed yield per plant and seed quality parameters in brinjal (*Solanum melongena* L.). *Journal of Applied and Natural Science*, **9**(4):2338-2343.
- Vandana, P. and Verma, L. R. (2014). Effect of spray treatment of growth substances at different stages on growth and yield of sweet pepper (*Capsicum annuum* L.) cv. Indra under green house. *International Journal of Life Sciences Research*. **2**(4):235-240.
- Veishnav, N., Singh, B. K., and Singh, A. K. (2012). Effect of NAA on growth and yield of chilli (*Capsicum annuum* L.). *Environment & Ecology*. **30**(4):126.