

# **Ethanollic** Extract of *Xylopi aethiopica* (African Negro Pepper) Fruit adversely perturbed Semen Qualities in Male Wistar Rats

## **ABSTRACT**

**Aim:** This study was designed to assess the effect of *Xylopi aethiopica* fruit on the sperm qualities of male Wistar rats.

**Methodology:** The fruits of *Xylopi aethiopica* were air-dried and extracted by Soxhlet extractor using **ethanollic** as solvent. The median lethal dose (LD<sub>50</sub>) of the extract was assessed using standard method. Thirty adult Wistar rats were divided into five groups of six rats each. Animals in groups 1, 2, 3, and 4 were treated with 130, 259, 389 and 518 mg/kg body weight of *X. aethiopica* fruit extract respectively, while those in group 5 received normal animal feeds and water only. The administration was done once daily for 28 days via oral route. At the end of 28 days treatment, animals were sacrificed under ether anaesthesia in a desiccator after an overnight fast. The cauda epididymis were separated from both testes and tinged with 2 mL of normal saline then teased the cauda epididymis of each rat. The suspension was mixed through a metallic net to avoid any other tissue contamination. This suspension was used for the determination of the sperm parameters.

**Results:** Ethanollic extract of *Xylopi aethiopica* fruit was observed to significantly perturbed sperm parameters of animals after 28 days of treatment. Sperm count and motility were significantly reduced by *Xylopi aethiopica* fruit in a dose-dependent manner when compared with those of the control group ( $P<0.05$ ). Administration of *Xylopi aethiopica* fruit increased sperm mortality and abnormality when compared with the control animals ( $P<0.05$ ). Seminal pH was decreased by **ethanollic** extract of *Xylopi aethiopica* fruit administration when compared with those in control animals ( $P<0.05$ ).

**Conclusion:** The findings of this study revealed that **ethanollic** extract of *Xylopi aethiopica* fruit adversely perturbed sperm quality of Wistar rats. This might not automatically translate to same effect in human. However, men interested in child-bearing should minimize its consumption.

**Keywords:** Potent contraceptive; sperm qualities; *Xylopi aethiopica* fruit consumption


## **1. INTRODUCTION**

Infertility has been defined to be the inability to get pregnant after one year of unprotected intercourse [1]. Infertility in male has been discovered in 50% of infertile couples [2]. Speroff and Fritz [3] have observe that 55% of the causes of infertility are discovered to be male-related while only 35% are said to be female-related, while the remaining 10% are

infertility of unidentified origin [3]. Some of the causes of decreasing male fertility could be traced to declining levels of androgen, reduced sexual activity, perturbations in semen qualities, especially, motility, morphology, and DNA integrity [4]. Gonadotropin releasing hormone (GnRH) evokes the distribution of gonadotropins i.e. follicle stimulating hormone (FSH) and luteinizing hormone (LH) from the pituitary gland [5]. Luteinizing hormone is a known glycoprotein

that controls the synthesis of testosterone through the extra-tubular Leydig cells. The second gonadotropic hormone, follicle stimulating hormone regulates spermiogenesis as well as spermiocytogenesis by altering Sertoli cells as well as germinal epithelium [6]. The concentrations of these sex hormones are subjected to negative feedback regulation by the gonads [7]. Testosterone is responsible for normal growth and development of male sexual organs, as well as the maintenance of secondary sex characteristics. A high intra-testicular concentration of this hormone is a basic requirement for the synthesis of sperm and its function. Testosterone enhances the motility of sperm as well as epididymis function [8]. Inadequate secretion of FSH and LH by the pituitary gland could lead to interference in the function of the testicles which can contribute to infertility [9].

Semen is an organic fluid with spermatozoa as its constituent. It is released by sexual glands as well as other accessory sex organs of male, and can fertilize the ova of the female. In humans, semen has different constituents outside spermatozoa [10]. Infertility in male could be examined by analyzing semen and hormonal profile [11].



**Fig 1: Sperm Morphology**

Male impotence popularly known as erectile dysfunction (ED) is a frequent health condition which influence the sexual behaviour of several of men all over the world [12,13]. ED is the incapability of a man to attain and sustain an erection good enough for natural satisfactory intercourse.

Sexual dysfunction is a severe medical and social sign that occurs in 10-52% of men and 25-63% of women [14]. It is the iterated failure to get usual sexual intercourse. ED is a common medical issue that may add to infertility cases [15]. ED is negatively impacted by by some

medications such as antihypertensive, antidiabetic agents, antipsychotic, antidepressant therapeutic and antimalarial drugs [16].

Literarily, medications from plant sources are believed to have less side effect when compared with orthodox drugs. This belief comes from the conception of "green is safe" [17,18]. Owing to this perception, medicinal plants are utilized without recourse to their toxicity. The situation is the same with *Xylopia aethiopica* fruit. Despite its use as traditional medication and some animal studies that have been carried out to ascertain its therapeutic uses, not much has been done in evaluating its safety or toxicity of most of its bioactive constituents. However, a qualitative/semi quantitative test for toxicity has been reported [19]. In a related study, the essential oil of the fruits of *X. aethiopica* was reported to be toxic to *Artemia salina* at concentrations between 10 and 1000  $\mu\text{g/mL}$  [20]. Koba *et al.* [21] investigated the *in vitro* cytotoxicity of essential oil from *X. aethiopica* fruits. The test was carried out using the human epidermal cell line HaCaT. In that study, it was identified that at concentrations between 50 and 1500  $\mu\text{g/mL}$ , the examined essential oil did not portray any cytotoxicity instead it caused a noticeable rise in the viability of cells (up to 130 %), indicating their ability to be cytoprotectors or antioxidants. At higher concentrations between 1600 and 3000  $\mu\text{g/mL}$ , a close result was reported. Xylopic acid isolated from the dried fruits of *Xylopia aethiopica*, in a study conducted by Woode *et al.* [22], when administered at doses of 10, 30 and 100 mg/kg to male albino rats, caused significant cytotoxicity by removing all matured spermatozoa, germ cells and other cell in the seminiferous tubules when compared with their control group. However, these impacts were alleviated when the experimental animals were given 14 days to recover. Therefore, their results predict that xylopic acid contains reversible spermatotoxic and antifertility effects at the experimented dosages. An ethanolic extract of a combination of equal quantities of *Alstonia congensis* bark and *Xylopia aethiopica* fruits have been investigated for acute and sub-acute toxicity [23]. In the acute toxicity study, no significant changes in the behaviour and sensory nervous system responses were observed. Similarly, no negative gastrointestinal impacts were noticed in both male and female mice used in the study. At a dose of 20.0 g/kg, all the treated mice survived after the one day of

observation. Therefore, it was concluded that the median acute toxicity value ( $LD_{50}$ ) of the extract must be more than 20.0 g/kg body weight. Although not entirely representative, these findings to some extent show that **ethanolic** extract of *Xylopia aethiopica* fruits is relatively safe [23].

*Xylopia aethiopica* is characterized with numerous chemical components with various medicinal potentials. The chemical components of this plant have been investigated to include saponins, sterols, carbohydrates, glycosides, mucilage, acidic compounds, tannins, balsams, cardiac glycosides, volatile aromatic oils, phenols [24,25], alkaloids, rutin and fixed oils [26,27]. The plant has also been known to contain vitamins such as vitamin A, vitamin B, vitamin C, vitamin D, and vitamin E, and proteins as well as several minerals such as copper, manganese and zinc [25,27]. The impact of the fruit on body weight and glucose concentration of animals has been reported [28]. The fruit has also been reported to induce dyslipidemia [29], hepatotoxicity [30] as well as renal toxicity [31]. This present study focused on examining its impact on the oxidative stress biomarkers of Wistar rats.

## 2. MATERIALS AND METHODS

### 2.1 Collection and Authentication of Plant Materials

The fruits of *Xylopia aethiopica* were obtained from new market in Aba, Abia State and were identified and authenticated by Prof. (Mrs) Margaret Bassey of the Department of Botany and Ecological Studies, University of Uyo with the voucher number UU/PH/4e. The plant was deposited in the Herbarium of the Department of Pharmacognosy and Natural Medicine, University of Uyo, Akwa-Ibom State, Nigeria.

### 2.2 Extraction of Plant Materials

The extraction was carried out in the Post-graduate Laboratory of Department of Pharmacognosy and Natural Medicine, Faculty of Pharmacy, University of Uyo, Nigeria. It was carried out according to the method described by Ogbuagu et al. [32]. The fruits were washed under running tap water to remove contaminants and air-dried. The plant material was pulverized using laboratory blender to provide a greater surface area. The pulverized plant material was

macerated in 250 mL of 99.8% **ethanolic** (Sigma Aldrich) contained in round bottom flask, which was then attached to a Soxhlet extractor coupled with condenser and heating mantle (Isomantle). It was then loaded into the thimble, which is placed inside the Soxhlet extractor. The side arm is lagged with glass wool. The mixture was heated using the heating mantle (Isomantle) at 60 °C and as the temperature increases it begins to evaporate, moving through the apparatus to the condenser. The condensate then drips into the reservoir containing the thimble. Once the level of solvent reaches the siphon it pours back into the flask and the cycle begins again. This continues until it is exhaustively extracted. The process runs for a total of 13 hours. Once it was set up, it was left to run without interruption as long as water and power supply were not interrupted. The equipment was turned on and off and overnight running was not permitted, and the time split over a number of days. The extract was poured into 1000 mL beaker and concentrated to dryness in water bath (A3672-Graffin Student Water Bath) at 35 °C. The total weight of the marc (residue) and the concentrated extract were recorded, these processes took several days. The dried extract was preserved in the refrigerator at 4°C for further analysis.

### 2.3 Determination of Median Lethal Dose ( $LD_{50}$ )

The median lethal dose ( $LD_{50}$ ) of the extract was estimated using albino mice according to the method described by Airaodion et al. [33]. This method involves two phases:

In Phase one, five groups containing five mice each weighing between 20 g and 27g were fasted for 18 hours. They were respectively administered 1000 mg/kg, 2000 mg/kg, 3000 mg/kg, 4000 mg/kg and 5000 mg/kg body weight intraperitoneally (i.p) and were observed for physical signs of toxicity and mortality for 24 hours. A dosage of 1000 mg/kg recorded 0% mortality while 2000 mg/kg, 3000 mg/kg 4000 mg/kg and 5000 mg/kg recorded 100% mortality within 24 hours. Based on the value of phase one, phase two was conducted.

In Phase two, twenty-five albino mice weighing between 20 - 27g were grouped into five of five mice per group and were fasted for 18 hours. Each group was administered 1200 mg/kg, 1400 mg/kg 1600 mg/kg, 1800 mg/kg and 2000 mg/kg

body weight intraperitoneally (i.p) and was observed for physical signs of toxicity and mortality within 24 hours. 1200 mg/kg recorded 0% mortality while 1400 mg/kg, 1600 mg/kg, 1800 mg/kg and 2000 mg/kg recorded 100% mortality within 24 hours. The LD<sub>50</sub> was calculated as geometrical means of the maximum dose producing 0% (a) and the minimum dose producing 100% mortality (b).

$$LD_{50} = \sqrt{at}$$

## 2.4 Experimental Design

Thirty adult male Wistar rats obtained from the University of Uyo, Nigeria were used for this study. They were acclimatized for seven days before the commencement of the experiment. They were weighed and divided into five groups of six rats each. Groups A, B, C, D served as the experimental groups, while group E served as the control. Animals in group A were administered 130 mg/kg body weight (10% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, those in group B were administered 259 mg/kg body weight (20% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, those in group C were administered 389 mg/kg body weight (30% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, those in group D were administered 518 mg/kg body weight (40% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, while those in group E (control) received normal feeds and water only. The administration was done once daily for 28 days via oral route. At the end of 28 days treatment, animals were sacrificed under ether anaesthesia in a desiccator after an overnight fast. The cauda epididymis were separated from both testes and tinged with 2 mL of normal saline then teased the cauda epididymis of each rat. The suspension was mixed through a metallic net to avoid any other tissue contamination. This suspension was used for the determination of the sperm parameters.

## 2.5 Determination of Sperm Qualities

### 2.5.1 Determination of Sperm Count

Sperm count was determined using the haemocytometer method [34]. A 1:20 dilution from each well-mixed sample was prepared by diluting 50 µL of liquefied semen with 950 µL diluent. The latter was prepared by adding 50 g of sodium carbonate (NaHCO<sub>3</sub>), 10 mL of 35% (v/v) formalin and, 0.25 g of trypan blue or 5 mL of saturated aqueous gentian violet to distilled

water and the solution was made up to a final volume of 1000 mL. Both chambers of the haematocytometer are scored and the average count is calculated,

### 2.5.2 Determination of Sperm Motility

Sperm motility was determined according to the method described by Larsen *et al.* [35]. The sample was thoroughly mixed and an aliquot was immediately removed, allowing no time for the spermatozoa to settle out of suspension. The sample was remixed before removing a replicate aliquot. For each replicate, a wet preparation approximately 20 µm deep was prepared. The sample was allowed to stop drifting (within 60 seconds). The slide was examined with phase-contrast optics at ×200 or ×400 magnifications.

### 2.5.3 Determination of Sperm Abnormality

Abnormality of spermatozoa was determined according to the method described by Airaodion *et al.* [36]. A film of semen was prepared on slide. These films on slide were fixed in methanolic. The slides were stained in eosine for 40 minutes. The films were washed in tap water and after drying, the slides were examined under the microscope to see abnormality of spermatozoa.

### 2.5.4 Determination of Sperm Mortality

Sperm mortality was determined as the difference between sperm motility and abnormality.

### 2.5.5 Determination of Seminal pH

Seminal pH was determined using pH paper in the range 6.0 to 10.0 according to the method described by Airaodion *et al.* [37]. The sample was thoroughly mixed and a drop was evenly spread on the pH paper. The colour of the impregnated zone became uniform after about 30 seconds and the colour was compared with the calibration strip to read the pH, and the corresponding value was recorded.

## 2.6. Statistical Analysis

Results are expressed as mean ± standard deviation. The levels of homogeneity among the groups were assessed using One-way Analysis of Variance (ANOVA) followed by Tukey's test. All analyses were done using Graph Pad Prism

Software Version 5.00 and P values < 0.05 were considered statistically significant.

### 3. RESULT

#### 3.1 Median Lethal Dose (LD<sub>50</sub>) Result

The visible symptoms of toxicity seen in the experimental animals included excitation, paw licking, increased respiratory rate, decreased motor activity, gasping and coma which was followed by death. In the first phase of the median lethal dose determination, no mortality was recorded in the group treated with 1000 mg/kg body weight of *X. aethiopica* fruit extract. However, 100 % mortality was recorded in the groups treated with 2000, 3000, 4000, and 5000 mg/kg body weight of *X. aethiopica* fruit extract respectively. Similarly, in the second phase of medial lethal dose determination, no mortality was recorded in the group treated with 1200 mg/kg body weight of *X. aethiopica* fruit extract while 100% mortality was recorded in the groups treated with 1400, 1600, and 1800 mg/kg body weight of *X. aethiopica* fruit extract respectively as presented in table 1.

The median lethal dose (LD<sub>50</sub>) was calculated as geometrical means of the maximum dose producing 0% (a) and the minimum dose producing 100% mortality (b).

$$LD_{50} = \sqrt{ab}$$

Where a = 1200 mg/kg

b = 1400 mg/kg

LD<sub>50</sub>= 1296.15 mg/kg

#### 3.2: Effect of ethanolic extract of *Xylopi aethiopica* fruit on Sperm Parameters of Animals after 28 days of Treatment

Ethanolic extract of *Xylopi aethiopica* fruit was observed to significantly perturbed sperm parameters of animals after 28 days of treatment, as presented in Figs. 1-5. Sperm count and motility were significantly reduced by *Xylopi aethiopica* fruit in a dose-dependent manner when compared with those of the control group ( $P<0.05$ ). Administration of *Xylopi aethiopica* fruit increased sperm mortality and abnormality when compared with the control animals ( $P<0.05$ ). Seminal pH was decreased by ethanolic extract of *Xylopi aethiopica* fruit administration when compared with those in control animals ( $P<0.05$ ).

**Table 1: The Median Lethal Dose (LD<sub>50</sub>) of *Xylopi aethiopica* Fruit Extract**

Study (Animal)	Phase/ Dosage of Extract (mg/kg) b.w	No of Mice per Group	No. of Death Recorded	% Mortality
<b>PHASE ONE</b>				
I	1000	5	0	0
II	2000	5	5	100
III	3000	5	5	100
IV	4000	5	5	100
V	5000	5	5	100
<b>PHASE TWO</b>				
I	1200	5	0	0
II	1400	5	5	100
III	1600	5	5	100

IV	1800	5	5	100
V	2000	5	5	100

---

UNDER PEER REVIEW




Fig. 1: Effect of *X. aethiopica* fruit extract on Sperm Count of Animals after 28 days of Treatment

Each bar represents mean  $\pm$  SD with n = 6.




Fig. 2: Effect of *X. aethiopica* fruit extract on Sperm Motility of Animals after 28 days of Treatment

Each bar represents mean  $\pm$  SD with n = 6.






Fig. 3: Effect of *X. aethiopica* fruit extract on Sperm Mortality of Animals after 28 days of Treatment

Each bar represents mean  $\pm$  SD with  $n = 6$ .




Fig. 4: Effect of *X. aethiopica* fruit extract on Seminal pH of Animals after 28 days of Treatment

Each bar represents mean  $\pm$  SD with n = 6.




Fig. 5: Effect of *X. aethiopica* fruit extract on Sperm Abnormality of Animals after 28 days of Treatment

Each bar represents mean  $\pm$  SD with n = 6.

## DISCUSSION

The acute toxicity study of the plant extracts recorded 100% mortality at a dose of 1400 mg/kg bodyweight and above (table 1). This shows that the fruit of *Xylopi aethiopica* might

be highly toxic. The visible symptoms of toxicity seen in the animals were excitation, paw licking, increased respiratory rate, decreased motor activity, gasping and coma and death.

In this study, it could be clearly demonstrated that sperm count declined significantly ( $P<0.05$ ) when animals treated with *Xylopi aethiopica* fruits were compared with the control groups at all doses of treatment (Fig. 1). This could be an indication that *Xylopi aethiopica* fruit extract interfered with steroid hormone biosynthesis, which resulted in impaired spermatogenesis [38]. Disturbance in steroid hormone biosynthesis as well as spermatogenesis might affect the seminal quality of animals. This agreed with the decrease in sperm count of animals treated with *Xylopi aethiopica* fruits reported by Uyovwiese vwa *et al.* [39] and Abarikwu *et al.* [40] respectively. The decrease in sperm count observed in this study is dose-dependent. This indicates that consumption of *Xylopi aethiopica* fruit at high doses will lead to significant reduction in sperm count and thus infertility potential of male animals. Nwangwa, [41] and Eze [42] had independently reported the adverse impact of ethanolic fruit extract of *Xylopi aethiopica* on male sex organ of Wistar rats. The result of this present study is in consonance with their respective findings.

There was a noticeable ( $P<0.05$ ) decline in sperm motility of animals treated with ethanolic extract of *Xylopi aethiopica* fruit when compared with the control group at all doses in this study (Fig. 2). The reduced sperm motility observed in this study might be an indicator that *Xylopi aethiopica* fruit extract has the ability to reduce the ATPase activity in all tissue of the animals [43]. This causes suppression of energy metabolism. If ATPase activity is decreased, it could suppress the motility rate of sperm, as ATP is the main energy source of sperm and it is directly related to sperm motility [44]. Ogbuagu *et al.* [45] recently reported that ethanolic extract of *Xylopi aethiopica* fruit induced oxidative stress in Wistar rats. The inhibitory motility seen in the sperm of animals exposed to ethanolic extract of *Xylopi aethiopica* fruit in this study could also be associated with oxidative stress induced by the consumption of the extract. The sperm is particularly vulnerable to lipid peroxidation because of the molecular anatomy of its plasma membrane. The increased oxidative stress can damage the sperm membrane leading to reduced motility. This agreed with the work of Kalender and Yel [46] as well as that of Sachder and Davies [47]. The inhibitory motility seen in the sperm of rats administered ethanolic extract of *Xylopi aethiopica* fruit in this study is dose-dependent.

This indicates that consumption of *Xylopi aethiopica* fruit at high doses might cause a significant reduction in sperm motility and consequently infertility in male animals.

In this study, ethanolic extract of *Xylopi aethiopica* fruit was observed to increase the number of abnormal spermatozoa when compared with those of the control animals after 28 days of treatment (Fig. 3). Increased abnormality of spermatozoa in *Xylopi aethiopica* treated-animals might be as a result of damage of Sertoli cells [48]. For normal testicular function, Sertoli cells plays vital role in maintaining conducive environment for spermatogenesis. Damage to the Sertoli cells might affect the maturation process of spermatozoa, culminating in increased abnormality of sperms observed in this study. This result corresponded to the findings of Adienbo *et al.* [49] who observed a noticeable elevation in sperm abnormality in animals treated with *Xylopi aethiopica* fruit for 30 days.

The seminal pH shows the balance between the pH values of the various accessory gland secretions, majorly the alkaline seminal vesicular secretion and the acidic prostatic release. In this study, seminal pH was observed to decline when animals treated with fruit extract of *Xylopi aethiopica* were compared with the control animals after 28 days of treatment (Fig 4). This might be that *Xylopi aethiopica* fruit extract affects the normal pH range of treated animals. If the pH is decreased, the medium of seminal plasma becomes acidic which in turn makes sperms highly fragile, thus leading to higher rate of mortality [50].

A significant increase was observed in sperm mortality of animals treated with ethanolic extract of *Xylopi aethiopica* fruit when compared with the control group (Fig. 5). This might be attributed to the significant ( $P<0.05$ ) decrease in seminal pH in experimental animals. Low pH of epididymal fluid of bovine has been reported to result in increased rate of mortality of spermatozoa [43,51]. The exact mechanism by which *Xylopi aethiopica* fruit reduced sperm count is unknown, but it has been reported that it contain a compound called xylopic acid which possibly cross the blood testes barrier to adversely perturbed the seminiferous tubules of the testes.

The antifertility effect of ethanolic extract of

*Xylopia aethiopica* fruit observed in this study agreed with the findings of Nnodim *et al.* [52] who studied the impact of fruit extract of *Xylopia aethiopica* on the production of sperm and testicular oxidative status in male Wistar rats. Adienbo *et al.* [49] had previously reported the impairments in testicular function indices in male Wistar rats following the administration of *Xylopia aethiopica* fruit extract. Similarly, Nnodim *et al.* [52], reported the negative effect of *Xylopia aethiopica* fruit on male reproductive hormones. *Xylopia aethiopica* fruit had also been reported to produce negative impact on female reproductive hormones of animals [53]. In fact, Adienbo *et al.*, [49] has recommended *Xylopia aethiopica* fruit as a potent contraceptive.

## 5. CONCLUSION

The findings of this study revealed that ethanolic extract of *Xylopia aethiopica* fruit adversely perturbed sperm quality of Wistar rats. This might not automatically translate to same effect in human. However, men interested in child-bearing should minimize its consumption.

## CONSENT

It is not applicable.

## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCES

1. Agarwal A, Majzoub A, Esteves SC. Infertility, recurrent pregnancy loss and

- sperm DNA fragmentation, have we found the missing link. *Avi Harlev*. 2016;5:935-941.
2. Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HWG, Behre HM. World Health Organization reference values for human semen characteristics. *Human Reproduction*. 2010;16:231–245.
3. Speroff L, Fritz MA. Female infertility, Chapter 27, Clinical Gynecologic Endocrinology and Infertility, Eighth Edition. Lippincott Williams & Wilkins, syf. 2010;11:57.
4. Sartorius G, Nieschlag E. Paternal age and reproduction. *Hum. Reprod. Update*. 2010;16:65-79.
5. De Kester DM. Endocrinology of male infertility. *Brit Med Bullett*. 1979;35:187-192.
6. Amelar RD. Infertility in man. F. A Davis Company, Philadelphia, U.S.A. 1966;30-53.
7. Jarow JP. Endocrine causes of male infertility. *Urol Clin North Am*. 2003;30:83-90.
8. Gray EN. Testosterone, sexual function, and cognition. *J Clin Endocrinol Metab*. 2005;90(7):3838–3846.
9. Weinbauer GF, Nieschlag E. Gonadotropin control of testicular germ cell development. *Adv Exp Med Biol*. 1995;317:55-65.
10. Ali M, Al-Dosary UH, Ahmed M. Refaat, Saranya R. Babu, Abdul Rauf Choudhry. Y-str profiling of semen stain evidences of azoospermic individuals. *International Journal of Forensic Science & Pathology (IJFP)*, *International Journal of Forensic Science and Pathology*. 2015;3(11):210-214.
11. Jungwirth A, Diemer T, Dohle GR, Giwercman A, Kopa Z, Tournaye H, Krausz C. Guidelines for the investigation and treatment of male infertility. *Eur Urol*. 2012;61(1):159-163.
12. Montorsi F, Salonia A, Dehaf, Cestari A, Guazzoni G, Rigatti P, Steef C. Pharmacological management of erectile dysfunction, *British Journal of Urology*. 2003;8:211-216.
13. Shabsigh B, Anastasiadis AG. Erectile dysfunction, *Annual Review of Medicine*. 2003;45:153-168.
14. Porst H. Phosphodiesterase type-5 inhibitors a critical comparative analysis. *EAU update ser*. 2004;2:56-63.
15. Yakubu MT, Bilbis LS, Lawal M, Akanji MA. Effect of repeated administration of sildenafil citrate on selected enzyme activities of liver

- and kidney of male albino rats. *Nigerian Journal of Pure and Applied Sciences*. 2003;18:1395-4000.
16. Mendoza-Lujambio I, Nachtigall LB, Dowsing AT, Chase CD. Infertility in Male, 2008.
  17. Airaodion AI, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Ethanolic extract of *Garcinia kola* (Heckel) seed possesses antiparasmodial properties against *Plasmodium berghei*. *Asian Journal of Medical Principles and Clinical Practice*. 2020;3(1):33-39.
  18. Airaodion AI, Airaodion EO, Ekenjoku JA, Ogbuagu EO, Ogbuagu U. Antiparasmodial potency of ethanolic leaf extract of *Vernonia amygdalina* against *Plasmodium berghei* in infected swiss albino mice. *Asian Journal of Medical Principles and Clinical Practice*. 2019;2(2):1-8.
  19. Somova LI, Shode FO, Moodley K, Govender Y. Cardiovascular and diuretic activity of kaurene derivatives of *Xylopia aethiopica* and *Alepidea amatymbica*. *Journal of ethnopharmacology*. 2001;77(2-3):165-174.
  20. Asekun OT, Adeniyi BA. Antimicrobial and cytotoxic activities of the fruit essential oil of *Xylopia aethiopica* from Nigeria, *Fitoterapia*, 2004;75: 368-370.
  21. Koba K, Sanda K, Raynaud C, Guyon C, Chaumont JP, Nicod L. Chemical composition and *In vitro* cytotoxic activity of *Xylopia aethiopica* (Dun) A. Rich.(Annonaceae) fruit essential oil from Togo. *Journal of Essential Oil Research*. 2008;20(4):354-357.
  22. Woode E, Alhassan A, Abaidoo CS. Effect of xylopic acid on sex hormones and spermatogenesis in male rats. *Al Ameen j Med Sci.*; 2012;5(3): 28-29.
  23. Ogonnia S, Adekunle AA, Bosa MK, Enwuru VN. Evaluation of acute and subacute toxicity of *Alstonia congensis* Engler (Apocynaceae) bark and *Xylopia aethiopica* (Dunal) A. Rich (Annonaceae) fruits mixtures used in the treatment of diabetes. *African Journal of Biotechnology*. 2008;7(6):701-705.
  24. Esekhiagbe M, Agatemor MMU, Agatemor C. Phenolic content and antimicrobial potentials of *Xylopia aethiopica* and *Myristica argentea*. *Macedonian Journal of Chemistry and Chemical Engineering*. 2009;28(2):159-162.
  25. John-Dewole O, Agunbiade S, Alao O, Arojojoye O. Phytochemical and antimicrobial studies of extract of the fruit of *Xylopia aethiopica* for medicinal importance. *Journal of Biotechnology and Pharmaceutical Research*. 2012;29(6):118-122.
  26. Asekun O, Kunle O. The chemical constituents of the fruit essential oil of *Xylopia aethiopica* (Dunal) A. Rich from Nigeria. *Journal of Essential Oil Bearing Plants*. 2004;7(2):186-189.
  27. Nwaichi E, Igbinobaro O. Effects of some selected spices on some biochemical profile of Wistar albino rats. *American Journal of Environmental Engineering*. 2012;2(1):8-11.
  28. Ogbuagu EO, Nweke IN, Unekwe PC, Airaodion AI, Ogbuagu U. Weight gain reduction and hypoglycemic effects of *Xylopia aethiopica* fruit extract on Wistar rats. *International Journal of Research and Reports in Hematology*. 2020;5(3):1-8.
  29. Ogbuagu EO, Unekwe PC, Airaodion AI, Nweke IN, Ogbuagu U. Hypolipidemic propensity of ethanolic extract of *Xylopia aethiopica* fruit in Wistar rats. *Asian Journal of Research in Cardiovascular Diseases*. 2020;3(5):1-11.
  30. Ogbuagu EO, Unekwe PC, Airaodion AI, Nweke IN, Ogbuagu U. Hepatotoxic effect of *Xylopia aethiopica* fruit in Wistar rats. *International Research Journal of Gastroenterology and Hepatology*. 2021; 4(1):1-16
  31. Ogbuagu EO, Airaodion AI, Ogbuagu U, Nweke IN, Unekwe PC. Nephrotoxicity of Ethanolic Extract of *Xylopia aethiopica* Fruit in Wistar Rats. *International Journal of Advances in Nephrology Research*. 2021; 4(1): 1-16.
  32. Ogbuagu EO, Ogbuagu U, Unekwe PC, Nweke IN, Airaodion AI. Qualitative determination of the phytochemical composition of ethanolic extract of *Xylopia aethiopica* fruit. *Asian Journal of Medical Principles and Clinical Practice*. 2020;4(3):1-5
  33. Airaodion AI, Ekenjoku JA, Ogbuagu EO, Okoroukwu VN, Ogbuagu U. *Carica papaya* leaves might cause miscarriage. *Asian Research Journal of Gynaecology and Obstetrics*. 2019;2(2): 1-9.
  34. WHO. World Health Organization Laboratory Manual for the Examination of Human Semen and Semen-Cervical Mucus

- Interaction. 3rd edn. Cambridge University Press, Cambridge. 1992
35. Larsen L, Scheike T, Jensen TK, Bonde JP, Ernst E, Hjollund NH, Zhou Y, Skakkebaek NE, Giwercman A. Computer-assisted semen analysis parameters as predictors for fertility of men from the general population. The Danish First Pregnancy Planner Study Team. *Human Reproduction*. 2000;15(7):1562-1567.
  36. Airaodion AI, Ayanleke IA, Agunbiade AP, Ogbuagu EO, Airaodion EO, Ogbuagu U. Antifertility Propensity of *Jatropha curcas* Linn. Leaves on Male Wistar Rats. *International Journal of Research and Reports in Gynaecology*. 2020;3(2): 21-29.
  37. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Okoroukwu VN, Ogbuagu U, Airaodion EO. Antifertility effect of ethanolic leaf extract of *Carica papaya* in male Wistar rats. *Merit Research Journal of Medicine and Medical Science*. 2019;7(10):374-381.
  38. Lakhsman J, Changanma C. Antispermato-genic effect of *Vernonia amygdalina* seed extract on steroidogenesis in albino rats. *Int. Journal of Pharmacy and pharmaceutical science*. 2013;5 (1):23-35
  39. Uyovwiese-vwa AJ, Aloamaka PC, Avwioro OG. Effect of *Xylopi-aethiopica* plant extract on semen quality of the Sprague Dawley rats. *International Journal of Recent Scientific Research*. 2011;2(6):179 - 181
  40. Abarikwu SO, Ogunlaja A, Otuechere CA, Gideon OB. Effect of Ethanolic Extract from Seeds or Pods of *Xylopi-aethiopica* (Dunal) A. Rich (Annonaceae) on the Testicular Function of Adult Male Rats. *Indian Journal of Clinical Biochemistry*. 2017;32(4):420–428.
  41. Nwangwa EK. Antifertility Effects of Ethanolic Extract of *Xylopi-aethiopica* on Male Reproductive Organ of Wistar Rats. *American Journal of Medicine and Medical Sciences*. 2012;2(1): 12-15.
  42. Eze KN. Antifertility Effects of Ethanolic Extract of *Xylopi-aethiopica* on Male Reproductive Organ of Wistar Rats. *American Journal of Medicine and Medical Sciences*, 2012;2:12-15.
  43. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Okoroukwu VN, Ogbuagu U, Airaodion EO. Antifertility Effect of Ethanolic Leaf Extract of *Carica papaya* in Male Wistar Rats. *Merit Research Journal of Medicine and Medical Science*. 2019;3(9):121-129.
  44. Airaodion AI, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Consumption of coconut (*Cocos nucifera* L.) water improved fertility parameters in male Wistar rats. *Asian Journal of Pregnancy and Childbirth*. 2019;2(3):1-7.
  45. Ogbuagu EO, Ogbuagu U, Airaodion AI, Uche CL, Nweke IN, Unekwe PC. Effect of ethanolic Extract of *Xylopi-aethiopica* Fruit on Oxidative Stress Indices of Wistar Rats. *Asian Journal of Immunology*. 2022;3(4):1-11
  46. Kalender Y, Yell M. Doxorubicin hepatotoxic and hepatic free radical metabolism in rats. The effect of Vitamin E and Catechin. *Toxicology*. 2005;209:39-45
  47. Sachder S, Davies K. Molecular Biology. *Free Radicals Biol Med*. 2008;44:215-223.
  48. Manivannan B, Mittal R, Goyal S, Ansari AS, Lohiya NK. Sperm characteristics and ultrastructure of testes of rats after long-term treatment with the methanolic subfraction of *Vernonia amygdalina* seeds. *Asian J Androl*; 2009;11:583–599.
  49. Adienbo OM, Nwafor A, Dapper DV. Impairments in testicular function indices in male wistar rats: a possible mechanism for infertility induction by *Xylopi-aethiopica* fruit extract. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2015;4(1):71-75.
  50. Ekenjoku JA, Airaodion AI, Okoroukwu VN, Ogbuagu EO, Ogbuagu U. Oral administration of ethanolic extract of *Vernonia amygdalina* leaves might impact negatively on fertility in male Wistar rats. *Asian Journal of Medical Principles and Clinical Practice*. 2019;2(3):1-8.
  51. Ogbuagu EO, Airaodion AI. Tiger Nut (*Cyperus esculentus* L.) Boosts Fertility in Male Wistar Rats. *Asian Research Journal of Gynaecology and Obstetrics*. 2020;3(3): 8-18
  52. Nnodim JK, Nwanjo HU, Okolie NJ, Opara AU, Nwosu DC, Okoroiwu I, Dike J, Okorie H, Nwadike CN, Uduji HI. Effects of *Xylopi-aethiopica* Fruits on reproductive hormonal level in rats. *Global Journal of Medicinal Plant Research*, 2013;1(1): 29-31.
  53. Onuka AE, Okechukwu NC, Maxine KM. A Comparative Study between *Xylopi-aethiopica* Dried Fruit Extract and Ibuprofen Inhibiting Effects on Some Reproductive Hormones Irrespective of the Estrous Cycle. *International Journal of Complementary & Alternative Medicine*. 2017;8(5):1-6.