

Chromosomal Engineering through CRISPR–Cas Technology: A Way Forward

ABSTRACT

The breeding of crops is dependent on the potential to interrupt or maintain genetic links between characteristics, and the availability of genetic variability. CRISPR-Cas is a new genome-editing technique that has made it possible for breeders to introduce regulated and site-specific genetic diversity while simultaneously improving qualities with high efficacy. The existence of genomic linkage is a barrier in transferring desirable features among domesticated species from their wild counterparts. One way to address this issue is to create mutants with deficiencies in the meiotic recombination machinery, thereby enhancing global crossover frequencies between homologous parental chromosomes. Although this seemed to be a promising approach at first, thus far, no crossover frequencies could be enhanced in recombination-cold regions of the genome. Consequently, attempts have been made to induce site-specific DSBs in both somatic and meiotic plant cells by utilizing CRISPR–Cas techniques to achieve preset crossovers among homologs. Nonetheless, this method has not yielded significant heritable homologous crossings which were recombination-based. Lately, CRISPR–Cas has been used to achieve hereditary chromosomal rearrangements (CRs), including translocations and inversions, in plants. This method allows for the development of megabase CRs by DSB repair through non-homologous end-joining after insertion of DSBs in somatic plant cells. This technique may potentially make it possible to restructure genomes on a more global scale, culminating in the creation not just of synthetic plant chromosomes, but also that of new plant species.

Keywords: CRISPR-Cas; genetic linkage; crossover induction; chromosome engineering; reciprocal translocations; synthetic chromosomes; CRISPR-associated protein

1. INTRODUCTION

The world's population continues to grow at an exponential rate and it is becoming increasingly difficult to meet future food demands due to restrictions in food production resulting from limited farmland availability or unpredictable yields. The progression of innovative and rapid methods to enhance prevailing crops is therefore critical to fulfilling upcoming food requirements. Such enhancements include greater productivity, improved nutritional content, and better resistance to both biological and environmental stresses [1]. When using traditional breeding strategies, this procedure may need years to complete because of various factors, for example, the linkage-drag, i.e., linked inheritance of unfavorable genetic material coupled with desirable traits, or diminished genetic variation owing to the cultivation approach [2]. A case of linkage-drag in plants is the connection of tolerance to TMV with lower tobacco production [3,4].

A major shift in genetic enhancement has begun with the introduction of "clustered regularly interspaced short palindromic repeats–CRISPR-associated protein" (CRISPR–Cas) systems as a gene-editing tool [5–9]. This technology uses an endonuclease that can cause double-strand breaks (DSBs) practically anywhere in the genetic structure, and a synthetic sgRNA to direct the endonuclease towards its intended cut site [10]. DNA repair systems, including "homology-directed repair" (HDR) and "non-homologous end-joining" (NHEJ), are used to repair the DSBs induced by the endonuclease [11]. Although the NHEJ mechanism, which is prevalent in somatic plant cells, is prone to errors and, has been shown to result in minor deletions or insertions at the DSB site [11]. The NHEJ repairing may also result in CRs, including translocations, inversions, or deletions, when multiple DSBs are triggered concurrently [12,13]. Rearrangements of chromosomes that occur naturally in mammals are mostly linked to cancer and other genetic diseases [14–16]. Cancer may be caused by these rearrangements that alter the function of a tumor suppressor gene or trigger an oncogene [17]. Hemophilia A, EDMD, and Hunter syndrome are a few of the conditions that may be induced by CRs [18–21]. CRs, on the other hand, play an essential role in speciation and adaptation [22], particularly in crops. For example, heterozygous rearrangements may result in the creation of imbalanced gametes during the process of meiotic recombination, which can induce reproductive isolation or impair fertility [22,23]. These may also adversely affect gene expression regulation across the entire genome [24]. As the activity of meiotic recombination is hindered inside the rearranged region in heterozygotes, the genomic information stored in chromosomal translocations and inversions cannot be used for gene shuffles [25,26]. Thus, breeding operations can become complicated with the suppression of genetic exchange among two varieties with the occurrence of these arrangements. CRISPR–Cas has so far been used to make relatively minor genetic modifications in plants. Nevertheless, in numerous modern experiments in both *Zea mays* and *Arabidopsis thaliana*, the introduction of megabase-scale CRs has been reported [27–29]. In this study, we discuss chromosomal engineering, for the breeding of plants, as a unique technique for stabilizing or breaking genetic markers. Later on, we also explore the possibility for alternate strategies of generating recombination across homologous chromosomes.

2. MANIPULATING DNA REPAIR PATHWAYS TO BREAK GENETIC LINKAGES IN PLANT BREEDING

Allelic combinations in plants are generated by crossovers or the mutual interchange of genetic information, among homologous chromosomes during meiosis. These crossovers help for the combination of beneficial traits in elite crops while also allowing for the elimination of detrimental traits. Nonetheless, the majority of chromosomes do not exchange genetic information because distribution and crossover rates are extremely restricted and unregulated in natural processes [30]. As a result, the number of suitable recombination events is quite restricted, with linkage drag being mostly unavoidable. During meiotic division, the parental genomic sets are split throughout two successive nuclear divisions, with no interstitial replication, resulting in the generation of haploid reproductive cells [31]. Crossovers are required for the precise homologous chromosome segregations during the initial meiotic division because they allow the physical linkage of the coupled homologous chromosomes, known as bivalents [32]. This is guaranteed in all living things by the preservation of one mandatory crossover per chromosomal pair, also considered as the smallest crossover number required for recombination events to take place. Despite the fact that meiotic recombination is the primary source of genetic variation in plant progeny, crossover counts hardly surpass three per bivalent.

The production of DSBs by SPO11 homologs, which are highly conserved, sets the foundation for meiotic recombination [33]. A crossover's likelihood depends on the details of the succeeding meiotic DSB repair processes [31]. To facilitate the establishment of the D-loop, i.e., initial repair phase leading to the formation of the crossover, DSB terminals are removed and strand infiltration inside homologous sequences takes place. For the development of non-crossovers in which the invading strand gets lengthened, SDSA is an essential process. The D-loop is subsequently discarded and the DSB is spanned for repair assembly. Alternatively, in a second-end capture, the D-loop may be lengthened such that the rejected strand can anneal with the contrary side of the DSB. The creation of the second key intermediate, the double Holliday junction, is enabled as a result. This is the only process in crossover production that uses endonucleolytic cleavage to disintegrate the double Holliday junction. According to research, there are a variety of conditions in crops that might cause the balance of crossover origination to move in different directions. It seems that using this information by overexpressing crossover-promoting or crossover-limiting elements is a potential strategy for boosting genetic diversity in crops.




Fig. 1. Control of genetic exchange by targeted induction of crossovers and inversions

The protein-coding gene FANCM was the first factor discovered that limited crossover in plants. When *fancm* mutations were introduced into *A. thaliana*, its crossover frequency tripled [34]. The enzyme BLM homolog RECQ4 has been identified as another crossover inhibiting factor. *Arabidopsis* has two homologs, RECQ4A and RECQ4B, that are closely linked to each other [35,36]. The crossover frequency of the *recq4a recq4b* double mutant has been shown to increase by 6.2 times [36]. The SDSA-promoting destabilization of D-loops is facilitated by the actions of FANCM and RECQ4 [37,38]. As a result, it was hypothesized that the combined mutation of these enzymes represented an almost total lack of SDSA [39]. These factors suggest that every DSB results in a double Holiday junction-like structure that is resolved by resolvases, with crossovers and NCOs prevailing in equal amounts. Other elements also play a role in the regulation of crossover control in a plant system. The unique AAA-ATPase FIGL-1, in conjunction with its partner FLIP, is essential for the regulation of recombinases that initiates the strand invasion [40,41]. It has been shown that mutations taking place in *Arabidopsis figl1* enhance the number of crossovers in both hybrid and inbred lines. When utilizing these mutations in crops, the drawback is their sterility, as has been shown in tomato, pea, and rice mutants. This leads to a breeding dead end which is undesirable [42,43]. As an alternative to utilizing anti-crossover factor mutations, it is possible to overexpress crossover-promoting factors, for instance, the E3 ligase HEI10 [44]. *HEI10* artificial overexpression in *Arabidopsis* results in a spike in crossover frequencies, particularly in the domain of subtelomeric euchromatin [45]. The reported crossover increase, however, was again confined to the chromosomal arms when the results were paired with a *recq4a recq4b* mutant background, with the greatest impact found in subtelomeres and no effect observed in centromeres [46]. Surprisingly, the rice synaptonemal complex protein ZEP1 had a comparable effect on the number of crossings as HEI10 overexpression, suggesting that other factors may be at play [47]. As numerous mechanisms have an impact on crossover abundance, combining several approaches has the potential to further increase crossover frequency. As an example, the combined mutation of *recq4a recq4b* and *figl1*, or *recq4a recq4b* and *fancm*, in *Arabidopsis* inbred strains led to a tenfold rise in crossovers. Triple mutants of the *recq4a recq4b figl1* gene showed an increase of 7.8 times. Since a quadruple mutant with *fancm* showed no further increase, a threshold was hypothesized to have been achieved [48]. RECQ4 homologs were shown to have a major role in the frequency of crossovers in a 2018 research, demonstrating that these results might be applied to a variety of crops. Tomato, pea, and rice hybrids that had RECQ4 homologs inactivated had crossover frequencies that were increased by 2.7-, 4.7-, and 3.2-fold, respectively, indicating the possibility of employing mutants to speed up breeding procedures [43]. While modifying crossover routes to enhance meiotic recombination is highly promising, care should be used when modulating DNA repair components as the most effective technique for modifying *RECQ4* homologs is not without its drawbacks. RECQ4, a component of the RTR (RecQ/Top3/Rmi1) complex, not only plays a critical role in crossover regulation but also maintains somatic genome integrity [35,49]. Such as, in *A. thaliana*, when utilizing *recq4* mutants, it is possible that secondary mutations may accumulate and that the overall fitness of the plant would decline [50,51]. Similar findings were made in the case of FANCM, which plays a critical role in the preservation of genomic integrity in crops [52]. Finally, the methods for increasing crossover frequencies that have been presented have only been effective in increasing crossover frequency and had no effect over crossover positioning, resulting in recombination-depleted regions like the pericentromeric regions, unaffected by these changes.

3. CRISPR-CAS-MEDIATED CROSSOVER INDUCTION

Meiotic mutants, as described above, only increase crossover frequencies, but do not alter the distribution of crossovers which results in, a large portion of the genome that cannot be triggered for recombination [30]. Therefore, a simple technique that can target these areas directly during meiosis to induce DSBs and promote homologous recombination (HR) should be adopted [53]. In theory, two separate techniques can be employed to accomplish this objective: either utilizing a programmable DNA nuclease for DSB creation or utilizing its DNA-binding capabilities to lead the innate DSB-inducing mechanism to the desired target position in the genome.

SPO11, a naturally occurring regulator of meiotic DSB induction, was first revealed to be adaptable to genome-editing tools in a comprehensive yeast research by employing ZFNs, TALENs, and the CRISPR–Cas network as DNA binding proteins [54]. A considerable rise was seen in crossover frequency when these SPO11 fusions were directed to regions where DSB induction is normally low during meiosis. Nonetheless, natural constraints on the SPO11-mediated DSB induction mechanism were evident in parts of the genome where DSBs are inhibited during meiosis (e.g., the centromeric and pericentromeric regions). Recently, the same strategy was applied in *A. thaliana* [55] by fusing the SPO11 complex partner MTOPVIB, which is essential for SPO11-mediated DSB induction during meiosis, to a dead Cas9 and then guiding it to a crossover hotspot previously demonstrated to be accessible for crossover regulation. Similarly, the production of DSBs in yeast had been accomplished by directing SPO11 to hotspots. As a result, a rise in the crossover rate was generally anticipated. However, there was no discernible effect on crossover frequency or distribution from dCas9–MTOPVIB. Following these findings, it is possible that only recruiting endogenous DSB-inducing machinery will not be adequate to influence crossover incidence in plants.

The direct generation of meiotic site-specific DSBs has not yet been reported, however, Cas9-generated DSBs have effectively triggered recombination among homologous chromosomes in tomato somatic cells [56]. To induce an allele-specific DSB in the *PHYTOENE SYNTHASE 1* gene in hybrid plants, two genetically unique tomato accessions were used. The use of a fruit-color test allowed the researchers to distinguish between NHEJ and homologous recombination repair results. Additionally, somatic HR events were also discovered in the study of SNP redistribution data. Gene conversions and a possible crossover were among the results of the study. This work showed that CRISPR–Cas production of targeted DSBs may be used to control HR in plants, a previously unknown possibility.

Recently, a similar strategy was undertaken by the same researchers, who used hybrids of genetically diverse tomato accessions—which had heterozygous mutations in the *CAROTENOID ISOMERASE* gene—to serve as a selection system to identify HR-mediated somatic recombination [57]. According to the findings of this research, one crossover incidence with at least 1Mbp transitions in the two directions was reported.

Given the low frequency of HR in higher eukaryotic organisms, recombination across homologous chromosomes through NHEJ may be preferred to recombination facilitated by HR. For example, if the homologous chromosomes are subjected to DSB induction at the same time, NHEJ may result in the exchange of genetic material. HR has recently been shown in human cells by end-joining and targeting the *CD44* gene [58]. By using DSB induction and compound heterozygous mutations in the interval across these mutations, it was possible to identify future reciprocal recombination by the straightforward revival of genomic function. There was reciprocal recombination for both double-stranded breaks and nicks when they were directed towards each homologous chromosome. There was a ten time increase in the frequency of recombination when DSBs were induced compared to nicks, reaching frequencies of

roughly 0.1%. An interesting finding was the failure to induce recombination when double-strand breaks and nicks were steered to a single homologous chromosome. This suggests that Non-homologous end joining is preferred to homologous recombination when it comes to the focused recombination stimulation in higher eukaryotic organisms.

4. CRISPR–CAS-MEDIATED CHROMOSOME ENGINEERING IN YEAST AND MAMMALS

Among the first approaches for designing ways to disrupt genetic linkages is the global enhancement or site-specific induction of crossovers, as stated above. The CRISPR–Cas technology, on the other hand, might be utilized to reorganize and reorder chromosomes by the production of massive CRs. Using this method, significant progress has been achieved in various organisms, including mammals and yeast, and it may be possible to draw useful lessons from this experience for the use of identical strategies in crops.

For instance, in animal cells, Using the CRISPR–Cas approach, researchers have been able to mimic oncogenic CRs to understand cancer onset [59,60]. CRISPR–Cas has been used successfully by a number of researchers to reproduce chromosomal abnormalities in animal cells [60–62]. Several translocations, such as the t(11;22) and t(8;21), which induce acute myeloid leukemia and Ewing's sarcoma, respectively, have been successfully replicated [61]. A translocation on human chromosome 2 is another instance of a cancerous CR, which results in the creation of a defective fusion protein owing to the alignment of the two genes *ALK* and *EML4*. It is related to a specific form of cancer in the lungs and has been effectively induced in mice and human cells using the CRISPR–Cas system [63]. So far, the CRISPR–Cas system has been demonstrated to be useful for modifying animal chromosomes in multiple different ways.

CRISPR–Cas techniques for reshaping chromosomes and genomic characteristics in yeast have also made significant development over the past few years. In *Saccharomyces cerevisiae*, scientists were able to reduce its 16 chromosomes to just 1 and 2 in two separate experiments [64,65]. This was made possible by inducing a variety of CRISPR–Cas-mediated fusions and translocations. Since the wild type having 16 chromosomes could not be backcrossed, it was hypothesized that the strain having 2 chromosomes may be considered a separate species under the conventional biological species concept [64]. Moreover, cases of chromosomal shuffling in yeast have been documented as a result of the CRISPR–Cas-facilitated production of CRs [66, 67].

5. REVERSION OF NATURAL CHROMOSOMAL INVERSIONS CAN RESTORE GENETIC EXCHANGE IN RECOMBINATION-DEAD REGIONS IN PLANTS

The presence of CRs in a wide spectrum of crops has been shown [29,68–70]. It is possible to discover sequence variation across a crop plant by performing sequence assemblies on a chromosome-scale in order to capture the supragenome [68]. Greater than 5Mb-long inversion polymorphisms in genome arrangement of contemporary elite barley germplasm were found [68]. Thus, a significant amount of genomic material is now unavailable for breeding purposes, implying that an effective technology to generate or negate evolutionary-derived CRs in a targeted way would be very beneficial to crop breeders. A few years ago, the use of Cas9 from a gram-positive bacterium *Staphylococcus aureus* allowed researchers to demonstrate that precise inversions of approximately 18 kb can be generated in

Arabidopsis [71]. Following their initial findings, they went further another step and reverted an inversion based on the evolution of the popular heterochromatic knob on the short arm of chromosome 4 (hk4S), which would be carried by numerous accessions and has a 1.17Mb size [23,28,72]. In the 38 main transformants, the authors found 7 distinct inversion incidences. Crop plants may also be able to trigger or revert inversions in the Mb range, according to this study. Another goal of the researchers was to determine if the meiotic recombination process could be reintroduced in hybrids harboring the reversed knob in a heterozygous condition.




Fig. 2. Future perspective on chromosome engineering in plants

Previously, no evidence of gene flow between Columbia and Landsberg *erecta*, the former of which had a knob, has been found inside the rearranged region [73]. Schmidt et al. used a recombination assay based on SNP to see whether their theory was correct and discovered that hybrids' crossover rates could be restored [28]. This significant analysis implies that undoing evolutionary-derived CRs may liberate the genetic information contained inside the CR, allowing for meiotic crossover to occur. With regards to this work, the first crop plant inversion facilitated by CRISPR–Cas, in *Zea mays*, was recently procured [29]. This suggested that chromosomal engineering may be viable in other crops that are also sensitive to transformation similar to maize crops. In crop improvement, the restoration or production of meiotic chromosomal inversions may be utilized to replace crossovers in previously altered parts of the genome or to disrupt linkage groups by physical separation (Figure 1). However, inversions might also be utilized to maintain genetic links between desirable characteristics (Figure 1). These artificial inversions prevent HR between chromosomes during meiosis.

6. INDUCTION OF RECIPROCAL TRANSLOCATIONS IN PLANTS

Large translocations, like inversions, are widely observed in crop plants and may result in a decrease in meiotic recombination [25,69]. Semi-sterility may be caused by reciprocal translocations in plants, which can also be linked to male and female sterility [74]. As a result, the controlled production of these translocations will almost certainly become a valuable tool for crop breeders in the process of maintaining or breaking maintaining genetic linkages. A linkage may be established and disrupted by joining desirable genes from non-identical chromosomes inside the same chromosome and by physical separation, respectively (Figure 2). Recently, in plants, the first directed mutual translocations have been successfully induced. Translocations between *A. thaliana* 1 and 2 as well as 1 and 5 chromosomes were successfully achieved by Beying et al. The translocated segments were about 1 Mb and 0.5 Mb in size. There were translocation rates of about 2.5% and 3.75% in the natural background and in the NHEJ variant ku70, respectively, in discrete T2 lines. This hard evidence research gives reason to be optimistic that the similar would be feasible in crops in the near future.

7. POTENTIALS OF RESTRUCTURING THE PLANT CHROMOSOME

It's intriguing to imagine what more could be conceivable in the future now that it's possible to rearrange the order of genes inside and across chromosomes. However, one should remember that chromosomal engineering in crop plants is still very much in its infancy despite recent advances. CRs are very rare compared to the development of mutations based on DSBs. In fact, a quantitative examination of mutual translocations found that only 1 out of 10,000 cells had translocation events, meaning that very few repair responses result in the formation of CRs [27]. Wild-type host proteins owned by the cNHEJ pathway preserve the matching damaged DNA ends in immediate contact after a DSB to ensure their re-ligation, hence preventing the development of CRs. Because of this, it may be advantageous to induce CRs within a cNHEJ-genetic background in certain cases. In *A. Thaliana*, translocation and inversion rates are both increased, the former by approximately to five times, when the Ku70 enzyme, a key participant in cNHEJ, is absent [27]. DSB repair is taken over by the secondary NHEJ pathway when Ku70 is absent. This NHEJ pathway has no affinity for retaining the damaged DNA ends together. Hence the usage of DSB repairing mutants may increase the likelihood of acquiring CRs. Nonetheless, the disadvantage of such a strategy is that a general repair defect may lead to additional, undesirable modifications in the genome. Furthermore, enhancing CR frequencies may also be achieved by using sgRNAs and Cas nucleases that have been designed for effective DSB generation in the relevant plant species [75,76]. Moreover, detecting unusual CRs requires an effective screening protocol. *A. thaliana* has successfully used a mass screening methodology that was initially designed to find infrequent gene targeting occasions [77].

A variety of CRs have occurred throughout evolution and could perhaps set a standard for the types of genomic modifications that CRISPR–Cas-mediated chromosomal engineering may be able to achieve. CRs are common during polyploidization mechanisms that result in dysploid changes, which in turn lead to a diploid chromosome pair [78]. The decrease in the number of chromosomes should be conceivable, either by end-to-end translocation or nested insertions, both of which have been recorded in natural CRs. Polyploids like potatoes and wheat might be suitable candidates for chromosomal editing techniques like these. The recent in-vivo observation of centromere relocation in maize following gamma irradiation further supports the theory that centromere repositioning arises regularly throughout plant genomic evolution [78,79]. It may be conceivable in the near future to imitate nature by altering centromere positions

and chromosomal numbers in the same way that it has been feasible to cause translocations and inversions, which take place with great frequency upon an evolutionary level (Figure 2). Nevertheless, it is important to remember that if numerous DSBs are induced concurrently, additional unexpected genetic alterations may result [80]. Along with present techniques, this would enable fundamental issues to be answered about how gene expression and chromatin state are affected by different locations of a chromosome. Additionally, chromosomal engineering may enable the reconstruction of the chromosome sets of contemporary plant species' ancestors.

8. THE CREATION OF SYNTHETIC CHROMOSOMES AND NEW PLANT SPECIES

Since the turn of the millennium, a unique method of modifying genomes has surfaced, i.e., the fabrication of artificial chromosomes. Owing to their tiny size, the earliest manufactured chromosomes were viral in origin. Following this achievement, artificial bacterial chromosomes of several strains and artificial yeast chromosomes were constructed [81]. Although theoretically feasible, the introduction of completely synthesized chromosomes from other creatures into crop plants has been hampered by a number of technological challenges, such as limits throughout the transformation procedure. Conventional plant transformation techniques, including *Agrobacterium* and biolistics, can only transfer so much DNA at a time until they run out of efficiency. The ability of an *Agrobacterium* to transfer is reliant on the normal *E. coli* vector capacity [82] of around 150 kb. Biolistic transformation, on the other hand, is capable of transferring up to 1,050 kb, although it may cause substantial degradation to the host genome and transgene and is typically incapable of delivering intact molecules exceeding a few kilobases [83,84]. Furthermore, synthetic chromosomes require a certain size in order to accommodate at least a single crossover throughout meiosis. In doing so, we may avoid the early separation of sister chromatids, which might lead to deterioration and the removal of genetic information [82]. Given the existing limitations of constructing synthetic crop plant chromosomes from the beginning, the next phase in the synthesis of these synthetic chromosomes will most likely be accomplished using CRISPR–Cas-mediated chromosomal engineering, in conjunction with the insertion of shorter lengths of artificial DNA.

9. CONCLUSION

Novel plant species, on the other hand, will be possible much sooner. For the first time, researchers were successfully able to engineer *Saccharomyces cerevisiae* cells with a single- and two-chromosome using CRISPR–Cas technology [64,65]. CRISPR–Cas-mediated chromosomal reorganization of crop plant chromosomes has made considerable progress, and it may be conceivable to use the same method to create a new species of plants via the process of reproductive isolation. Engineered plants may become reproductively isolated from their wild-type parent by the decrease or combination of distinct forms of CRs. If this is the case, the modified line might be considered a new species of plant. This might be an intriguing way to avoid agricultural plants from crossing with wild counterparts in an unintended way. Using chromosomal engineering in crop plants has opened new doors in the domain of agricultural development and will enable scientists better understand how genetics and evolution work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bailey-Serres J, Parker JE, Ainsworth EA, Oldroyd GE, Schroeder JI. Genetic strategies for improving crop yields. *Nature*. 2019;575(7781):109-118.
2. Wolter F, Schindele P, Puchta H. Plant breeding at the speed of light: the power of CRISPR/Cas to generate directed genetic diversity at multiple sites. *BMC Plant Biol*. 2019;19(1):1-8.
3. Lewis RS, Rose C. Agronomic Performance of Tobacco Mosaic Virus-Resistant Tobacco Lines and Hybrids Possessing the Resistance Gene N Introgressed on Different Chromosomes. *Crop Sci*. 2010;50(4):1339-1347.
4. Li J, Chitwood J, Menda N, Mueller L, Hutton SF. Linkage between the I-3 gene for resistance to Fusarium wilt race 3 and increased sensitivity to bacterial spot in tomato. *Teor. Appl. Genet*. 2018;131(1):145-155.
5. Zhang Y, Pribil M, Palmgren M, Gao C. A CRISPR way for accelerating improvement of food crops. *Nat. Food*. 2020;1(4):200-205.
6. Zaidi SS, Mahas A, Vanderschuren H, Mahfouz MM. Engineering crops of the future: CRISPR approaches to develop climate-resilient and disease-resistant plants. *Genome Biol*. 2020;21(1):1-9.
7. Schindele A, Dorn A, Puchta H. CRISPR/Cas brings plant biology and breeding into the fast lane. *Curr. Opin. Biotechnol*. 2020;61:7-14.
8. Atkins PA, Voytas DF. Overcoming bottlenecks in plant gene editing. *Curr. Opin. Plant Biol*. 2020;54:79-84.
9. Zhang Y, Malzahn AA, Sretenovic S, Qi Y. The emerging and uncultivated potential of CRISPR technology in plant science. *Nat. Plants*. 2019;5(8):778-794.
10. Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*. 2012;337(6096):816-821.
11. Puchta H. The repair of double-strand breaks in plants: mechanisms and consequences for genome evolution. *J. Exp. Bot*. 2005;56(409):1-4.
12. Schmidt C, Schindele P, Puchta H. From gene editing to genome engineering: restructuring plant chromosomes via CRISPR/Cas. *aBIOTECH*. 2020;1(1):21-31.
13. Rönspies M, Schindele P, Puchta H. CRISPR/Cas-mediated chromosome engineering: opening up a new avenue for plant breeding. *J. Exp. Bot*. 2021;72(2):177-183.
14. Thompson SL, Compton DA. Chromosomes and cancer cells. *Chromosome Res*. 2011;19(3):433-444.
15. Park CY, Sung JJ, Kim DW. Genome editing of structural variations: modeling and gene correction. *Trends Biotechnol*. 2016;34(7):548-561.
16. Brunet E, Jasin M. Induction of chromosomal translocations with CRISPR-Cas9 and other nucleases: understanding the repair mechanisms that give rise to translocations. *Adv. Exp. Med. Biol*. 2018;33(6):15-25.
17. Hasty P, Montagna C. Chromosomal rearrangements in cancer: detection and potential causal mechanisms. *Mol. Cell. Oncol*. 2014;1(1):29-39.

18. Lakich D, Kazazian HH, Antonarakis SE, Gitschier J. Inversions disrupting the factor VIII gene are a common cause of severe haemophilia A. *Nat. Genet.* 1993;5(3):236-241.
19. Bondeson ML, Dahl N, Malmgren H, Kleijer WJ, Tønnesen T, Carlberg BM, Pettersson U. Inversion of the IDS gene resulting from recombination with IDS-related sequences in a common cause of the Hunter syndrome. *Hum. Mol. Genet.* 1995;4(4):615-621.
20. Small K, Iber J, Warren ST. Emerin deletion reveals a common X-chromosome inversion mediated by inverted repeats. *Nat. Genet.* 1997;16(1):96-99.
21. Park CY, Lee DR, Sung JJ, Kim DW. Genome-editing technologies for gene correction of hemophilia. *Hum. Genet.* 2016;135(9):977-981.
22. Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and speciation. *Genetics.* 2006;173(1):419-434.
23. Frasz P, Linc G, Lee CR, Aflitos SA, Lasky JR, Toomajian C, Ali H, Peters J, van Dam P, Ji X, Kuzak M. Molecular, genetic and evolutionary analysis of a paracentric inversion in *Arabidopsis thaliana*. *Plant J.* 2016;88(2):159-178.
24. Harewood L, Fraser P. The impact of chromosomal rearrangements on regulation of gene expression. *Hum. Mol. Genet.* 2014;23(1):76-82.
25. Martin G, Baurens FC, Hervouet C, Salmon F, Delos JM, Labadie K, Perdereau A, Mournet P, Blois L, Dupouy M, Carreel F. Chromosome reciprocal translocations have accompanied subspecies evolution in bananas. *Plant J.* 2020;104(6):1698-1711.
26. Wellenreuther M, Bernatchez L. Eco-evolutionary genomics of chromosomal inversions. *Trends Ecol. Evol.* 2018;33(6):427-440.
27. Beying N, Schmidt C, Pacher M, Houben A, Puchta H. CRISPR–Cas9-mediated induction of heritable chromosomal translocations in *Arabidopsis*. *Nat. Plants.* 2020;6(6):638-645.
28. Schmidt C, Frasz P, Rönspies M, Dreissig S, Fuchs J, Heckmann S, Houben A, Puchta H. Changing local recombination patterns in *Arabidopsis* by CRISPR/Cas mediated chromosome engineering. *Nat. Commun.* 2020;11(1):1-8.
29. Schwartz C, Lenderts B, Feigenbutz L, Barone P, Llaca V, Fengler K, Svitashv S. CRISPR–Cas9-mediated 75.5-Mb inversion in maize. *Nat. Plants.* 2020;6(12):1427-1431.
30. Taagen E, Bogdanove AJ, Sorrells ME. Counting on crossovers: controlled recombination for plant breeding. *Trends Plant Sci.* 2020;25(5):455-465.
31. Wang Y, Copenhaver GP. Meiotic recombination: mixing it up in plants. *Annu. Rev. Plant Biol.* 2018;69:577-609.
32. Lambing C, Franklin FC, Wang CJ. Understanding and manipulating meiotic recombination in plants. *Plant Physiol.* 2017;173(3):1530-1542.
33. Bergerat A, de Massy B, Gadelle D, Varoutas PC, Nicolas A, Forterre P. An atypical topoisomerase II from Archaea with implications for meiotic recombination. *Nature.* 1997;386(6623):414-417.
34. Crismani W, Girard C, Froger N, Pradillo M, Santos JL, Chelysheva L, Copenhaver GP, Horlow C, Mercier R. FANCM limits meiotic crossovers. *Science.* 2012;336(6088):1588-1590.
35. Hartung F, Suer S, Puchta H. Two closely related RecQ helicases have antagonistic roles in homologous recombination and DNA repair in *Arabidopsis thaliana*. *Proc. Natl Acad. Sci. USA.* 2007;104(47):18836-18841.
36. Séguéla-Arnaud M, Crismani W, Larchevêque C, Mazel J, Froger N, Choinard S, Lemhemdi A, Macaisne N, Van Leene J, Gevaert K, De Jaeger G. Multiple mechanisms limit meiotic

crossovers: TOP3 α and two BLM homologs antagonize crossovers in parallel to FANCM. *Proc. Natl Acad. Sci. USA*. 2015;112(15):4713-4718.

37. Dorn A, Rohrig S, Papp K, Schropfer S, Hartung F, Knoll A, Puchta H. The topoisomerase 3 α zinc-finger domain T1 of *Arabidopsis thaliana* is required for targeting the enzyme activity to Holliday junction-like DNA repair intermediates. *PLoS Genet*. 2018;14(9):100-110.
38. Seguela-Arnaud M, Choinard S, Larcheveque C, Girard C, Froger N, Crismani W, Mercier R. RMI1 and TOP3 α limit meiotic CO formation through their C-terminal domains. *Nucleic Acids Res*. 2017;45(4):1860-1871.
39. de Maagd RA, Loonen A, Chouaref J, Pelé A, Meijer-Dekens F, Fransz P, Bai Y. CRISPR/Cas inactivation of RECQ 4 increases homeologous crossovers in an interspecific tomato hybrid. *Plant Biotechnol J*. 2020;18(3):805-813.
40. Girard C, Chelysheva L, Choinard S, Froger N, Macaisne N, Lemhemdi A, Mazel J, Crismani W, Mercier R. Correction: AAA-ATPase FIDGETIN-LIKE 1 and helicase FANCM antagonize meiotic crossovers by distinct mechanisms. *PLoS Genet*. 2015;11(9):129-141.
41. Fernandes JB, Duhamel M, Seguela-Arnaud M, Froger N, Girard C, Choinard S, Solier V, De Winne N, De Jaeger G, Gevaert K, Andrey P. FIGL1 and its novel partner FLIP form a conserved complex that regulates homologous recombination. *PLoS Genet*. 2018;14(4):89-92.
42. Zhang P, Zhang Y, Sun L, Sinumporn S, Yang Z, Sun B, Xuan D, Li Z, Yu P, Wu W, Wang K. The rice AAA-ATPase OsFIGNL1 is essential for male meiosis. *Front. Plant Sci*. 2017;8:1639-1652.
43. Mieulet D, Aubert G, Bres C, Klein A, Droc G, Vieille E, Rond-Coissieux C, Sanchez M, Dalmais M, Mauxion JP, Rothan C. Unleashing meiotic crossovers in crops. *Nat. Plants*. 2018;4(12):1010-1016.
44. Pyatnitskaya A, Borde V, De Muyt A. Crossing and zipping: molecular duties of the ZMM proteins in meiosis. *Chromosoma*. 2019;128(3):181-198.
45. Ziolkowski PA, Underwood CJ, Lambing C, Martinez-Garcia M, Lawrence EJ, Ziolkowska L, Griffin C, Choi K, Franklin FC, Martienssen RA, Henderson IR. Natural variation and dosage of the HEI10 meiotic E3 ligase control *Arabidopsis* crossover recombination. *Genes Dev*. 2017;31(3):306-317.
46. Serra H, Lambing C, Griffin CH, Topp SD, Nageswaran DC, Underwood CJ, Ziolkowski PA, Séguéla-Arnaud M, Fernandes JB, Mercier R, Henderson IR. Massive crossover elevation via combination of HEI10 and *recq4a recq4b* during *Arabidopsis* meiosis. *Proc. Natl Acad. Sci. USA*. 2018;115(10):2437-2442.
47. Wang K, Wang C, Liu Q, Liu W, Fu Y. Increasing the genetic recombination frequency by partial loss of function of the synaptonemal complex in rice. *Mol. Plant*. 2015;8(8):1295-1298.
48. Fernandes JB, Séguéla-Arnaud M, Larchevêque C, Lloyd AH, Mercier R. Unleashing meiotic crossovers in hybrid plants. *Proc. Natl Acad. Sci. USA*. 2018;115(10):2431-2436.
49. Hartung F, Suer S, Knoll A, Wurz-Wildersinn R, Puchta H. Topoisomerase 3 α and RMI1 suppress somatic crossovers and are essential for resolution of meiotic recombination intermediates in *Arabidopsis thaliana*. *PLoS Genet*. 2008;4(12):100-115.
50. Mannuss A, Dukowic-Schulze S, Suer S, Hartung F, Pacher M, Puchta H. RAD5A, RECQ4A, and MUS81 have specific functions in homologous recombination and define different pathways of DNA repair in *Arabidopsis thaliana*. *Plant Cell*. 2010;22(10):3318-3330.
51. Schröpfer S, Kobbe D, Hartung F, Knoll A, Puchta H. Defining the roles of the N-terminal region and the helicase activity of RECQ4A in DNA repair and homologous recombination in *Arabidopsis*. *Nucleic Acids Res*. 2014;42(3):1684-1697.

52. Knoll A, Higgins JD, Seeliger K, Reha SJ, Dangel NJ, Bauknecht M, Schröpfer S, Franklin FC, Puchta H. The Fanconi anemia ortholog FANCM ensures ordered homologous recombination in both somatic and meiotic cells in *Arabidopsis*. *Plant Cell*. 2012;24(4):1448-1464.
53. Peciña A, Smith KN, Mézard C, Murakami H, Ohta K, Nicolas A. Targeted stimulation of meiotic recombination. *Cell*. 2002;111(2):173-184.
54. Sarno R, Vicq Y, Uematsu N, Luka M, Lapierre C, Carroll D, Bastianelli G, Serero A, Nicolas A. Programming sites of meiotic crossovers using Spo11 fusion proteins. *Nucleic Acids Res*. 2017;56(4):78-92.
55. Yelina NE, Gonzalez-Jorge S, Hirsz D, Yang Z, Henderson IR. CRISPR targeting of MEIOTIC-TOPOISOMERASE VIB-dCas9 to a recombination hotspot is insufficient to increase crossover frequency in *Arabidopsis*. *Plant Cell*. 2021;34(9):516-526.
56. Filler Hayut S, Melamed Bessudo C, Levy AA. Targeted recombination between homologous chromosomes for precise breeding in tomato. *Nat. Commun*. 2017;8(1):1-9.
57. Ben Shlush I, Samach A, Melamed-Bessudo C, Ben-Tov D, Dahan-Meir T, Filler-Hayut S, Levy AA. CRISPR/Cas9 induced somatic recombination at the CRTISO locus in tomato. *Genes*. 2021;12(1):59-66.
58. Davis L, Khoo KJ, Zhang Y, Maizels N. POLQ suppresses interhomolog recombination and loss of heterozygosity at targeted DNA breaks. *Proc. Natl Acad. Sci. USA*. 2020;117(37):22900-22909.
59. Cheong TC, Blasco RB, Chiarle R. The CRISPR/Cas9 system as a tool to engineer chromosomal translocation in vivo. *Adv. Exp. Med. Biol*. 2018:39-48.
60. Torres-Ruiz R, Martinez-Lage M, Martin MC, Garcia A, Bueno C, Castaño J, Ramirez JC, Menendez P, Cigudosa JC, Rodriguez-Perales S. Efficient recreation of t (11; 22) EWSR1-FLI1+ in human stem cells using CRISPR/Cas9. *Stem Cell Rep*. 2017;8(5):1408-1420.
61. Torres RA, Martin MC, Garcia A, Cigudosa JC, Ramirez JC, Rodriguez-Perales S. Engineering human tumour-associated chromosomal translocations with the RNA-guided CRISPR–Cas9 system. *Nat. Commun*. 2014;5(1):1-8.
62. Lagutina IV, Valentine V, Picchione F, Harwood F, Valentine MB, Villarejo-Balcells B, Carvajal JJ, Grosveld GC. Modeling of the human alveolar rhabdomyosarcoma Pax3-Foxo1 chromosome translocation in mouse myoblasts using CRISPR-Cas9 nuclease. *PLoS Genet*. 2015;11(2):25-39.
63. Choi PS, Meyerson M. Targeted genomic rearrangements using CRISPR/Cas technology. *Nat. Commun*. 2014;5(1):1-6.
64. Luo J, Sun X, Cormack BP, Boeke JD. Karyotype engineering by chromosome fusion leads to reproductive isolation in yeast. *Nature*. 2018;560(7718):392-396.
65. Shao Y, Lu N, Wu Z, Cai C, Wang S, Zhang LL, Zhou F, Xiao S, Liu L, Zeng X, Zheng H. Creating a functional single-chromosome yeast. *Nature*. 2018;560(7718):331-335.
66. Fleiss A, O'Donnell S, Fournier T, Lu W, Agier N, Delmas S, Schacherer J, Fischer G. Reshuffling yeast chromosomes with CRISPR/Cas9. *PLoS Genet*. 2019;15(8):11-25.
67. Yadav V, Sun S, Coelho MA, Heitman J. Centromere scission drives chromosome shuffling and reproductive isolation. *Proc. Natl Acad. Sci. USA*. 2020;117(14):7917-7928.
68. Jayakodi M, Padmarasu S, Haberer G, Bonthala VS, Gundlach H, Monat C, Lux T, Kamal N, Lang D, Himmelbach A, Ens J. The barley pan-genome reveals the hidden legacy of mutation breeding. *Nature*. 2020;588(7837):284-289.

69. Walkowiak S, Gao L, Monat C, Haberer G, Kassa MT, Brinton J, Ramirez-Gonzalez RH, Kolodziej MC, Delorean E, Thambugala D, Klymiuk V. Multiple wheat genomes reveal global variation in modern breeding. *Nature*. 2020;588(7837):277-283.
70. Crow T, Ta J, Nojoomi S, Aguilar-Rangel MR, Torres Rodríguez JV, Gates D, Rellán-Alvarez R, Sawers R, Runcie D. Gene regulatory effects of a large chromosomal inversion in highland maize. *PLoS Genet*. 2020;16(12):277-285.
71. Schmidt C, Pacher M, Puchta H. Efficient induction of heritable inversions in plant genomes using the CRISPR/Cas system. *Plant J*. 2019;98(4):577-589.
72. Fransz PF, Armstrong S, de Jong JH, Parnell LD, van Drunen C, Dean C, Zabel P, Bisseling T, Jones GH. Integrated cytogenetic map of chromosome arm 4S of *A. thaliana*: structural organization of heterochromatic knob and centromere region. *Cell*. 2000;100(3):367-376.
73. Drouaud J, Camilleri C, Bourguignon PY, Canaguier A, Bérard A, Vezon D, Giancola S, Brunel D, Colot V, Prum B, Quesneville H. Variation in crossing-over rates across chromosome 4 of *Arabidopsis thaliana* reveals the presence of meiotic recombination “hot spots”. *Genome Res*. 2006;16(1):106-114.
74. Zhang Y, Cheng Z, Ma J, Xian F, Zhang X. Characteristics of a novel male–female sterile watermelon (*Citrullus lanatus*) mutant. *Sci. Horticulturae*. 2012;140:107-114.
75. Steinert J, Schiml S, Fauser F, Puchta H. Highly efficient heritable plant genome engineering using Cas9 orthologues from *Streptococcus thermophilus* and *Staphylococcus aureus*. *Plant J*. 2015;84(6):1295-1305.
76. Schindele P, Puchta H. Engineering CRISPR/LbCas12a for highly efficient, temperature-tolerant plant gene editing. *Plant Biotechnol. J*. 2020;18(5):1118.
77. Wolter F, Klemm J, Puchta H. Efficient in planta gene targeting in Arabidopsis using egg cell-specific expression of the Cas9 nuclease of *Staphylococcus aureus*. *Plant J*. 2018;94(4):735-746.
78. Mandáková T, Lysak MA. Post-polyploid diploidization and diversification through dysploid changes. *Curr. Opin. Plant Biol*. 2018;42:55-65.
79. Liu Y, Su H, Zhang J, Shi L, Liu Y, Zhang B, Bai H, Liang S, Gao Z, Birchler JA, Han F. Rapid birth or death of centromeres on fragmented chromosomes in maize. *Plant Cell*. 2020;32(10):3113-3123.
80. Wimmer E, Mueller S, Tumpey TM, Taubenberger JK. Synthetic viruses: a new opportunity to understand and prevent viral disease. *Nat. Biotechnol*. 2009;27(12):1163-1172.
81. Coradini AL, Hull CB, Ehrenreich IM. Building genomes to understand biology. *Nat. Commun*. 2020;11(1):1-1.
82. Dawe RK. Charting the path to fully synthetic plant chromosomes. *Exp. Cell. Res*. 2020;390(1):111-126.
83. Liu J, Nannas NJ, Fu FF, Shi J, Aspinwall B, Parrott WA, Dawe RK. Genome-scale sequence disruption following biolistic transformation in rice and maize. *Plant Cell*. 2019;31(2):368-383.
84. Zhang H, Phan BH, Wang K, Artelt BJ, Jiang J, Parrott WA, Dawe RK. Stable integration of an engineered megabase repeat array into the maize genome. *Plant J*. 2012;70(2):357-365.

