

Original Research Article

OCT Macular Changes in Amblyopic Children

Abstract

Background: Amblyopia is defined as a decrease in best-corrected visual acuity in one or both eyes that is not solely due to a defect in the structure of the eye. This study aimed to investigate the OCT changes occurring in the amblyopic eye retina compared to normal eye.

Methods: This prospective study was carried out on 30 cases diagnosed with amblyopia using the better eye as a control. Patients were subjected to Assessment (visual acuity was measured by Landolt's chart. Refraction error measured by autorefractometer) and macular analysis.

Results: CMT was statistically insignificant difference between amblyopic and non-amblyopic eyes. The difference in AMT between amblyopic and non-amblyopic eyes was statistically significant ($p = 0.001$). BCVA in different ages before and after treatment was statistically significant difference ($P=0.001$). The various types of refractive errors of the studied eyes regarding BCVA, before and after treatment was statistically significant difference ($P=0.001$).

Conclusions: CMT and AMT of the amblyopic eye more than the sound eye and after 6 months of patching, there was a decrease in the CMT and AMT of the amblyopic eye.

Keywords: OCT, Macular Changes, Amblyopic Children.

Introduction

Amblyopia is defined as a decrease in best-corrected visual acuity in one or both eyes that is not solely due to a defect in the eye structure. The prevalence of amblyopia worldwide is approximately 1%-5%. Amblyopia may be bilateral and this is called isoametropic amblyopia or unilateral commonly in hyperopia and astigmatism and less common in myopia. Amblyopia is a condition in which the normal development of the cortical visual pathway is disrupted during childhood. It is defined as a discrepancy of two or more lines in best-corrected visual acuity between the eyes clinically ^[1].

Amblyopia lowers visual acuity and contrast sensitivity due to unusual visual processing in the primary visual cortex (V1). Amblyopia can be caused by a variety of ocular disorders, including uncorrected refractive problems, strabismus, and obstruction of the central visual axis. During the "critical time" of brain plasticity, amblyopia is treated by addressing the underlying cause and penalising the healthy eye. Treatment was traditionally suggested for children until they were 8–9 years old. Recent research suggests, nevertheless, that therapy of older people may be successful since the visual system's plasticity may stretch into adulthood. Furthermore, in children and adults with amblyopia, medication that addresses inhibitory interactions within the visual cortex may enhance both binocular and monocular visual performance ^[2]. Amblyopia can be caused by uncorrected refractive defects, strabismus, visual axis obstruction, or an integration of these factors ^[3]. The evidence for amblyopic retinal changes is still inconclusive and debatable. In the amblyopic eye, retinal ganglion cells are disturbed, which results in greater thickness of the retinal nerve fiber layer (RNFL). Disruption of the macula's physiological development, such as the relocation of Henle's fibers away from the foveola, and a decline in foveal cone diameter, were also believed to cause an increase in foveal thickness ^[4]. Previous investigations of the retina utilizing optical coherence tomography (OCT) imaging have yielded contradictory findings.

In amblyopia, researchers began to measure the thickness of the fovea's layers. Till now, the studies have produced contradictory findings. They discovered that in anisohypermetropic amblyopic eyes, the length of the outer segment (OS) layer of a photoreceptor in the fovea was substantially thinner than in companion eyes ^[6]. They also noticed reduced length in foveal cones and a thicker overall fovea in amblyopia eyes ^[7]. In both the inner plexiform layer and the ganglion cell layer, they discovered a significant difference between amblyopic and fellow eyes, but not in the outer segment layer ^[8].

The eyes of amblyopic and non-amblyopic showed insignificant difference, however there was a substantial difference between amblyopic and control eyes ^[9]. These inconsistencies could be related to differences in ethnicity, patient age, amblyopia severity, or assessment procedures for foveal layer thickness ^[10]. The goal of this investigation was to make a comparison in the OCT alterations in the retina of an amblyopic eye to a normal eye and measure each retinal layer thickness in various parts of the fovea (within a 0.5mm radius of the foveal center) in the children eyes diagnosed with unilateral amblyopia with those from the age-matched control eyes.

Patients and Methods

This prospective study was conducted on 30 cases diagnosed with amblyopia using the better eye as a control, age 4-6 years and difference in best-corrected visual acuity of 2 or more lines of acuity between the eyes. All participants signed an informed consent form. Children with sensory lesions as: (Congenital cataract - Corneal opacity), previous squint surgery, uncooperative patient: (young age and mentally retarded), previous surgical intervention and previous amblyopia treatment were excluded.

Methods: Assessment (visual acuity was measured by Landolt's chart. Refraction error measured by autorefractometer using cycloplegic refraction test. cycloplegia induced by instillation of cyclopentolate 1% eye drops. Detailed ophthalmologic and fundoscopic

examination by slit lamp biomicroscopy. Patching the sound eye of children 4-6 year according to severity of amblyopia if mild 2hours per day and if sever 6 hours per day for 1 month and follow up for 6 months by OCT. Macular thickness will be measured by SD-OCT. The Spectral domain optical coherence Tomography CIRRUS™ (Cirrus-4000 OCT, ZEISS corporation, Carl-Zeiss, Germany) was used to measure the macular thickness. CIRRUS™ normative databases matched the average anatomic age and ethnical diversity values and obtained clearance from the US food and drug administration (FDA)).

Macular analysis: The print shows a line scanning ophthalmoscope (LSO) fundus image with an ILM-RPE retinal thickness map overlay. With Fovea Finder, the ETDRS grid is automatically centered on the fovea. The values of retinal thickness, measured in microns from the ILM to the RPE, are compared to normative data. Slice navigator allows you to see a selected point on an LSO picture, an OCT fundus image, a retinal thickness map, layer maps, and an OCT image all at the same time. The Fovea Finder allows for exact ETDRS grid placement. The macular thickness map is a topographical numerical and colored coded depiction of retinal thickness. Chart format expressed showing central subfield thickness (ILM-RPE thickness), average macular thickness, and volume. For both eyes, the average thickness of all spots along the calculation circle is displayed. Each quadrant's average thickness is also displayed. The patient's data is compared to normative data in each of these charts. Symmetry is highlighted in the data table, which is also compared to normative data.

Statistical analysis

Statistical analysis was done by SPSS v26 (IBM Inc., Chicago, IL, USA). The mean and standard deviation (SD) of quantitative values were reported. Qualitative variables were presented as frequency and percentage (%). A P value ≤ 0.05 was used to determine statistical significance.

Results

Error! Reference source not found. showed gender and age distribution of patients and Subdivision of studied eyes as regard refraction

Table 1: Gender and age distribution of patients and Subdivision of studied eyes as regard refraction (n=30)

Gender	Male	6(20.0%)
	Female	24(80.0%)
	Total	30(100.0%)
Age	4 years	5(16.7%)
	5 years	9(30%)
	6 years	16(53.3%)
	Total	30(100.0%)
Refractive error	Myope	9(30%)
	Hypermetrope	6(20%)
	Strabismus	4(13.3%)
	Hypermetropic astigmatism	11(36.7%)
	Total	30(100%)

Data are represented by number (%)

BCVA in different ages of our patients (4, 5, 6 y) before and after treatment was statistically significant difference (P=0.001). The different types of refractive errors of the studied eyes regarding BCVA, before and after treatment was statistically significant difference(P=0.001).

(Error! Reference source not found.)

Table 2: The mean values of (BCVA) in different age groups and in studied eyes with different refractive errors

		Range (BCVA) logMAR			Mean	±	S. D	t. test	p. value
4 Years	Before	1	–	1.2	1.08	±	0.11	20.246	0.001*
	After	0.2	–	0.2	0.20	±	0.00		
5 Years	Before	1	–	1.2	1.13	±	0.10	17.964	0.001*
	After	0.3	–	0.5	0.32	±	0.07		
6 Years	Before	0.8	–	1.2	0.98	±	0.16	10.641	0.001*
	After	0.3	–	0.5	0.40	±	0.10		

Myope	Before	0.8	–	1.2	1.00	±	0.17	12.203	0.001*
	After	0.2	–	0.3	0.27	±	0.05		
Hypermetrope	Before	0.8	–	1.2	1.00	±	0.18	10.413	0.001*
	After	0.2	–	0.3	0.28	±	0.04		
Hypermetropic astigmatism	Before	0.8	–	1.2	1.02	±	0.17	9.961	0.001*
	After	0.3	–	0.5	0.45	±	0.09		
Strabismus	Before	1	–	1.2	1.10	±	0.12	5.563	0.001*
	After	0.5	–	0.8	0.63	±	0.13		

*: Statistically significant at $p \leq 0.05$, BCVA, Best Corrected Visual Acuity

CMT was statistically insignificant difference between amblyopic and non-amblyopic eyes.

AMT was statistically significant difference between amblyopic and non-amblyopic eyes (p value= 0.001). (**Error! Reference source not found.**)

Table 3: CMT and AMT evaluation of studied amblyopic and non-amblyopic eyes

	Amblyopic eye	Non-amblyopic eye	T. test	P. value
CMT	219 – 270	209 – 264	1.320	0.192
	237.7 ± 18.63	231.4 ± 18.35		
AMT	270 – 305	249 – 292	3.426	0.001*
	286.1 ± 10.61	275.4 ± 13.42		
Part time occlusion (PTO)	2 – 6	5.07 ± 1.72		

*: Statistically significant at $p \leq 0.05$, Data are represented by range or mean \pm SD, CMT: Central Macular Thickness, AMT: Average Macular Thickness

CMT was statistically insignificant difference in amblyopic eyes before and after patching.

AMT was statistically significant difference in amblyopic eyes before and after patching (p value= 0.004). (**Error! Reference source not found.**)

Table 4: CMT and AMT evaluation of studied amblyopic eyes before and after patching

	Before	After	T. test	P. value
CMT	209 – 270	210 – 268	1.405	0.163
	234.55 ± 18.60	229.70 ± 19.20		
AMT	249 – 305	212 – 291	2.921	0.004*
	280.75 ± 13.15	272.30 ± 18.14		

*: Statistically significant at $p \leq 0.05$, CMT: Central Macular Thickness, AMT: Average Macular Thickness

As regard best corrected visual acuity (BCVA) of amblyopic eye, BCVA was evaluated before patching, every month till 6 months after patching and the results were statistically significant (p value= 0.001) (**Error! Reference source not found.**)

Table 5: Follow up of BCVA of amblyopic eyes

BCVA of amblyopic eye (log MAR)		1.2	1	0.8	0.6	0.5	0.3	0.2	Total	X ²	P-value
Before	N	11	11	8	0	0	0	0	30	205.051	0.001*
	%	36.7 %	36.7 %	26.7 %	0.0%	0.0%	0.0%	0.0%	100.0 %		
1 month	N	4	15	9	2	0	0	0	30		
	%	13.3 %	50.0 %	30.0 %	6.7%	0.0%	0.0%	0.0%	100.0 %		
2 months	N	0	13	4	10	3	0	0	30		
	%	0.0%	43.3 %	13.3 %	33.3 %	10.0 %	.0%	.0%	100.0 %		
3 months	N	0	12	4	9	4	1	0	30		
	%	0.0%	40.0 %	13.3 %	30.0 %	13.3 %	3.3%	.0%	100.0 %		
4 months	N	0	2	3	8	11	6	0	30		
	%	0.0%	6.7%	10.0 %	26.7 %	36.7 %	20.0 %	.0%	100.0 %		

5 month s	N	0	0	2	5	13	9	1	30		
	%	0.0%	.0%0	6.7%	16.7%	43.3%	30.0%	3.3%	100.0%		
6 month s	N	0	0	1	2	9	14	4	30		
	%	0.0%	.0%	3.3%	6.7%	30.0%	46.7%	13.3%	100.0%		

Discussion

Amblyopia is treated mostly by occluding the superior eye with a patch that is opaque to improve vision. Two to six hours of partial occlusion each day is demonstrated to produce similar results as full-time occlusion when used as directed. One to two hours of maintenance patching every day is recommended frequently to inhibit amblyopia from recurring following patching with success because the length of the patch-on and patch-off intervals should be proportional to the severity of amblyopia^[12].

As regard refractive error with the mean BCVA, our study included 9 myope (1.00 ± 0.17), 6 hypermetrope (1.00 ± 0.18), 4 strabismus (1.02 ± 0.17) and 11 hypermetropia astigmatism (1.10 ± 0.12) whereas Yoon and Chun,^[13] showed unilateral anisometropic amblyopia; 10 hyperopia (3.5 ± 2.1) and 12 myopia (2.2 ± 1.9) as the remaining patients missed in follow up, EID et al.,^[12] revealed 26 Myopia (-7.07 ± 4.55) and 14 Hypermetropia (3.93 ± 1.96), moreover, Liu et al.,^[14] illustrated 20 anisometropic, 12 strabismic anisometropic, 5 strabismic amblyopia and 7 ametropic amblyopia but Kavitha et al.,^[15] included Unilateral anisometropic amblyopia 7 myopia, 5 hypermetropia and 18 astigmatism (Mean = 0.63 ± 0.405 for all types).

In our study, The duration of part-time occlusion varied from 2 – 6 hours per day for a month according to the severity of amblyopia with the mean value was 5.07 ± 1.72 ; if mild 2 hours per day and if severe 6 hours per day and follow up for 6 months. In Yoon and Chun,^[13] study PTO was 6 to 8 hours of partial occlusion for 11.6 ± 2.3 months, whereas Kavitha et

al., ^[15] study revealed occlusion therapy at the first visit (baseline) and every 3 months for one year.

On comparison amblyopic eyes with non-amblyopic eyes as regard CMT in our study, the mean values were 237.7 ± 18.63 and 231.4 ± 18.35 respectively that was statistically insignificant (p value=0.192), whereas on comparison as regard AMT, the mean values were 286.1 ± 10.61 and 275.4 ± 13.42 respectively that was statistically significant. In Liu et al., ^[14] study, between amblyopic and normal fellow eyes, no significant differences were detected in macular volume, or choroidal thickness, macular thickness.

In Kavitha et al., ^[15] investigation, in children with anisometropic amblyopia, the MT was more in amblyopic eyes ($286.9 \pm 6.52 \mu\text{m}$) in comparison with non-amblyopic fellow eyes ($240 \pm 10.45 \mu\text{m}$) and normal children with normal eyes ($239.8 \pm 4.294 \mu\text{m}$).

Pang et al., ^[16] studied children with an average age of 9.56 years and found that the amblyopic eye has a narrower inner and outer macula than the normal eye in amblyopic children with unilateral severe myopia. These results suggest that anatomic changes in the retinas of amblyopic children with unilateral high myopia are possible.

Dickmann et al., ^[17] studied OCT to measure macular thickness in 20 strabismic amblyopes, and it was discovered that the macular thickness in the amblyopic eye was greater than in the normal fellow eye in patients with strabismic amblyopia.

Kavitha et al., ^[15] reported that MT in amblyopic eyes was higher than normal eyes, and it reduced when BCVA improved after occlusion therapy.

In our study, as regard CMT of studied eyes, the mean was 234.55 ± 18.60 before patching and was 229.70 ± 19.20 after patching and this was statistically insignificant whereas the mean AMT was 280.75 ± 13.15 before patching and was 272.30 ± 18.14 after patching that was statistically significant. in contrary to our study, in Eid et al., ^[12] study the mean CMT was 188.30 ± 28.33 before patching and was 195.08 ± 33.88 after patching and this was

statistically significant but the mean AMT was 267.65 ± 20.60 before patching and was 268.40 ± 21.28 after patching that was statistically insignificant.

Liu et al.,^[14] found that the recovered amblyopic eyes' corrected macular and choroidal biometric values were not substantially different from the persistent amblyopic eyes'.

Yoon and Chun,^[13] before to and after occlusion therapy, found no statistical difference in the thickness of the inferior, superior, temporal, and nasal quadrants in the inner (3 mm diameter) and outer (6 mm diameter) macula of individuals with unilateral anisometropic amblyopia. None of these measures were significantly changed after occlusion therapy.

Kavitha et al.,^[15] observed statistically significant decrease in MT on each visit ($5 = 251.40 \pm 12.51$) compared to baseline (288.20 ± 6.62) [visit 1] however, in amblyopic eyes after occlusion therapy, there was no statistically significant difference in RNFLT.

Pang et al.,^[16] documented that before amblyopia treatment, the peripheral maculae of amblyopic eyes in children with myopic anisometropia were thinner than those of the other eyes. Following treatment for amblyopia, central maculae became thinner, but peripheral maculae remained same.

In our study, BCVA was evaluated before patching, every month till 6 months after patching and the results were statistically significant (p value= 0.001).

In Eid et al.,^[12] study, regarding BCVA, they noted a statistically significant difference when compared pre – (0.27 ± 0.12) and post - amblyopia treatment (0.49 ± 0.20).

Liu et al.,^[14] found the type of amblyopia (strabismic anisometropic amblyopia) and the initial severity of the amblyopia ($BCVA \leq 0.3$) were risk factors for persistent amblyopia in 47 – 96.3% of amblyopic people.

Kavitha et al.,^[15] observed the effects of occlusion therapy on BCVA after three, six, nine, and twelve months, and they discovered an enhancement in BCVA across all patients after one year of occlusion therapy.

Conclusions

The amblyopic eye has more CMT and AMT than the healthy eye and after 6 months of patching, there was a decrease in the CMT and AMT of the amblyopic eye.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

Ethical approval:

The Research Ethical Committee of Tanta University's Faculty of Medicine gave its approval to this study.

Consent

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

References

1. Repka MX, Beck RW, Holmes JM, Birch EE, Chandler DL, Cotter SA, et al. A randomized trial of patching regimens for treatment of moderate amblyopia in children. *Arch Ophthalmol*. 2003;121:603-11.
2. Thompson JR, Woodruff G, Hiscox FA, Strong N, Minshull C. The incidence and prevalence of amblyopia detected in childhood. *Public Health*. 1991;105:455-62.
3. Sjöstrand J, Abrahamsson M. Risk factors in amblyopia. *Eye (Lond)*. 1990;4 (Pt 6):787-93.
4. Liu H, Zhong L, Zhou X, Jin QZ. Macular abnormality observed by OCT in children with amblyopia failing to achieve normal visual acuity after long-term treatment. *J Pediatr Ophthalmol Strabismus*. 2010;47:17-23, quiz 4-5.
5. Chow KL. Failure to demonstrate changes in the visual system of monkeys kept in darkness or in colored lights. *Journal of Comparative Neurology*. 1955;102:597-606.
6. Rasch E, Swift H, Riesen AH, Chow KL. Altered structure and composition of retinal cells in darkreared mammals. *Exp Cell Res*. 1961;25:348-63.
7. Wendell-Smith CP. Effect of light deprivation on the postnatal development of the optic nerve. *Nature*. 1964;204:707.
8. Yen MY, Cheng CY, Wang AG. Retinal nerve fiber layer thickness in unilateral amblyopia. *Invest Ophthalmol Vis Sci*. 2004;45:2224-30.
9. Altintas O, Yüksel N, Ozkan B, Caglar Y. Thickness of the retinal nerve fiber layer, macular thickness, and macular volume in patients with strabismic amblyopia. *J Pediatr Ophthalmol Strabismus*. 2005;42:216-21.
10. Al-Haddad CE, Mollayess GM, Cherfan CG, Jaafar DF, Bashshur ZF. Retinal nerve fibre layer and macular thickness in amblyopia as measured by spectral-domain optical coherence tomography. *Br J Ophthalmol*. 2011;95:1696-9.

11. Nishi T, Ueda T, Mizusawa Y, Semba K, Shinomiya K, Mitamura Y, et al. Effect of optical correction on subfoveal choroidal thickness in children with anisohypermetropic amblyopia. *PLoS One*. 2017;12:e0189735.
12. Zakaria Eid M, Mahmoud Amin A, El-Sayed Mohamed M. Changes in macular thickness after part time occlusion in children with amblyopia using optical coherence tomography (OCT). *Al-Azhar Medical Journal*. 2020;49:117-24.
13. Yoon DH, Chun BY. Comparison of the Thickness and Volume of the Macula and Fovea in Patients with Anisometropic Amblyopia Prior to and after Occlusion Therapy. *Korean J Ophthalmol*. 2018;32:52-8.
14. Liu CH, Ong SJ, Huang CY, Wu WC, Kao LY, Yang ML. Macular Thickness, Foveal Volume, and Choroidal Thickness in Amblyopic Eyes and Their Relationships to the Treatment Outcome. *J Ophthalmol*. 2018;2018:1967621.
15. Kavitha V, Heralgi MM, Harishkumar PD, Harogoppa S, Shivaswamy HM, Geetha H. Analysis of macular, foveal, and retinal nerve fiber layer thickness in children with unilateral anisometropic amblyopia and their changes following occlusion therapy. *Indian J Ophthalmol*. 2019;67:1016-22.
16. Pang Y, Goodfellow GW, Allison C, Block S, Frantz KA. A prospective study of macular thickness in amblyopic children with unilateral high myopia. *Invest Ophthalmol Vis Sci*. 2011;52:2444-9.
17. Dickmann A, Petroni S, Salerni A, Dell'Omo R, Balestrazzi E. Unilateral amblyopia: An optical coherence tomography study. *J aapos*. 2009;13:148-50.