

# THE IMPACT OF PERIODONTAL THERAPY ON PERIODONTAL PATIENT WITH TYPE 2 DIABETES MELLITUS.

## ABSTRACT:

Diabetes is a common among general population with many oral manifestations; persistent deprived glycemic control has been associated with the incidence and progression of diabetes related complications including gingivitis and periodontitis which the most common cause of tooth loss, evidence suggests that periodontitis affects glucose control in diabetes.

The disease evoked bacteremia can cause elevation in serum pro inflammatory cytokines leading to elevated lipid levels and ultimately inflicting a hypoglycemic agent resistance syndrome and tributary to destruction of duct gland beta cells. Treating chronic odontology infections is crucial for managing polygenic disorder. Aim of this literature review is to look at the impact of odontology medical aid on glycemic management in sort II diabetic patients..

## INTRODUCTION:

Diabetes mellitus and obesity are the severe nutritional disorders of multifactorial nature, which altered almost over 100 million population. ( Gursoy UK, Marakoglu, Ersan S et al, 2006) Patient with polygenic disease have redoubled incidence and severity of periodontitis not accounted for by variations within the gingival microbic infection poor glycemic management has been systematically related to periodontitis severity. additionally recent proof suggests that hyper glycaemia might induce inflammatory protein production. (Engbretson et al, 2000). Studies by Saito and his coauthors studies that the periodontitis patients develop deep pockets , if deep pockets , depth. greater than 20 were there is significant relationship with individual current glucose tolerance and the

patient with insulin resistance has more deep pockets. Further studies described that periodontitis has effect on glucose control in diabetic patients . (Saito T, et al 2002.) . Diabetic individual had smaller amount caries and dental plaque as compare to non-diabetic population, lower salivary flow rates and buffer effect, and more frequent growth of yeasts than their non-diabetic control. Periodontal disease is most typical reason for tooth loss. it's insidious onset, chronic course and ordinarily result because of accumulative impact of dietary habits, oral hygiene ways and oral habits practiced over the years.( Shah N , Sundaram KR et al, 2003.) polygenic disease could be a cluster of infections and lesions poignant the odontology tissues that kind that attachment equipment of a tooth. (Stegeman CA ,2005.)

Laminin 5(ln-5) is concerned within the top migration of epithelial cells throughout the event of dental pockets. Low-dose Vibramycin (LDD) will therapeutically modulate the host response with its non-anti microorganism properties. (Emingil G et al,2004) In addition, proof supports the observation that dental medicine infections contribute to issues with glycemic control( Taylor GW,2003.) for e.g.: adults with polygenic disorder United Nations agency received inaudible scaling surgical procedure together with consistently administrated at 3 months, important reduction in mean HbA1c nearly 100 percent from pretreatment worth.(Grossi SG, Skrepinski FB, DeCaro ,1997) proof suggests that disease induced bacteriemia can cause elevation in humor professional inflammatory cytokines, resulting in hyperlipidermia associated ultimately inflicting an hormone resistance syndrome and conducive to destruction of duct gland beta cells.( Iacopino AM et al,2001) Treating chronic dental medicine infections is important for managing polygenic disorder.

### **Literature Review:**

Diabetes mellitus (DM) is a major health problem that affects people all over the world. According to the International Diabetes Federation's 2013 report, 8.3% of adults (or 382 million people) worldwide have diabetes, with the figure expected to climb to 592 million by 2035. (Diabetic federation 2017). Type 2 diabetes mellitus patient were 2.8 times

more likely to have destructive periodontal disease( Emrich LJ, Shlossman M, Genco RJ (1991) and 4.2 times more likely to have alveolar bone loss progression.( Taylor G, Burt B, Becker M, Genco RJ, ShlossmanM,1998). With poorly controlled polygenic disease, disease worsens, and with severe periodontic conditions, there's usually poorer glycemic management. revealed papers conclude that periodontic treatment results in a discount in haemoprotein A1c; but, different studies show restricted or no improvement. Some patients don't respond well to skilled periodontics medical care, even while not polygenic disease, that might be associated with poor oral hygiene and/or host factors together with microbe profiles. There area unit few revealed papers work the efficaciousness of periodontics medical care in diabetics and fewer that embody assessment of the oral microbiome. The investigators can examine general changes in polygenic disease standing and microbiome influences on clinical response to periodontics medical care in an exceedingly randomized test. it's been shown that antiseptic gluconate rinse results in vital clinical improvement over commonplace periodontics medical care alone, and interdental tooth cleaners take away a lot of plaque (leading to reduced animal tissue inflammation) effectively than brushing alone. periodontal disease has been considered to be another complication of diabetes mellitus and evidence by some author (Lo" e H et al,1993). also support poorer glycemic control contributing to poorer periodontal health.( Ainamo J, Lahtinen A, Uitto V 1990; Unal T, Firatli E, Sivas A, Meric H, Oz H 1993; Novaes AJ, Gutierrez F, Novaes A 1996; Taylor G, Burt B, Becker M 1998)

Recent studies exemplified the synergistic association between diabetes and periodontics. Sever periodontics was associated with low glucose control and exacerbated symptoms of diabetes included hyperglycemia.

Although it's been declared by some studies that improved glycemic levels might result in improved oral health,( Sastrowijoto, S. H., et al, 1990) it's still unclear whether or not the management of microorganism oral infection might improve or not the metabolic management among polygenic disorder. The aim of the current study was to analyze the impact of improved dentistry health on metabolic management in Diabetic patients. it's is studied that infections are usually common among tissue with glucose resistance Vki-

Jarvinen, H., Sammalkorpi, K., Koivisto, V. & Nikkila, E. 1989) A recent study has incontestable that in the acute part of a microorganism infection, endocrine resistance raised by thirty third whereas throughout the improve amount it raised by twenty eighth (sammalkorpi 1989)

Insulin resistance increased by 33% during the acute phase of a bacterial infection, but only by 28% throughout the convalescence stage, according to previous research (sammalkorpi 1989). moreover ,Grossi, S. G., Skrepinski, F.B., DeCaro, T., Zambon, J.J., Cummin, D. & Genco, R. J. 1996 have instructed that chronic gram-negative infections and chronic endotoxemia, like is seen in disease may additionally induce endocrine resistance and a worsening of metabolic management in diabetic patients. Seeable of those facts, it's been hypothesized that management of dentistry infections improves metabolic management of polygenic disorder. one study investigated many microorganism treatments in sort two diabetics. to work out their effectiveness compared to traditional non surgical medical aid.(Grossi 1996) Another report examined the correlation between animal tissue hemorrhage and blood sugar levels in nine diabetics with moderate –to – sever periodontal disease studied by some author(Jonsson B et al; 2009) during this study ,glycated hemoprotein (HbA 1c)and glycated simple protein were determined before dentistry medical aid, and four and eight weeks when medical aid. Periodontal medical aid consisted of one and a couple of sessions of root coming up with, oral hygiene directions, antibiotic drug one hundred mg for fourteen days, and sharpening at two weeks. A non important decrease in HbA1c from nine.44% to 9.01% was ascertained within the nine subjects. Glycated simple protein levels didn't modification when medical aid five to nine subjects had a uniform reduction in hemorrhage on inquiring (reduced at four to eight weeks) In these five subjects ,HbA1c was reduced from eight.7% to 7.8%,which was statistically important so, this small , short-run pilot study instructed that dominant dentistry inflammation might improve metabolic management of diabetics. (Grossi S. G., Skrepinski, F.B., DeCaro, T.,Zambon, J.J., Cummin, D. & Genco, R. J.1996).

## **Discussion**

The prevalence of diabetes in Saudi Arabia (24%) was among the top ten worldwide. More than 60% of the dentists who took part reported that diabetes patients make up 11 to 40% of their practise. However, because our sample was limited to two large cities without include rural areas, the percentage of diabetes participants in our study may not be applicable to the entire population of Saudi Arabia. At the same time, our analysis indicated that, when compared to dentists, dental assistants answered that less than 10% of their patients were diabetic, which may be explained by the assumption that medically challenged individuals choose to see dentists rather than dental assistants.

Study of 85 type 2 diabetes mellitus Native Americans (Grossi S. G., Skrepcinski, F.B., DeCaro, T., Zambon, J.J., Cummin, D. & Genco, R. J. 1996). Fasting blood glucose did not improve much following periodontal therapy, however there was a statistically significant drop in HbA1c. This improvement was thought to be the result of doxycycline inhibiting haemoglobin glycation. These findings were expanded upon in a subsequent publication that included 28 additional participants. The HbA1c level in the treatment group that showed the most improvement was 10.5 percent at the start and fell to 9.6 percent three months following periodontal treatment. Despite further improvement in periodontal probing depths and attachment levels, HbA1c levels returned to baseline at the 6-month evaluation. Periodontal therapy, in addition to doxycycline, was acknowledged in this study for the improvement in HbA1c. Study of eighty five subjects of diabetes mellitus Native Americans (Grossi S. G., Skrepcinski, F.B., DeCaro, T., Zambon, J.J., Cummin, D. & Genco, R. J. 1996). showed that fast blood sugar failed to considerably improve once odontology treatment, however a statically vital decrease in HbA1c was ascertained . This improvement was attributed to be a potential result of antibiotic inhibition of glycation of hemoglobin. Epidemiological data confirm that diabetes is a key risk factor for periodontitis; susceptibility to periodontitis is increased by approximately threefold in diabetic population. There is a significant relationship between degree of raised glucose levels and complications of periodontitis. The mechanisms that underpin the links between these two conditions are not completely understood, but involve aspects of immune functioning, neutrophil activity, and cytokine

biology. These findings were mentioned in any detail during a ulterior report by some researcher enclosed an extra twenty eight subjects with same finding and conclusion. In the treatment cluster that exhibited the best improvement, baseline HbA1c level was 10.5% and shriveled to 9.6% in three months once odontology treatment. However, by the six months examination, HbA1c levels came back to baseline, in spite of constant improvement in odontology inquiring depths and attachment levels. during this report, odontology medical aid, additionally to antibiotic, was attributable for the development in HbA1c. There is robust proof to counsel that the incidence and severity of odontology is influenced by the presence or absence of polygenic disorder of mellitus, in addition because the degree to that the sickness on top of things by patients (. Bacic M, Plancak D, Granic M 1988 ,Emrich IJ, Shlossman M, Genco RJ.1991, Lalla E, Lamster IB, statesman AM1998, Nelson RG, Shlossman M, Budding IM, PettitDJ, Saad MF, Genco Rj, Knowler Wc 1998 Novaes AB Jnr, Gutierrez FG, Novaes AB.1996.) Oliver RC and his colleagues in their study reported that the existence sever generalized {periodontitis|periodontal sickness|disease} might adversely influence the management of underlying general disease (Grossi SG, Genco RJ. 1998; Grossi SG, Skrepinski FB, DeCaro Tet al; .1996, . Miller LS, Manwell MA, Newbold D, Reding ME, RasheedA, Blodgett J, Kornman K. 1992Sammalkorpi K.1989, Taylor GW, Shlossman M.Kinder WC,Pettitt DJ et al;1998, William R,Mahan C.1960, Y ki-Jarvinen, H.et al; 1989)). Studies also acknowledge that the growing older population incorporates a way higher incidence of kind a pair of DM than do younger age teams (Gottsegen R.1990, Mealey BL 1998, Nathan DM, Dale DC, Federman Doctor of Divinity, 1996, u. s. formulary Drug Information1998).

Certain disorders, such as diabetes mellitus (DM), are acknowledged to be at higher risk to develop various infections (Grossi,S., Zambon, J, Ho,1994; Page, R. C. & Beck, J. D. et al;1997). Less research attention has been directed to account that chronic oral infections are common with systematic disease like metabolic syndrome, even with the opportunity that the morbidity can be reduced and survival rates of some diseases might be raised by an improvement in periodontal health in these population with risk factors.

(Beck , J. D et al; 1996, Offenbacher, S et al;1996) .Periodontitis as a chronic localized oral infection that triggers a systematic as well as local host immune-inflammatory response and that can be a source of bacteremia, because of the large epithelial surface with ulcerated periodontal, pockets (Ebersole ,J.L.&Cappelli,D.2000).Periodontitis, particularly in its intense scientific shape, is presently taken into consideration to persuade the pathogenesis or growth the threat of a few systematic illnesses (Garsia R,Henshaw, M.&Krall,E.2001).the organic dating among DM and periodontal disorder is nicely documented (Mattson J. S. & Cerutis, R et al; 2001, ,Soskoline ,W.A. & Klinger, A et al ; 2001).Periodontal disorder and DM are intently related and are exceptionally universal persistent illnesses with many similarities in pathobiology. Inflammation is a crucial participant with inside the affiliation,& its significance is simply now coming to light(Mealey, B. L,& Oates, T.W. 2006) DM is characterized as a metabolic disease related to a persistent hyperglycemic state. It became first proven that DM became a threat thing for periodontitis and substiquently the inverse relation is suggested proposed,i.e. that periodontitis can be a risk for diabetic complication, and this speculation has been supported with the aid of using numerous research (Grossi, S.G.2001, Iacopino AM. 2001, Lalla E, Lamster IB, Stern DM, Schmidt AM2001, Taylor GW. 2001, Katz, J, Bhattacharyya, I, Farkhondeh- Kish, F, Perez, F.M, Caudle, R.M.& Heft, M. W. et al :2005, Takeda, M, et al: 2006, Lim, L.P and Thait al; A.C.2007 ). Several current experimental research have addressed the mechanisms underlying the interplay in among DM and periodontitis. All said a robust inflammatory reaction characterized with the aid of using a big secretion of irritation mediators, in particular pro-inflammatory cytokines, that can have each neighborhood (periodontal destruction) and systemic (impaired glycaemic manipulate) outcomes( Grossi ,S.G.(2001) Iacopino ,A.2001, Lalla ,E,Lamster, I, Stern, D.& Schmidt, A.M.(2001), Nishimura ,Fet al; 2003, Genco ,R.J et al ; 2005). Various research had been posted at the impact of periodontal remedy on DM manipulate. Although a few authors found (Miller,L. S., Manwell, M,A., Newbold, D. 1992, . Grossi, S. G.et al;. 1996, Grossi SG et al ;2001, Rodrigues, D et al; 2003, Kiran , M et al; 2005) that periodontal remedy might also additionally have a

useful impact on glycemic control, now no longer all reported this developme( Seppala, M.& Ainamo, J;1993, Seppala & Ainamo ,J.1994, Aldrdge,J et al; . 1995, Smith,G.T., Greenbaum, C.J et al; 1996, Rothendler, J.A.& Garcia, R. I et al; 2007). Even now, at the start of twenty first century, the medical proof stays insufficient and inconclusive. With this background, a cynical have a look at became designed to decide whether or not an development with inside the periodontal fame of kind 2 diabetic topics is observed with the aid of using an development of their metabolic manipulate. Microbial dental plaque is the initiator of periodontal disorder. Systemic elements regulate all styles of periodontitis basically via their ordinary outcomes at the ordinary immune and inflammatory defences. Periodontal disorder has been characterised because the 6th hassle of diabetes. (Loe H. 1993) .A 1995 report by Aldridge JP showed that metabolic manipulate can be the maximum critical thing among periodontal fitness and IDDM. These facts help the speculation that diabetes and the extent of metabolic manipulate are critical modifiers of periodontitis. Data at the efficacy of periodontal care on enhancing glycemic manipulate in poorly managed diabetes are equivocal stated by Taylor GW in 1999. According to US Department of Health and Human & Services ,among diabetes and periodontitis has been postulated, however helping facts are sparse. Several research endorse that enhancements in periodontal circumstance will enhance glycaemic manipulate (Miller LS, Manwell MAet al ;1992,Taylor, G. Wet al; 1996, Grossi SG, Skrepcinski FB, DeCaro 1997; St Iwamoto ,Y et al ; 2001).Epidemiological research have proven an association among each type of diabetes and periodontal disorder (Emrich LJ et al; 1991, Moore PA et al; 1999). However accelerated periodontal threat is frequently associated with period and adequacy of manipulate of the diabetic state. For example, it's been referred to that people with non-insulin-structured diabetes mellitus have a 3-fold accelerated risk of growing periodontal disorder that cannot in any other case be defined on the idea of age, socioeconomic or oral hygiene (Emrich LJ et al;1991). In another study by Sastrowijoto, S. H et al in 1990 reported that people with terrible metabolic disorder had accelerated loss in comparison to control hyperglycemic population. It is likewise nicely documented that diabetic sufferers have a compromised capacity to reply



to bacterial infections, and it's been proposed that it's far this compromised host reaction that during component might also additionally growth diabetics threat of periodontal disorder. Interestingly, the opposite opportunity that the periodontal contamination might also additionally exacerbate the diabetic circumstance is now starting to get hold of growing attention (Grossi, S. G. & Genco, R. J. 1998) . The presence of glycated hemoglobin withinside the movement and in tissues, due to the hyperglycemia of diabetes, is assumed to be a contributing thing to the degenerative micro vascular and arterial adjustments which can be not unusualplace sequelae of diabetes. A have a look at of the Gila river Indian community, a populace having a incidence of non-insulin-structured diabetes mellitus of approximately 50%. Has lately examined the speculation that intense periodontitis in people with non-insulin-structured diabetes mellitus will increase the attention of glycated hemoglobin. (Taylor GW, Burt BA, Becker MP, Genco RJ, Shlossman M1998) The consequences confirmed that intense periodontitis at baseline became related to accelerated threat of getting terrible glycaemic manipulate at follow-up 2 or greater years later. If periodontal disorder does have an effect on diabetic fame, we might count on that treating periodontal disorder might lessen the severity of diabetes. A current systematic overview of the literature with the aid of using Grossi, S. G. & Genco, R. J. 1998). Concluded that the impact on diabetic fame became structured upon the remedy modality. Studies that look into the impact of best mechanical debridement had been not able to illustrate any impact on blood glucose degree or glycated hemoglobin degree no matter periodontal disorder severity or diploma of diabetes manipulate Aldridge JP, Lester V, Watts TL, Collins A, Viberti G, Wilson RF, 1995, O'Neil TC 1979, Seppala B, Ainamo J. 1994). However, all 3 research that introduced systemic antibiotics to mechanical debrident proven advanced metabolic manipulate of diabetes 2. (Miller LS, Manwell MA, Newbold D, Reding ME, Rasheed A, Blodgett J, Kornman KS 1992, O'Neil TC 1979, Williams RC Jr, Mahan CJ 1960). Results suggest that each one topics that had been handled with doxycycline skilled a discount in glycated hemoglobin (Grossi SG et al; 1996) These consequences endorse that periodontal antimicrobial treatment might also additionally lessen the extent of glycated hemoglobin in diabetic

topics and can in the long run keep the ability to lessen diabetic sequelae. There is an intensive frame of literature that factor to an emerging “-manner dating” among diabetes/terrible glycaemic manipulate and periodontal disorder. Specifically, the severity of the diabetic circumstance, whether or not insulin-structured diabetes mellitus is associated with the occurrence and severity of periodontal disorder (Soskolne WA, Klinger A. 2001) Conversely, there's proof that the severity of periondal disorder might also additionally have an effect on the extent of glycemic manipulate in diabetic sufferers (Grossi SG. 2001, Miller LS et al ; 2001, Taylor GW et al; 2002). A vast number of research have now been conducted to look into the impact of treating periodontitis on diabetic glycemic control. In persons with diabetes and periodontitis, some of these studies were conducted as randomised controlled trials in which periodontal treatment was compared to no periodontal treatment (or delayed periodontal treatment). This might bring about an improved blood glucose, which in flip ought to react with hematogenous proteins together with hemoglobin to shape glycosylated hemoglobin. Further glycation and oxidation of proteins and lipids might then cause formation pf superior glycation cease merchandise that would sell sequelae of diabetes together with neighborhood unfavorable inflammatory responses and tissue damage (Stewart JE et al; 2001)

## **Conclusion;**

It's been confirmed that periodontal disorder has a devastating impact on glycemic manipulate amongst kind 2 diabetic sufferers and a widespread discount of Glycated hemoglobin(HA1c), 0.40 %, became located after 3–four months of periodontal remedy finished on sufferers with each kind 1 and a couple of DM and periodontitis. For the dental practitioner, an critical vicinity of scientific studies facilities on enhancing glycemic manipulate (as measured with the aid of using the glycosylated hemoglobin) via periodontal remedy. Studies the usage of disinfectant, irrigation, and/or systematic antibiotics on each NIDDM and IDDM sufferers have proven small, however in a few instances widespread, discounts in glycosylated hemoglobin after periodontal therapy.

We concluded that periodontal remedy itself improves metabolic manipulate and reinforces that T2DM is a threat thing for periodontitis

#### REFERENCES:

1. Ainamo J, Lahtinen A, Uitto V (1990). Rapid periodontal destruction in adult humans with poorly controlled diabetes. A report of 2 cases. *J Clin Periodontol* 17: 22–28
2. Aldridge JP, Lester V, Watts TL, Collins A, Viberti G, Wilson RF (1995). Single-blind studies of the effects of improved periodontal health on metabolic control in type 1 diabetes mellitus. *J Clin Periodontol* 22: 271–275.
3. Basic M, Placak D, Granic M. (1988) CIPTN assessment of periodontal status in diabetics. *J Periodontol* 59: 816–822.
4. Beck, J. D., García, R., Heiss, G., Vokonas, P.S. & Offenbacher, S. (1996) Periodontal disease and cardiovascular disease. *Journal of Periodontology* 67, 1123–1137.
5. Christgau, M., Palitzsch, K.D., Schmalz, G., Kreiner, U. & Frenzel, S. (1998) Healing response to non-surgical periodontal therapy in patient with diabetes mellitus; clinical microbiological, and immunologic results *Journal of clinical Periodontology* 25, 112–124.
6. Ebersole, J.L. & Cappelli, D. (2000) Acute phase reactants in infections and inflammatory diseases. *Periodontology* 2000 23, 19–49.
7. Emrich LJ, Shlossman M, Genco RJ (1991). Periodontal disease in non-insulin-dependent diabetes mellitus. *J Periodontol* 62: 123–131.
8. Emrich G, Emingil G, Atilla G, Sorsa T, Savolainen P, Baylas H. *Periodontol*, 2004; 75 (10): 1387–96.
9. Engebretson SP, Hey-Hadavi J, Ehrhardt FJ, Hsu D, Celenti RS, Grbic JT, Lamster IB, Garcia, R, Henshaw, M. & Krall, E. (2001) Relationship between periodontal disease and systemic health. *Periodontology* 2000 25, 21–36.

10. Genco, R.J., Grossi, S.G., Ho, A., Nishimura, F. & Murayama, Y. (2005) A proposed model linking inflammation to obesity, diabetes and periodontal infections. *Journal of Periodontology* 76, 2075-2084.
11. Gottsegen R. (1990) Diabetes mellitus, cardiovascular diseases and alcoholism. In: Schluger S, Yuodelis R, Page RC, Johnson RH, ed. *Periodontal diseases*. 2nd edn. Philadelphia: Lea and Febiger, 273-279.
12. Grossi SG, Skrepcinski FB, DeCaro T et al (1997). Treatment of periodontal disease in diabetics reduces glycated hemoglobin. *J Periodontol* 68:713-9.
13. Grossi SG, Genco RJ. (1998) Periodontal disease and diabetes mellitus: a two-way relationship. *Ann Periodontol* 3: 51-61.
14. Grossi SG. (2001) Treatment of periodontal disease and control of diabetes: an assessment of the evidence and need for future research. *Ann Periodontol* 6:138-145
15. Gursoy UK, (2006) Marakoglu, Ersan S. Periodontal status and cytoplasmic enzyme activities in gingival crevicular fluid of type 2 diabetic and/or obese patient with chronic periodontitis. *Jan*;8(1);2-5.
16. Gustke CJ. (1999) Treatment of periodontitis in the diabetic patient. A critical review. *J Clin Periodontol* 26:133-137.
17. Hagiwara, S., Ogaawara, Y. & Tanaka, A. (2002) Effect of non surgical periodontal therapy on diabetic metabolic control. *Journal of Dental Research* 81(Special issue), 206(abstr 1551).
18. Hall R. Oral medications for diabetes. *Periodontal Insights* 1998;5:12-13.
19. Iacopino AM (2001). Periodontitis and diabetes interrelationships: role of inflammation. *Ann Periodontol*. 6(1):125-37
20. International Diabetes Federation. *IDF diabetes atlas*. 2017. <http://www.diabetesatlas.org/> (accessed 13/10/2017).
21. Iwamoto, Y. Nishimura, F., Nakagawa, M. (2001) The effect of antimicrobial periodontal treatment on circulating TNF- $\alpha$  and glycated hemoglobin level in patients with type 2 diabetes. *Journal of Periodontology* 72, 774-748.

22. Jonsson B, Ohrn K, Oscarson N, Lindberg P. An individually tailored treatment programme for improved oral hygiene: introduction of a new course of action in health education for patients with periodontitis. *Int J Dent Hyg.* 2009;7:166–175. doi: 10.1111/j.1601-5037.2008.00350.x
23. Jones , J. A., Miller, D. R., Wehler, C. J, Rich, S.E & Krall-Kaye, E. A. & Garcia ,R.I.(2007) Does periodontal care improve glycemic control ?The department of veterans affairs dental diabetes study. *Journal of Clinical Periodontology* 34, 46-52.
24. Katz, J., Bhattacharyya, I., Farkhondeh-kish, F., Perez, F. M., Caudle, R. M. & Heft, M.W.(2005) Expression of the receptor of advanced glycation end products in gingival tissue of type 2 diabetes patients with chronic periodontal disease: a study utilizing immunohistochemistry and RT-PCR. *Journal of clinical Periodontology* 32, 40-44.
25. Kiran ,M. Arpak, N. Unsal, E. & Erdogan, M.F.(2005) Effect of improved periodontal health on metabolic control in type 2 diabetes mellitus. *Journal of Clinical Periodontology* 32, 266-272.
26. Lotfy M, Adeghate J, Kalasz H, Singh J, Adeghate E. Chronic complications of diabetes mellitus: a mini review. *Curr Diabetes Rev.* 2017;13(1):3–10
27. Lalla E, Lamster IB, Stern DM, Schmidt AM (2001). Receptor for advanced glycation end products, inflammation, and accelerated periodontal disease in diabetes: mechanisms and insight into therapeutic modalities, *Ann Periodontol* 6:113-118.
28. Lim L.P., Tay, F.B.K., Sum, C.F. & Thai, A.C.(2007) Relationship between markers of metabolic control and inflammation on severity of periodontal disease in patient with diabetes mellitus. *Journal of Clinical Periodontology* 34, 118-123.
29. Loe H (1993). The sixth complication of diabetes mellitus. *Diabetes Care* 16: 476–480.
30. Mattson, J.S. & Ceruti, R. (2001) Diabetes Mellitus. A review of the literature and dental implication. *Compendium of Continuing Education in Dentistry* 22, 757-772.

31. Mealey BL(1998). Impact of advances in diabetes care on dental treatment of the diabetic patient. *Compendium Contin Educ Dent* 19: 41–58.
32. Mealey BL & Aots T.W.(2006) Diabetes mellitus and periodontal disease. *Journal of Periodontology* 77, 1289-1303.
33. Miller LS, Manwell MA, Newbold D, Reding ME, Rasheed A, Blodgett J, Kornman KS (1992). The relationship between reduction in periodontal inflammation and diabetes control : a report of 9 cases, *J Periodontol* 63:843-848.
34. Moore PA, Weyant RJ, Mongelluzzo MB, Myers DE, Rossie K, Guggenheimer J, Block HM, Huber H, Orchard T. (1999) Type I diabetes mellitus and oral health: assessment of periodontal disease. *J Periodontol* 70: 409–417.
35. Nathan DM, Dale DC, Federman DD, ed. Diabetes mellitus. *Sci Am Med* 1996: 6: 1–24.
36. Nelson RG, Shlossman M, Budding IM, Pettit DJ, Saad MF, Genco RJ, Knowler WC (1990) .Periodontal disease and NIDDM in Pima Indians. *Diabetes Care* 13: 836–840.
37. Nishimura, F., Iwatomo, Y., Mineshiba, J., Shimizu, A., Soga, Y. & Murayama, Y. (2003) Periodontal disease and diabetes mellitus; the role of tumor necrosis factor - $\alpha$  in a 2 way relationship. *Journal of Periodontology* 74,97-102.
38. Novaes AJ, Gutierrez F, Novaes A (1996). Periodontal disease progression in type II non-insulin-dependent diabetes mellitus patients (NIDDM). Part I – Probing pocket depth and clinical attachment. *Braz Dent J* 7: 65–73
39. Offenbacher, S., Katz, V., Fertik, G. et al.(1996) Periodontal infection as a possible risk factor for preterm low birth weight. *Journal of Periodontology* 67,1103-1113.
40. Oliver RC, Tervonen T(1993). Periodontitis and tooth loss. Comparing diabetics with the general population. *J Am Dent Assoc* 124: 71–76.
41. O’Neil TC.(1997) Plasma female sex-hormone levels and gingivitis in pregnancy. *J Periodontol* 50: 279–282.

42. Page, R. C. & Beck, J. D. (1997c) Risk assessment for periodontal diseases. *International Dental Journal* 47, 61–67.
43. Ringelberg ML, Dixon DO, Francis AO, Plummer RW.(1997) Comparison of gingival health and gingival crevicular fluid flow in children with and without diabetes. *J Dent Res* 56: 108–111.
44. Rodrigues, D, Taba, M, Novaesd, A, Souza, S. & Grisi, M. (2003) Effect of non-surgical periodontal therapy on glycemic control in patients with type 2 diabetes mellitus. *Journal of Periodontology* 74, 1361-1367.
45. Sammalkorpi, K. (1989) Glucose intolerance in acute infections. *J Intern Med* 255, 15–19.
46. Sastrowijoto SH, van der Velden U, van Steenberghe TJ, Hilleman P, Hart AA, de Graaff J, Abraham-Inpijn L (1990). Improved metabolic control, clinical periodontal status and subgingival microbiology in insulin-dependent diabetes mellitus. *J Clin Periodontol* 17: 233–242.
47. Scannapieco, F. A. & Genco, R. J. (1999) Association of periodontal infections with atherosclerotic and pulmonary diseases. *Journal of Periodontal Research* 34, 340–345.
48. Seppala, B, Seppala M. & Ainamo, J. (1993) A Longitudinal study on insulin-dependent diabetes mellitus and periodontal disease. *Journal of Clinical Periodontology* 20, 161-165.
49. Seppala B, Ainamo J.(1994) A site-by-site follow-up study on the effect of controlled versus poorly controlled insulin-dependent diabetes mellitus. *J Clin Periodontol* 21: 161–165.
50. Shlossman M, Knowler WC, Pettitt DJ, Genco RJ.(1990) Type 2 diabetes mellitus and periodontal disease. *J Am Dent Assoc* 121: 532–536
51. Shah N , Sundaram KR (2003). Impact of socio-demographics variables, oral hygiene practices and oral habits on periodontal health status of Indian elderly: a community –based study .*Indian J Dent Res*.14(4):289-97.

52. Smith, G. T, Greenbaum, C.J ,Johnson, B.D.& Person, G. R.(1996) Short- term responses to periodontal therapy in insulin-dependent diabetic patients. Journal ofperiodontology 67, 794-802.
53. Soskolne WA, Klinger (2001) A. Relationship between periodontal disease and diabetes: an overview. Ann Periodontal :6:91-98.
54. Stewart JE, Wager KA, Friedlander AH, Zadeh HH. (2001) The effect of periodal treatment on glycemic control in patients with type 2 diabetes mellitus. Clin Periodontol :28:306-310
55. Stegeman CA (2005).Oral manifestations of diabetes. Home Health Nurse. 23(4): 233-40; quiz 241.
56. Steenbergen, T. J., Hillemans, P., Hart, A. A.,de Graff, J. & Abraham-Inpijn, L. (1990) Improved metabolic control, clinical periodontal status and subgingival microbiology in insulin-dependent diabetes mellitus. Journalof Clinical Periodontology 17, 233–242.
57. William R,Mahan C. (1960) Periodontal disease and dibetes in young adults.JAMA 172:776-778.
58. Yki-Jarvien H, Sammalkorpi K, Koivisto VA, Nikkila EA(1989). Severity,duration and mechanisms of insulin resistance during acute infections. J Clin Endocrinol Metab 69:317–323.