

Helicobacter pylori infection and circulating ghrelin levels- a review article.

ABSTRACT :

The correlation amidst *Helicobacter pylori* contamination & ghrelin levels flowing in the body is still an arguable subject. The enteric enteroendocrine system produces ghrelin, which is then octanoylated by, as of late, found ghrelin o-acyltransferase (GOAT) before being emitted into the circulatory system. Since ghrelin ties to the ghrelin neuroreceptor only after its acylation, this octanoylation is needed for a long time for ghrelin's natural components, like hunger incitement and calming characteristics (GHS-R). Given the site of ghrelin manufacture in the gut, it is expected that gastric mucosal injury impacts the flow of ghrelin levels among humans. *H. pylori* bacterium can contaminate > 50% of the world's citizens & can live for a lifetime once got rooted within the gastric mucosa. Chronic gastritis, stomach shrinkage, and ulceration, decreased appetite, and a decreased BMI are all connected to infection (BMI). The vast majority of research looking at flowing hunger hormone & ghrelin expression in the gut among patients with the contamination show that the bacteria inhibit ghrelin production and secretion. Ghrelin is restored once infection is eradicated, improving appetite and raising BMI. However, a causal association amidst *H. pylori*-related serum ghrelin reduction & edible consumption & fatness, and adiposity has yet to be shown in specific investigations. The majority of research looks at total ghrelin in the blood; however, the proportion of acyl/total hunger hormone may give a clear picture of how the acylated hunger hormone changes under the course of contamination & deterioration.

Keywords: *Helicobacter pylori*, Ghrelin, Gastritis, Enteroreceptor

Introduction:

❖ H. PYLORI: BACTERIUM

H. pylori characteristics : GRAM -ve .

Microaerophilic bacteria with helical structure.

Natural colonization of the stomach lamina of individuals and nonhuman mammal.

Not found in other animals.

Prevalence – high ; around 80% . (1)

Studies show that the frequency of *H. pylori* + status depends on a variety of components, including aging, zone, lifestyle, the standard of living, and socioeconomic status. (2) The main route of *H. pylori* infection is speculated as oral-oral transmission. This helps to explain why the infection is so common among family members, such as parents and children. In this approach, it appears that sharing utensils during meals is vital for setting up an infection. (3). Fecal-oral spread is a different way of contamination happens because of consumption of contaminated water, primarily due to inadequate sanitation. (4) It's also worth noting that rising socioeconomic position and better living conditions are both significant contributors to the decline in *Helicobacter pylori* contamination presence. (5)

W. JR, M. B discovered *Helicobacter pylori* contamination in the abdomen, until then. Because of its high acidity, the gastrointestinal environment was thought to be germ-free. (6) (7) (8)

The bacteria can use a variety of procedure to achieve successful colonization under such harsh conditions, including enhanced motility, a synthetic device that facilitates the development of a favourable environment during contamination maintenance as well as joining to epithelium. (9) (10) (11)

Ghrelin is restored once infection is eradicated, improving appetite and raising BMI. However, a causal association amidst *H. pylori*-related serum ghrelin reduction & edible consumption & fatness, and adiposity has yet to be shown in certain investigations. The majority of research look at total ghrelin in the blood however, the proportion of acyl/total hunger hormone may give a clear picture of how the mechanism of acylated hunger hormone changes under course of contamination & deterioration.

Furthermore, body defense mechanism have an important part in the process of contamination and its progression, most likely through a T helper cell one feedback against the bacteria. (12)

While almost *Helicobacter pylori* infections are not symptomatic, they increase the risk of developing illnesses such as peptic ulcers and stomach adenocarcinomas. As a result, competent clinical care, including a capable interpretation and successful treatment, are critical steps in improving a patient's clinical outcome. (13) (14)

Distinctive assortment of intrusive and non-obtrusive symptomatic techniques have been utilized to identify *h.pylori* and, in regards to treatment, bacterial obstruction addresses a significant test in contamination destruction. (15)

❖ **GHRELIN** :

LENTOMORELIN, also known as hunger hormone is a twenty eight - amino corrosive gastric-obtained peptide. Ghrelin-delivering cells are a different gathering of endocrine cells found all through the gastric mucosa, as well as the small intestine and the endocrine pancreas to a lesser extent. Ghrelin levels in the blood increase during fasting and chronic caloric restriction to encourage food intake and fat storage while also preventing life-threatening blood glucose dips.

Ghrelin inhibits the proliferation of breast, lung, and thyroid cell lines, as well as protecting the gastric mucosa. Ghrelin accelerates stomach emptying and increases acid secretion in the gastrointestinal tract via vagal stimulation. Ghrelin levels fluctuate a lot depending on how much

energy the body needs. Exogenous ghrelin has been shown to enormously decrease (NF)-B and plasma tumour necrosis factor activation. (16)

❖ **THE GUT- BRAIN AXIS AND GHRELIN :**

The aquaphobic octanoyl moiety esterhuizen to the 3rd serine remnant makes hungerhormone a twenty eight -aminoalkanoic -acid polypeptide autacid. "Reverse pharmacology" was utilized to uncover ghrelin, which was previously thought to be an founding G protein-linked enteroreceptor but is now recognised as the major ghrelin receptor, using the endogenous growth hormone secretagogue receptor (GHS-R). Following initiation of the GHS-R, it revealed that hunger hormone is a potent stimulant of GH production from mammalian somatotroph cells. It was discovered a fresh element of the somatotrophin for the first time. Phospholipase C, cAMP, and the nitric oxide/cGMP system are all involved in this activity.

The octanoylated mature hormone is released into the general bodyflow via the capillary networks of the gastric lamina propria. The stomach mucosa has the highest levels of ghrelin expression and secretion, whereas the pituitary and hypothalamus nuclei have the highest levels of GHS-R, leading to the theory that ghrelin and its receptor evolved to establish a connection.

The processes behind ghrelin's co- and post-translational alterations are only now beginning to be unravelled. The hydrophobicity imparted by acylation may allow ghrelin to pass the blood-brain barrier in both directions. This change also makes it easier for ghrelin to bind to the GHS-R, which is necessary for GHS-R-mediated ghrelin action. while the pancreatic may be the most prominent locus of expression in human tissues.

Moreover , it's tempting to think that paracrine GOA Texpressing cells could acylate unacylated ghrelin produced from the stomach. (17) (18)

❖ **ACYLATED GHRELIN IS NOT THE ONLY GHRELIN :**

Several ghrelin isoforms, including the splice variant des-Gln14-ghrelin and the unmodified des-octanoyl or unacylated ghrelin, have been found ghrelin/GHS-R axis may be broadened to include them. Unacylated ghrelin cant join to & thus doesn't initiate the GHS-R, inspite among the majority of prevalent species of ghrelin in serum. Unacylated ghrelin, once assumed to be a by-product of bioactive ghrelin deterioration , speculated to be a vital hormone with a wide range of biological functions, including cardiovascular function, bone physiology, reproductive axis, and foetal growth. (19) These seemingly inverse biological features are most likely conclusion of alternate receptor activation, which supports the idea of a non-discovered non-GHS-R ghrelin receptor (s) (20) (18) .

▪ **Ghrelin suppresses inflammation in rodent models of disease and in humans :**

Ghrelin and GHS-R are expressed in immune cells, and ghrelin/GHS-R activity and expression alter T cell function. Ghrelin antagonises leptin in immune cells, just as it does in the hypothalamus. Human_T cells activated by leptin produce more proinflammatory, anorectic cytokines like as IL-1, IL-6, and TNF, as well as enhanced GHS-R1a expression. In a dose-dependent manner, cotreatment with ghrelin suppresses leptin-induced cytokine levels. Ghrelin knockdown increases Th1 cytokine production and IL-17 secretion in primary human T cells,

implying a function for autocrine/paracrine ghrelin in the endogenous control of pro-inflammatory cytokine production and secretion. (21) (22)

The number of studies confirming ghrelin's anti-inflammatory effect in vivo is quickly expanding, and includes animal models of pancreatitis and colitis. Downregulation of pro-inflammatory cytokines, inflammation-suppressing regulatory T cells, and elevated levels of the anti-inflammatory cytokine IL-10 have all been linked to ghrelin's therapeutic effect. (23). Ghrelin therapy also reduces pro-inflammatory cytokine production in brain and spinal cord resident macrophages (microglia), DECREASEDing the severity of experimental autoimmune encephalomyelitis, a model of multiple sclerosis. (24) (18) Ghrelin has been successfully used as an anti-inflammatory medication in cachexic individuals with persistent respiratory infection and inflammation in clinical trials. Ghrelin medication boosted body weight and dramatically reduced inflammation in the lungs in these individuals by DECREASEDing neutrophil infiltration/accumulation and serum TNF-alpha . (25)

➤ **SYNTHESIS OF DATA :**

The information gathered was divided in 3 categories:

- 1) Statistics analyzing flowing ghrelin concentrations in H. pylori +VE and -VE participants.
- 2) Statistics analyzing flowing ghrelin concentrations prior and post H. pylori removal .
- 3) Statistics evaluating among the gastric ghrelin characteristics.

SR NO.	RESEARCH QUESTIONS :	EXPLANATORY DATA :
1	What is the link amidst helicobacter pylori & circulating ghrelin levels ?	Statistics analyzing flowing ghrelin concentrations in H. pylori +VE and -VE participants .
2	Consequences of helicobacter elimination on flowing quantity of ghrelin in body ?	Statistics analyzing flowing ghrelin concentrations prior and post H. pylori removal .
3	What are consequences on gherlin levels in the gut due to helicobacter pylori infection ?	Data assessing any of the gastric ghrelin parameter

TABLE 1 : Research questions explored by the review

DISCUSSION :

➤ SOURCES OF HELICOBACTER PYLORI AND GHERLIN :

GHERLIN : Ghrelin is mostly generated by the stomach, with smaller amounts coming from the bowel, pituitary, kidney, placenta, hypothalamus, and pituitary, kidney, placenta, and hypothalamus. (23) (26) .

The pancreatic islet's A-cells, the lung, and the kidney As a result, It's crucial to figure out which organ has the most influence .variations in ghrelin levels in a variety of people illnesses. Despite the fact that the liver produces the majority of the circulating ghrelin. Additional sources of ghrelin secretion, such as those generated in the stomach, can enhance or reduce ghrelin secretion in a compensatory manner. (27) (28) (29) (30)

H. pylori infects more than half of the adult population on the planet. Atrophic gastritis and intestinal metaplasia are the earliest symptoms of H pylori infection, which can progress to dysplasia and gastric cancer. Thus, in case of H. pylori contagiousness influences gastric ghrelin synthesis & , as a result, plasma ghrelin collection is a fascinating subject. (27)

➤ GHERLIN AND THE REGULATION OF GLUCOSE HOMEOSTAIS :

When injected into people, ghrelin causes an increment of plasma glucose & a decrement insulin levels. This coincided along the identification of the GHS-R in the pancreas islets . As a result, most research looking at the effects of ghrelin on GSIS have found that it inhibits it .

It's worth noting that the pancreas produces ghrelin, and ghrelin's influence on the endocrine pancreas could be via a paracrine mechanism. Endogenous ghrelin was found to increase insulin secretion when it was blocked, implying that ghrelin inhibits insulin production by acting directly on pancreatic -cells.

➤ RELATION AMIDST BMI OF HELICOBACTER PYLORI INFECTED PATIENTS AND PLASMA GHRELIN QUANTITY :

Several investigations have demonstrated that flowing ghrelin is high among persons with anorexia nervosa, DECREASED in adiposity , and normalised with increase in weight or decrease in weight , indicating that ghrelin has involvement in balance of energy on a long term basis . longitudinal and cross- sectional investigations of anorexia nervosa and obesity reveals strong link among flowing hungerhormone quantity in body and/with body fat percentage, fat mass, BMI, body weight, insulin, leptin, and T3. The link amidst Body Mass Index & flowing ghrelin quantity in body was not strong in Helicobacter pylori contaminated participants. However this revealed that plasma ghrelin quantity is highly impacted by H. pylori infection. (27) (31) (32)(33) (34) (27) (28) (29)

➤ **EFFECTS OF THE H.PYLORI ON GASTRIC ENDOCRINE SYSTEM :**

Helicobacter pylori contamination is antrum predominant in the majority of infected people, and acidic content produced mainly by not affected corpus is increased, increasing the risk of duodenal ulcers. (35) (29)

It was observed that ,Acid produced in the family members of patient with gastric cancer was lower in quantity compared to that of normal civilization , and that patients with antral predominant H. pylori infection and gastritis developed corpus predominant infection, the initial infection may start in the antrum and spread to the corpus among people in whom low basic acid is produced. (36) (37)

Hypochlorhydria allows other bacteria to infect you, which can increase the creation of carcinogenic (e.g. N-nitroso) chemicals. (34)

In chronic gastric inflammation and infection occurred due to Helicobacter pylori infection, have decreased control over the production of gastric juice, however the effect of helicobacter bacteria on endocrine system depends on the area of infection , or following influence of the endocrine system, like acid produced. (38)

➤ **CONSEQUENCES OF HELICOBACTER PYLORI ON ENDOCRINE CELLS :**

Helicobacter pylori colonisation causes dispersion of chemo attractants like IL-8, IL1, and TNF, which give signals to G cells, and the number of gastrin cells increases in Helicobacter pylori contaminated gastric lamina . (39)

Simultaneously, the amount of D cells making SST decrease in quantity . In the same way, some swine Helicobacter species change the no. of endocrine cells in the stomach lamina. (40)

A few changes among these , such as an increment among no. of G cells, a decrement among no. of D cells, and specially an increment in the G/D cell proportion , have been linked to the formation of gastroesophageal ulcers in pigs. (41)

➤ **CONSEQUENCE OF HELICOBACTER PYLORI ON CIRCULATING GHERLIN LEVELS :**

Because cells that produce hunger hormone are localised in stomach, it stands to reason that chronic gastritis and atrophy would impede hungerhormone manufacture , acylation, &/or emission , affecting hunger, weight, and BMI.

Some of the discrepancies in the literature about the effect of H. pylori infection on circulating ghrelin levels could be because of difference among people and civilization, illness severity (for ex, presence or absence of atropy), and Helicobacter pylori strain dissimilarity . This is further exacerbated by the fact that ghrelin is measured using a variety of immunoassays and that acylated ghrelin is not stable and deteriorate fast to unacylated ghrelin. (42)

The best way to assess plasma ghrelin is a debated topic; some researchers believe that measuring total ghrelin accurately reflects activated ghrelin quantity & is an appropriate method. The majority of studies looking into the effects of H. pylori have only looked at total ghrelin

levels, and the majority of these studies found that infection lowers plasma ghrelin levels. [TABLE :2]

TABLE 2. Total ghrelin levels

<u>SUBJECT</u>	<u>NATIONALITY</u>	<u>SAMPLE TYPE</u>	<u>TOTAL GHRELIN IN H.PYLORI PATIENT</u>	<u>REFERENCE</u>
39 adults , F	Turkish	Plasma	INDISTINGUISHABLE	(43)
256 adults , M	USA	Serum	INDISTINGUISHABLE	(44)
85 children , F+ M	ITALIAN	Serum	INDISTINGUISHABLE	(45)
196 adults , F	TAIWAN	Plasma	INDISTINGUISHABLE	(46)
63 adults , F+M	KOREAN	Plasma	INDISTINGUISHABLE	(47)
41 adults , F+M	KOREAN	Plasma	INDISTINGUISHABLE	(48)
50 adults , F + M	TURKISH	Plasma	INDISTINGUISHABLE	(48)
110adults , F+M	USA	Serum	INDISTINGUISHABLE	(49)
24 adults , F+M	USA	Plasma	INDISTINGUISHABLE	(50)
13 adults , F+M	—	Serum	INDISTINGUISHABLE	(51)
220 adults , F+M	Japanese	Plasma	DECREASED	(52)
81 adults , F+M	Japanese	Plasma	DECREASED	(53)
287 children , F+M	Poland	SERUM	DECREASED	(54)
160 adults , M	Japanese	Plasma	DECREASED	(55)
145 adults , M	Taiwan	Plasma	DECREASED	(46)
15 adults , F+M	Spain	Plasma	DECREASED	(56)
62 adults ,F+M	France	Plasma	DECREASED	(57)
100 adults, F+M	Chinese	Plasma	DECREASED	(58)
249 adults , F+M	Japanese	Plasma	DECREASED	(59)
74 adults ,	Japanese	Plasma	DECREASED	(60)

F+M				
100 adult , F	Poland	Serum	DECREASED	(61)
180 adult and children , F+M	Poland	Serum	DECREASED	(62)
68 adults , F+M	Japanese	Plasma	DECREASED	(63)
79 adults , F+M	Italian	Serum	DECREASED	(64)
89 adults ,F+M	Japanese	Plasma	DECREASED	(65)

However, a common thread running across these research is that abdominal epithelial cells degradation can lead to reduction in total plasma ghrelin ,& there is a negative link admist plasma ghrelin quantity & atropic seriousness and intesity in a subset of these investigations was found. (52) (66)

Because acyl- and unacylated ghrelin have potentially different, even inverse biological effects, therefore proportion of changed to unchanged hungerhormone is critical. Few studies have taken this under notice in the background of Helicobacter pylori contamination , and those that have found inconsistent results. [TABLE : 3]

TABLE : 3 ORIGINAL STUDIES MEASURING THE EFFECT OF H.PYLORI INFECTION ON PLASMA ACYL AND UNACYLATED GHRELIN LEVELS

Subjects	Nationality	Tissue	Acyl and total ghrelin in <i>H. pylori</i> infection (or cure)
69 Adults F + M	Japanese	P	↓ acyl ghrl in atrophy only
50 Adults F + M	Italian	P	↑ acyl ghrl and acyl/ total ratio in atrophy cf. healthy controls
220 Adults F + M	Japanese	P	↓ acyl ghrl associated with atrophy and ↑ after cure

Ref
(67)
(66)
(52)

The acylated ghrelin/total ghrelin proportion along with plasma acyl ghrelin levels are DECREASED in Japanese adults, whereas there is a notable rise in acyl ghrelin & the proportion of acylated ghrelin/total ghrelin in Western males, which the authors postulate may be because of an autogenous , balancing increase in the acylation mechanism as a result to a loss The degree of articulation and function of GOAT while Helicobacter pylori contamination , gastric infection and inflammation since long time , & degeneration of its mucosa would be a logical expansion of this research. (66-69)

CONCLUSION :

According to existing research, the concentration of circulating ghrelin in patients contaminated with Helicobacter pylori is DECREASED compared to those that are not contaminated with bacteria . Although , a more complicated connection between the amount of flow of gherlinin the body and helicobacter elimination. The strain of infecting H pylori, the length of follow-up, the amount of H pylori-induced gastritis, and other underlying disease may all influence this

connection. There is requirement of research is needed to fully understand the influence of *Helicobacter pylori* eradication on circulating ghrelin levels.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References:

1. Clyne M, Dolan B, Reeves EP. Bacterial factors that mediate colonization of the stomach and virulence of *Helicobacter pylori*. *FEMS Microbiol Lett*. 2007 Mar;268(2):135–43.
2. Prevalence of *Helicobacter pylori* infection worldwide: a systematic review of studies with national coverage - PubMed [Internet]. [cited 2021 Aug 19]. Available from: <https://pubmed.ncbi.nlm.nih.gov/24563236/>
3. Role of infected grandmothers in transmission of *Helicobacter pylori* to children in a Japanese rural town - PubMed [Internet]. [cited 2021 Aug 19]. Available from: <https://pubmed.ncbi.nlm.nih.gov/23560808/>
4. Goh K-L, Chan W-K, Shiota S, Yamaoka Y. Epidemiology of *Helicobacter pylori* infection and public health implications. *Helicobacter*. 2011 Sep;16 Suppl 1:1–9.
5. W L, F I, B I. Seroprevalence of *Helicobacter pylori* infection in Polish children and adults depending on socioeconomic status and living conditions. *Advances in medical sciences* [Internet]. 2014 Mar [cited 2021 Aug 19];59(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/24797992/>
6. Warren JR, Marshall B. Unidentified curved bacilli on gastric epithelium in active chronic gastritis. *Lancet*. 1983 Jun 4;1(8336):1273–5.
7. Wroblewski LE, Peek RM. *Helicobacter pylori*, Cancer, and the Gastric Microbiota. *Adv Exp Med Biol*. 2016;908:393–408.
8. Boquet P, Ricci V. Intoxication strategy of *Helicobacter pylori* VacA toxin. *Trends Microbiol*. 2012 Apr;20(4):165–74.
9. Eaton KA, Morgan DR, Krakowka S. Motility as a factor in the colonisation of gnotobiotic piglets by *Helicobacter pylori*. *J Med Microbiol*. 1992 Aug;37(2):123–7.

10. Alm RA, Bina J, Andrews BM, Doig P, Hancock RE, Trust TJ. Comparative genomics of *Helicobacter pylori*: analysis of the outer membrane protein families. *Infect Immun*. 2000 Jul;68(7):4155–68.
11. Camilo V, Sugiyama T, Touati E. Pathogenesis of *Helicobacter pylori* infection. *Helicobacter*. 2017 Sep;22 Suppl 1.
12. Bamford KB, Fan X, Crowe SE, Leary JF, Gourley WK, Luthra GK, et al. Lymphocytes in the human gastric mucosa during *Helicobacter pylori* have a T helper cell 1 phenotype. *Gastroenterology*. 1998 Mar;114(3):482–92.
13. Malfertheiner P, Venerito M, Schulz C. *Helicobacter pylori* Infection: New Facts in Clinical Management. *Curr Treat Options Gastroenterol*. 2018 Dec;16(4):605–15.
14. Abadi ATB, Kusters JG. Management of *Helicobacter pylori* infections. *BMC Gastroenterol*. 2016 Aug 12;16(1):94.
15. Safavi M, Sabourian R, Foroumadi A. Treatment of *Helicobacter pylori* infection: Current and future insights. *World J Clin Cases*. 2016 Jan 16;4(1):5–19.
16. Stec-Michalska K, Malicki S, Michalski B, Peczek L, Wisniewska-Jarosinska M, Nawrot B. Gastric ghrelin in relation to gender, stomach topography and *Helicobacter pylori* in dyspeptic patients. *World J Gastroenterol*. 2009 Nov 21;15(43):5409–17.
17. Kojima M, Kangawa K. Ghrelin: structure and function. *Physiol Rev*. 2005 Apr;85(2):495–522.
18. Jeffery PL, McGuckin MA, Linden SK. Endocrine impact of *Helicobacter pylori*: focus on ghrelin and ghrelin o-acyltransferase. *World J Gastroenterol*. 2011 Mar 14;17(10):1249–60.
19. Soares J-B, Leite-Moreira AF. Ghrelin, des-acyl ghrelin and obestatin: three pieces of the same puzzle. *Peptides*. 2008 Jul;29(7):1255–70.
20. Chen C-Y, Chao Y, Chang F-Y, Chien EJ, Lee S-D, Doong M-L. Intracisternal des-acyl ghrelin inhibits food intake and non-nutrient gastric emptying in conscious rats. *Int J Mol Med*. 2005 Oct;16(4):695–9.
21. Dixit VD, Schaffer EM, Pyle RS, Collins GD, Sakthivel SK, Palaniappan R, et al. Ghrelin inhibits leptin- and activation-induced proinflammatory cytokine expression by human monocytes and T cells. *J Clin Invest*. 2004 Jul;114(1):57–66.
22. Dixit VD, Yang H, Cooper-Jenkins A, Giri BB, Patel K, Taub DD. Reduction of T cell-derived ghrelin enhances proinflammatory cytokine expression: implications for age-associated increases in inflammation. *Blood*. 2009 May 21;113(21):5202–5.
23. Gonzalez-Rey E, Chorny A, Delgado M. Therapeutic action of ghrelin in a mouse model of colitis. *Gastroenterology*. 2006 May;130(6):1707–20.

24. Theil M-M, Miyake S, Mizuno M, Tomi C, Croxford JL, Hosoda H, et al. Suppression of experimental autoimmune encephalomyelitis by ghrelin. *J Immunol.* 2009 Aug 15;183(4):2859–66.
25. Kodama T, Ashitani J-I, Matsumoto N, Kangawa K, Nakazato M. Ghrelin treatment suppresses neutrophil-dominant inflammation in airways of patients with chronic respiratory infection. *Pulm Pharmacol Ther.* 2008 Oct;21(5):774–9.
26. Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature.* 1999 Dec 9;402(6762):656–60.
27. Osawa H. Ghrelin and *Helicobacter pylori* infection. *World J Gastroenterol.* 2008 Nov 7;14(41):6327–33.
28. Marshall BJ, Goodwin CS, Warren JR, Murray R, Blincow ED, Blackbourn SJ, et al. Prospective double-blind trial of duodenal ulcer relapse after eradication of *Campylobacter pylori*. *Lancet.* 1988 Dec 24;2(8626–8627):1437–42.
29. Uemura N, Okamoto S, Yamamoto S, Matsumura N, Yamaguchi S, Yamakido M, et al. *Helicobacter pylori* infection and the development of gastric cancer. *N Engl J Med.* 2001 Sep 13;345(11):784–9.
30. Wotherspoon AC, Doglioni C, Diss TC, Pan L, Moschini A, de Boni M, et al. Regression of primary low-grade B-cell gastric lymphoma of mucosa-associated lymphoid tissue type after eradication of *Helicobacter pylori*. *Lancet.* 1993 Sep 4;342(8871):575–7.
31. Shiiya T, Nakazato M, Mizuta M, Date Y, Mondal MS, Tanaka M, et al. Plasma ghrelin levels in lean and obese humans and the effect of glucose on ghrelin secretion. *J Clin Endocrinol Metab.* 2002 Jan;87(1):240–4.
32. Tolle V, Kadem M, Bluet-Pajot M-T, Frere D, Foulon C, Bossu C, et al. Balance in ghrelin and leptin plasma levels in anorexia nervosa patients and constitutionally thin women. *J Clin Endocrinol Metab.* 2003 Jan;88(1):109–16.
33. Wu M-S, Lee W-J, Wang H-H, Huang S-P, Lin J-T. A case-control study of association of *Helicobacter pylori* infection with morbid obesity in Taiwan. *Arch Intern Med.* 2005 Jul 11;165(13):1552–5.
34. Furuta T, Shirai N, Xiao F, Takashima M, Hanai H. Effect of *Helicobacter pylori* infection and its eradication on nutrition. *Aliment Pharmacol Ther.* 2002 Apr;16(4):799–806.
35. Vakil N, Talley NJ, Stolte M, Sundin M, Junghard O, Bolling-Sternevald E. Patterns of gastritis and the effect of eradicating *Helicobacter pylori* on gastro-oesophageal reflux disease in Western patients with non-ulcer dyspepsia. *Aliment Pharmacol Ther.* 2006 Jul 1;24(1):55–63.

36. Imagawa S, Yoshihara M, Ito M, Yoshida S, Wada Y, Tatsugami M, et al. Evaluation of gastric cancer risk using topography of histological gastritis: a large-scaled cross-sectional study. *Dig Dis Sci*. 2008 Jul;53(7):1818–23.
37. Kuipers EJ, Uytterlinde AM, Peña AS, Hazenberg HJ, Bloemena E, Lindeman J, et al. Increase of *Helicobacter pylori*-associated corpus gastritis during acid suppressive therapy: implications for long-term safety. *Am J Gastroenterol*. 1995 Sep;90(9):1401–6.
38. Bordi C, Ravazzola M. Endocrine cells in the intestinal metaplasia of gastric mucosa. *Am J Pathol*. 1979 Aug;96(2):391–8.
39. Maciorkowska E, Panasiuk A, Kondej-Muszyńska K, Kaczmarek M, Kemon A. Mucosal gastrin cells and serum gastrin levels in children with *Helicobacter pylori* infection. *Adv Med Sci*. 2006;51:137–41.
40. Dzierzanowska-Fangrat K, Michalkiewicz J, Cielecka-Kuszyk J, Nowak M, Celinska-Cedro D, Rozynek E, et al. Enhanced gastric IL-18 mRNA expression in *Helicobacter pylori*-infected children is associated with macrophage infiltration, IL-8, and IL-1 beta mRNA expression. *Eur J Gastroenterol Hepatol*. 2008 Apr;20(4):314–9.
41. Sapieryński R, Fabisiak M, Kizerwetter-Swida M, Cywińska A. Effect of *Helicobacter* sp. infection on the number of antral gastric endocrine cells in swine. *Pol J Vet Sci*. 2007;10(2):65–70.
42. Hosoda H, Doi K, Nagaya N, Okumura H, Nakagawa E, Enomoto M, et al. Optimum collection and storage conditions for ghrelin measurements: octanoyl modification of ghrelin is rapidly hydrolyzed to desacyl ghrelin in blood samples. *Clin Chem*. 2004 Jun;50(6):1077–80.
43. Gokcel A, Gumurdulu Y, Kayaselcuk F, Serin E, Ozer B, Ozsahin AK, et al. *Helicobacter pylori* has no effect on plasma ghrelin levels. *Eur J Endocrinol*. 2003 Apr;148(4):423–6.
44. Roper J, Francois F, Shue PL, Mourad MS, Pei Z, Olivares de Perez AZ, et al. Leptin and ghrelin in relation to *Helicobacter pylori* status in adult males. *J Clin Endocrinol Metab*. 2008 Jun;93(6):2350–7.
45. Pacifico L, Anania C, Osborn JF, Ferrara E, Schiavo E, Bonamico M, et al. Long-term effects of *Helicobacter pylori* eradication on circulating ghrelin and leptin concentrations and body composition in prepubertal children. *Eur J Endocrinol*. 2008 Mar;158(3):323–32.
46. Chuang C-H, Sheu B-S, Yang H-B, Lee S-C, Kao A-W, Cheng H-C, et al. Gender difference of circulating ghrelin and leptin concentrations in chronic *Helicobacter pylori* infection. *Helicobacter*. 2009 Feb;14(1):54–60.

47. Jun DW, Lee OY, Lee YY, Choi HS, Kim TH, Yoon BC. Correlation between gastrointestinal symptoms and gastric leptin and ghrelin expression in patients with gastritis. *Dig Dis Sci*. 2007 Oct;52(10):2866–72.
48. Cindoruk M, Yetkin I, Deger SM, Karakan T, Kan E, Unal S. Influence of H pylori on plasma ghrelin in patients without atrophic gastritis. *World J Gastroenterol*. 2007 Mar 14;13(10):1595–8.
49. de Martel C, Haggerty TD, Corley DA, Vogelmann JH, Orentreich N, Parsonnet J. Serum ghrelin levels and risk of subsequent adenocarcinoma of the esophagus. *Am J Gastroenterol*. 2007 Jun;102(6):1166–72.
50. Shak JR, Roper J, Perez-Perez GI, Tseng C, Francois F, Gamagari Z, et al. The effect of laparoscopic gastric banding surgery on plasma levels of appetite-control, insulinotropic, and digestive hormones. *Obes Surg*. 2008 Sep;18(9):1089–96.
51. Uzzan B, Catheline J-M, Lagorce C, Airinei G, Bon C, Cohen R, et al. Expression of ghrelin in fundus is increased after gastric banding in morbidly obese patients. *Obes Surg*. 2007 Sep;17(9):1159–64.
52. Kawashima J, Ohno S, Sakurada T, Takabayashi H, Kudo M, Ro S, et al. Circulating acylated ghrelin level decreases in accordance with the extent of atrophic gastritis. *J Gastroenterol*. 2009;44(10):1046–54.
53. Isomoto H, Ueno H, Saenko VA, Mondal MS, Nishi Y, Kawano N, et al. Impact of *Helicobacter pylori* infection on gastric and plasma ghrelin dynamics in humans. *Am J Gastroenterol*. 2005 Aug;100(8):1711–20.
54. Plonka M, Bielanski W, Konturek SJ, Targosz A, Sliwowski Z, Dobrzanska M, et al. *Helicobacter pylori* infection and serum gastrin, ghrelin and leptin in children of Polish shepherds. *Dig Liver Dis*. 2006 Feb;38(2):91–7.
55. Osawa H, Nakazato M, Date Y, Kita H, Ohnishi H, Ueno H, et al. Impaired production of gastric ghrelin in chronic gastritis associated with *Helicobacter pylori*. *J Clin Endocrinol Metab*. 2005 Jan;90(1):10–6.
56. Alonso N, Granada ML, Salinas I, Reverter JL, Flores L, Ojanguren I, et al. Plasma ghrelin concentrations in type 1 diabetic patients with autoimmune atrophic gastritis. *Eur J Endocrinol*. 2007 Dec;157(6):763–9.
57. Salles N, Ménard A, Georges A, Salzmann M, de Ledinghen V, de Mascarel A, et al. Effects of *Helicobacter pylori* infection on gut appetite peptide (leptin, ghrelin) expression in elderly inpatients. *J Gerontol A Biol Sci Med Sci*. 2006 Nov;61(11):1144–50.
58. Gao X-Y, Kuang H-Y, Liu X-M, Duan P, Yang Y, Ma Z-B. Circulating ghrelin/obestatin ratio in subjects with *Helicobacter pylori* infection. *Nutrition*. 2009 May;25(5):506–11.

59. Isomoto H, Nishi Y, Ohnita K, Mizuta Y, Kohno S, Ueno H, et al. The Relationship between Plasma and Gastric Ghrelin Levels and Strain Diversity in *Helicobacter pylori* Virulence. *Am J Gastroenterol*. 2005 Jun;100(6):1425–7.
60. [*Helicobacter pylori* infected gastric mucosa--inflammation, atrophy and carcinogenesis] - PubMed [Internet]. [cited 2021 Aug 29]. Available from: <https://pubmed.ncbi.nlm.nih.gov/16180662/>
61. Czesnikiewicz-Guzik M, Bielanski W, Guzik TJ, Loster B, Konturek SJ. *Helicobacter pylori* in the oral cavity and its implications for gastric infection, periodontal health, immunology and dyspepsia. *J Physiol Pharmacol*. 2005 Dec;56 Suppl 6:77–89.
62. Konturek PC, Cześnikiewicz-Guzik M, Bielanski W, Konturek SJ. Involvement of *Helicobacter pylori* infection in neuro-hormonal control of food intake. *J Physiol Pharmacol*. 2006 Nov;57 Suppl 5:67–81.
63. Isomoto H, Nakazato M, Ueno H, Date Y, Nishi Y, Mukae H, et al. Low plasma ghrelin levels in patients with *Helicobacter pylori*-associated gastritis. *Am J Med*. 2004 Sep 15;117(6):429–32.
64. D’Onghia V, Leoncini R, Carli R, Santoro A, Giglioni S, Sorbellini F, et al. Circulating gastrin and ghrelin levels in patients with colorectal cancer: correlation with tumour stage, *Helicobacter pylori* infection and BMI. *Biomed Pharmacother*. 2007 Apr;61(2–3):137–41.
65. Isomoto H, Ueno H, Nishi Y, Wen C-Y, Nakazato M, Kohno S. Impact of *Helicobacter pylori* infection on ghrelin and various neuroendocrine hormones in plasma. *World J Gastroenterol*. 2005 Mar 21;11(11):1644–8.
66. Campana D, Nori F, Pagotto U, De Iasio R, Morselli-Labate AM, Pasquali R, et al. Plasma acylated ghrelin levels are higher in patients with chronic atrophic gastritis. *Clin Endocrinol (Oxf)*. 2007 Nov;67(5):761–6.
67. Suzuki H, Masaoka T, Hosoda H, Nomura S, Ohara T, Kangawa K, et al. Plasma ghrelin concentration correlates with the levels of serum pepsinogen I and pepsinogen I/II ratio--a possible novel and non-invasive marker for gastric atrophy. *Hepatogastroenterology*. 2004 Oct;51(59):1249–54.
68. Khatib, Mahalaqua Nazli, Abhay Gaidhane, Shilpa Gaidhane, and Zahiruddin Quazi Syed. “Ghrelin as a Promising Therapeutic Option for Cancer Cachexia.” *CELLULAR PHYSIOLOGY AND BIOCHEMISTRY* 48, no. 5 (2018): 2172–88. <https://doi.org/10.1159/000492559>.
69. Khatib, Mahalaqua Nazli, Anuraj H. Shankar, Richard Kirubakaran, Abhay Gaidhane, Shilpa Gaidhane, Padam Simkhada, and Zahiruddin Quazi Syed. “Ghrelin for the Management of Cachexia Associated with Cancer.” *COCHRANE DATABASE OF SYSTEMATIC REVIEWS*, no. 2 (2018). <https://doi.org/10.1002/14651858.CD012229.pub2>.

UNDER PEER REVIEW