

BMI INDEX AND ABO BLOOD GROUPING INFLUENCE ON PAIN PERCEPTION : A SYSTEMATIC REVIEW

Abstract

Establishment on The stream exact review considered investigation dispersed inside the 10 years going before June 2020, dealing with the subject of obesity and pain and abo blood groups. Inside the setting of the perplexing natural and social interrelationships among these marvels, we hoped to perceive openings in the composition and to include key concentrations for future transdisciplinary research. The overall thought norms were that the included examinations could directly add to our understanding of these perplexing research.

We glanced through PubMed/Medline/Cochrane/science direct/Embase informational indexes returning 10 years, using the fundamental chase terms "weight" and "pain,abo blood grouping," and for a discretionary request we used the pursuit terms "pain" and "abo blood,obesity."

Results Included assessments (n= 90) are in a general sense human; regardless, some animal considers were joined to further develop appreciation of related fundamental natural wonders or possibly where human data were missing or inside and out limited.

Key words : ABO blood group, perception, pain, bmi index, obesity

Introduction:

“Different blood grouping might have their own qualities character characteristics. For example, individuals with blood bunch A will in general be more yearning, anxious and are alluded to as type A character. The ABO blood classifications are likewise observed to be related with different ailments, like cardiovascular infections, type 2 diabetes, and different

malignancies. Critical relationship of ABO bunches with the pervasiveness of toxemia has likewise been accounted for. We needed to analyze if ABO blood bunch has any impact on torment insight in sound volunteers initiated by chilly pressor test”[1].

“Few studies with blood group B are found to have most elevated pain perceptive and resistance while blood group AB showed least agony and blood group A showed least pain resilience. Extraordinary pain is described as an 'unpalatable, bewildering, dynamic psychophysiological response to tissue damage, disease or inflammation”[2,3] and “often continues for a short period. Its natural limit is to restrict rehearses that may incite diminished tissue healing”[4,5].”The ABO blood packs are polymorphic, gained, antigenic sugar present outwardly of erythrocytes and other tissues”[6].

“This structure is huge especially at the hour of blood holding since they can convey genuine reaction as antigens”.[7,8] “On the establishments of antigen present outwardly of erythrocyte, it might be amassed into four sorts: A, B, AB and O. The ABO blood antigens are encoded by genetic locus arranged on chromosome 9 at 9q34.1–q34.2. It has three allelic designs: A, B, and O. The A allele encodes a glycosyltransferase-A which puts a N-acetyl galactosamine to a H-antigen and designs A antigen while the B allele encodes a glycosyltransferase-B which moves D-galactose to a H-antigen and produces B antigen. The O allele encodes an impetus with no known limit and appropriately the H antigen stays unaltered and no antigens are produced”[8,9]. “The presence or nonappearance of antigen A or antigen B in red platelets will define the blood group an individual will have. In case antigen A or antigen B is accessible in the red platelets, the individual presents with blood A and B independently, however on the off chance that both antigen A and antigen B are accessible or missing in the red platelets, then the individual will have blood AB and O independently. ABO blood are conflicting circled at both area and overall scale”[10].

Agarwal et al[11] have done an examination of ABO blood arrangements and Rh (D) on 10,000 strong blood supporters giving in blood banks orchestrated in five assorted geological regions (North, South, East and Center) of India. Their assessment showed that O was the most broadly perceived blood pack (37.12%) in the country immovably followed by B at 32.26%, followed by A at 22.88% while AB was the most unescapable grouped at 7.74%. Their examination furthermore uncovered that 94.61% of the provider people was Rh positive and the rest were Rh adverse.

“Past assessment among the north Indian blood sponsor people has shown that the typical blood packs masterminded by repeat were B, O, A, and AB”[11]. “Study drove among the Medical studies in Assam showed that blood bundle A was the most broadly perceived followed by B, O and AB with 97.62% understudies having Rh positive”[12] Singh et al[13] finished an examination on the allocation of ABO and Rh (D) blood packs among Mao Naga group of Mao, Manipur, India. Out of the 775 people (359 people and 431 females), 354 individuals (46.2%) were found to have blood bundle O and shows the most significant allele repeat of 0.60 followed by 185 individuals having blood pack 'A' (23.9%) with a looking at allele repeat worth of 0.22 and 139 individuals with blood pack 'B' (17.9%) with a relating worth of 0.18 while the abundance 93 individuals were seen to have blood pack AB.

“Two research studies have concluded the connection between blood group and test incited pain. The principle assessment was done among the clinical studies and uncovered no colossal differentiation between blood arrangements and preliminary incited desolation in strong volunteers”[12].”The resulting examination was done among the strong volunteers and found that the blood pack B have the most insignificant mechanical exacerbation affectability while blood bundle AB have the highest”[13]. “Therefore, the place of this assessment was to dissect if ABO blood characterization has any impact on pain perception in solid volunteers actuated by crisp pressor test.

Both exceptional and steady pain unfavourably influence various pieces of real working. They sway the volume of real direct just as the multifaceted nature of genuine advancements” [14,15]. “The inescapability of consistent torture is 10% generally around 100 million Americans experience the impacts of steady pain” [16]. “Meanwhile, the transcendence of distress is 33% in individuals with obesity or can say with high bmi . The normality of solidified anguish with beefiness is impressive and weight reduction can chip away yet not improve steady pain perception” [17]. “Robustness related consistent pain recalls certain stress for joints and other musculoskeletal” [18]. “people with high bmi or overweight are typical in people with persevering pain” [19]. “Together, these propose an expected relating sway yet fairly very few studies consider weight and anguish altogether equivalent to such interrelationships. Also, extra verification in like manner proposes people with robustness are more misery sensitive” [20], which may make the general population or normal BMI people even more helpfully affected by exceptional pain sensation. In this way adding to the complexity of the united effects of pain and BMI. High BMI abatement can additionally foster determined pain stays an overall shortfall of examination concerning this subject all things considered. Also, colossal openings in the assessment composing (overviews and clinical primers) to the extent planning the likely friendly, healthy, and physiological covers between these two complex wonders are self-evident.

Methodology :

Study design : Systematic Review of Literature

We went through a clear marathon in search of related information and databases and published papers and chapters which were useful for our study, firstly with the help of following keywords we searched ,included keywords were :

Pain perception,

Obesity

BMI (body mass index)

ABO blood grouping

We downloaded the papers that included following papers we gone through the abstracts, if the data was enough related we searched the concerned data bases and took the whole paper, if could find the enough information in the abstracts we gone through the whole papers and also considered the synonyms of the key words and searched ,we also included few animal studies as there were no considerable evidences to include a total collection of 90 studies were found, in which 74 papers were included other papers were removed as it did have the relativity of the subject for our study.

Table 1 : Literature review databases

DATA BASES	NO OF PAPERS
PUBMED	54
EMBASE	12
COCHRANE	5
SCIENCE DIRECT	10
MEDLINE	9
TOTAL NO PAPERS INCLUDED	74
TOTAL NO	16

OF PAPERS	
EXCLUDED	

Discussion :

“BMI associates with consistent pain perceived by a couple of parts recalling a mechanical impedance from outrageous burden for skeletal muscle and joints and an extended proficient combustible status” [21]. “The composing suggests cause-and effect associations among bmi and pain are not yet clear. Caldwell et al. chosen individuals from a multidisciplinary center to dissect the relationship among BMI and pain” [22]. “Height and weight were taken from clinical records and various outcomes were assessed by studies. Mean age of 183 individuals was 38.2 years and 63% of them were women. Among this clinical people, 21% of the individuals were overweight and 30% had strength. Individuals experienced a typical of 60.0 ± 82.4 significant stretches of desolation and the larger part point by point leg/foot pain(62.7%) and hip pain (53.8%). Results showed that leg and foot pain were by and large associated with BMI in various regions (e.g., hip , spine , and back) were not related. Additionally, BMI was connected with perception of pain”.

“A cross-sectional assessment chose more than 3600 individuals to investigate the association among moderate and outrageous self-uncovered pain and bmi in southwestern US” [23]. In this general population, 36.4% adults experienced distress in some action month to month, in which 18% uncovered moderate pain and 6.6% had more pain. Backslide assessment showed colossal associations between extended BMI and extended report of moderate and outrageous pain. Furthermore, adults with power had basically higher risk of encountering moderate and outrageous misery appear differently in relation to people in the common weight and

underweight BMI classes. While 21% of people with underweight and normal weight uncovered moderate or outrageous pain, more than 30% of people with power definite moderate , and more than 40% of people with BMI 35–39.9 kg/m² reported moderate or genuine distress. Furthermore, the association among torture and BMI remained in the wake of adjusting to age, race, sex, and guidance. These examinations suggest that comorbidity of desolation and heaviness exists both in the general and torture treatment searching for masses and is likely free mature enough and a couple of other anticipated contributing segments.

“Another social class based assessment done by Wright et al. similarly showed an association among misery and strength in twins” [24]. Individuals were chosen from a twin library. Zygoty, continuous misery appearances, BMI status, bitterness, and sociodemographic factors were obtained through surveys. Among the 3471 individuals, 24.3% were overweight and 13.3% had weight. Results showed basic connection between continuous unlimited misery and robustness in the whole model (chances extent of enormous refrain common weight: 3.02; 95% CI: 2.14, 4.25). In within pair examination, connection between consistent misery and rotundity decreased resulting to adjusting to familial components (genetic and normal components), yet it remained tremendous (chances extent of huge refrain ordinary weight: 2.35; 95% CI: 1.14, 4.87). In controlling for the innate effects, this assessment had the choice to show that secret frameworks of comorbidity of torture and weight are conceivable a blend of strength express segments including an intriguing genetic tendency.

“Emery et al. guided an observational assessment to dissect if thin down quality is a mediator in the association among weight and torture” [25]. They enrolled 50 people with obesity and 50 people without heaviness. The mean BMI of individuals was 30.4 ± 7.8 kg/m² anyway it went from 18.2 to 53.3 kg/m². Body weight, height, and body piece were assessed, and one 24-h dietary audit was gotten from each subject through dietary gathering. D assessed through Healthy Eating Index-2010 (HEI-2010) using data from 24-h dietary audits. Self-

definite distress and mental factors were assessed through overviews. Results showed that more unmistakable considerable exacerbation was connected with higher BMI yet not related to sex or ethnicity. Ensuing to adding diet quality (i.e., HEI-2010) as the judge, the relationship among BMI and genuine misery became non-basic in showing assessments. Furthermore, HEI-2010 remained as an immense go between in the wake of adjusting to joint torment investigate, joint torture, utilization of anguish medication, mental hopelessness, and age. Muscle versus fat and waste border were used as markers of significant torture in two separate models which consolidated the identical covariates referred to above, and HEI2010 remained as a colossal centre individual in the association among BMI and genuine torture. To perceive which arrangements of food adding to HEI-2010 as a center individual in the association among BMI and significant distress, further relationship examinations were coordinated. The results showed that use of fish and plant protein were responsible for the intercession effect of HEI-2010 ($P= 0.005$), anyway no other dietary parts were recognized as judges of the association among BMI and genuine torture. While confined without assistance from any other person report dietary recording, this assessment suggests that diet quality, may be a huge idea in encouraging an all out perception of the association among BMI and considerable desolation.

“ABO blood grouping examination was coordinated among 80 first year clinical understudies in the age social occasion of 18-22 years. The typical BMI of the understudies goes from 18 to 21 Kg/m². Most outrageous number of the understudies were found to have blood pack B and blood bundle O (33.8 % each), followed by blood bundle A (26.3 %). Least number of understudies have blood pack AB (6.3 %). Similar disclosures were also reported by Agarwal et al¹⁰ India which showed that O was the most notable blood pack in with AB as the most un-unavoidable social occasion among 10,000 sound blood suppliers of North, south, east and central India. Past assessment coordinated among 775 Mao Naga family of

Mao, Manipur, North-East India found that the most notable blood bundle was O followed by A, B and AB. Another examination done among Assamese clinical understudies moreover showed that blood pack A was the most broadly perceived among the examination bundle, followed by B, O and AB"[25].

"Our study found that blood bundle O and B are the commonest and AB, the most different among the understudies. Globally, course of blood pack is conflicting among different population"[26].Moreover, our assessment was driven among a specific social event of subjects explicitly first year clinical understudies. These components possibly may have influenced the seen speed of ordinariness of different blood groups in our examination.

"Our examinations showed that both the typical disturbance edge and ordinary pain opposition were higher in folks when stood out from females which were really basic too. Like our revelations, an examination by Dixon et al²⁰ among patient get-togethers and sound controls showed higher misery affectability in females than in folks. Another assessment moreover definite that women were on a very basic level more delicate to cold desolation, moderate , and to ischemic desolation than men"[27]. Their result in like manner showed that women are more sensitive to a collection of hurtful lifts than men and month to month cycle stage doesn't appear to coordinate those differentiations in strong individuals. Regardless, in an assessment done by Larson et al no sex contrasts in torture affectability during CPT was taken note"[28].

"The specific covered up frameworks of sex contrasts in pain are yet to be set up. A cooperation of regular, mental and socio-social parts may be at risk for the uniqueness in torture insight among folks and females. The effect of sex synthetic compounds are acknowledged to be a basic reason for anguish related change saw among folks and females"[29]. "Effects of estradiol and progesterone on pain affectability are relative

multifaceted, both being great for nociceptive and against nociceptive while testosterone has all the reserves of being more adversary of nociceptive and protective in nature maintained by the confirmation of connection between decreased androgen center and consistent pain”[30,31]. It has been shown in an examination that women with high estradiol shows reduced torture affectability and extended endogenous opiate neurotransmission while the low estradiol is connected with lessened endogenous opiate neurotransmission.

“In our examination, blood B showed higher paincutoff or lower pain affectability and torture versatility among the different blood social events while blood AB have the least exacerbation edge and blood bundle A have the least irritation opposition. Similar revelations have been represented by Simoni et al. It was done up from their examination that blood b B showed most negligible anguish affectability with blood pack AB showing an inclination towards extended general pain affectability. Regardless, an assessment by Shankar et al saw a tendency of blood B showing a non-gigantic most diminished irritation cutoff and torture obstruction. In another examination done by Lausten et al in patients who had gone through reconstructive knee operation, there was no basic differentiation between post-employable misery assessed by the proportion of post-usable analgesics ate up and blood orders A, B and O”[32-35].

“Study drove by Maram at el found that there is an immense diminishing in fibromyalgia incidental effects including torture scores (VAS score) with low "fermentable oligo-, di-or mono-saccharides and polyols (FODMAP) diet. Past examination in rodents moreover suggested that High oily eating regimens, rather than strength all things considered, extended desolation behaviors. All the above revelations are suggestive of a connection between BMI

types and pain insight among the different blood groups , anyway further investigation is required. Another possible speculation is the incorporation of compound Glycosyltransferases in blend of ABO blood which may add to different nociceptive models as the glycosyltransferases are related with nerve myelination”[36-40].

Conclusion :

This study summarises that concerning pain perception and abo blood grouping and BMI index, there stays an complexity on the pain perception may play along BMI abo blood grouping and the vice-versa. Limited assessment has shown that BMI is decidedly associated with pain in the general population searching for tenacious torture the board everybody. AB Blood group has more pain tolerance,compared to blood group A,B,O blood group B has least pain tolerance.

Limitations :

Our study was confined to the a bounded variables, in depth of variables like genetics affecting the BMI ,gender in perceiving pain perception blood groups in different genders weren't explored much and would be grateful to see in future studies .

REFERENCES :

- [1] ChapmanCR,VierckCJ.The transition of acute postoperative pain to chronic pain: an integrative overview of research on mechanisms. J Pain 2017;18(4):359.e1-e38.
- [2] MeeksNM,GlassJS,CarrollBT. Acute pain management in dermatology: mechanisms and pathways. J Am Acad Dermatol 2015;73(4):533-40.
- [3] International Association for the Study of Pain (IASP). Taxonomy: Pain. 2017. www.iasp-pain.org

pain.org/Taxonomy?navItemNumber=576#Pain(accessed7December2017).

- [4] Brennan TJ. Pathophysiology of postoperative pain. *Pain* 2011; 152(Suppl3):S33-S40.
- [5] Henderson LA, Keay KA. Imaging acute and chronic pain in the human brain stem and spinal cord. *Neuroscientist* 2018; 24(1):84-96.
- [6] Meo SA, Rouq FA, Suraya F, et al. Association of ABO and Rh blood groups with type 2 diabetes mellitus. *Eur Rev Med Pharmacol Sci* 2016; 20(2):237-42.
- [7] Dean L. The ABO blood group. In: *Blood groups and red cell antigens*. Bethesda, MD: NCBI 2005.
- [8] Mitra R, Mishra N, Rath GP, et al. Blood group systems. *Indian J Anaesth* 2014; 58(5):524-8.
- [9] Garratty G, Glynn SA, McEntire R, et al. ABO and Rh(D) phenotype frequencies of different racial/ethnic groups in the United States. *Transfusion* 2004; 44(5):703-6.
- [10] Agrawal A, Tiwari AK, Mehta N, et al. ABO and Rh (D) group distribution and gene frequency: the first multicentric study in India. *Asian J Transfus Sci* 2014; 8(2):121-5.
- [11] Agarwal N, Thapliyal RM, Chatterjee K. Blood group phenotype frequencies in blood donors from a tertiary care hospital in north India. *Blood Res* 2013; 48(1):51-4.
- [12] Deori JP, De SK. Frequency and distribution of blood groups among medical students in a tertiary care hospital of north east India. *J Evolution Med & Dent Sci* 2016; 5(32):1704-06.
- 13. Soram JS, Panmei K, Biswas SP. Distribution of ABO and Rh (D) blood groups among the people of Mao Nagatribe of Mao, Senapati district of Manipur, North East India. *International Journal of Development Research* 2014; 4(1):184-7.
- S. Subramanian, J. M. Perkins, E. Özaltın, and G. D. Smith,

- “Weight of nations: a socioeconomic analysis of women in low- to middle-income countries,” *American Journal of Clinical Nutrition*, vol. 93, no. 2, pp. 413–421, 2011.
14. S. Subramanian, J. E. Finlay, and M. Neuman, “Global trends in body-mass index,” *The Lancet*, vol. 377, no. 9781, pp. 1915–1916, 2011.
 15. Obesity and Overweight, 2017, <http://www.who.int/mediacentre/factsheets/fs311/en/index.html>.
 16. J. A. Knight, “Diseases and disorders associated with excess body weight,” *Annals of Clinical and Laboratory Science*, vol. 41, no. 2, pp. 107–121, 2011.
 17. A. Irei, K. Takahashi, D. S. N. Le, P. Ha, and N. Hung, “Obesity is associated with increased risk of allergy in Vietnamese adolescents,” *European Journal of Clinical Nutrition*, vol. 59, no. 4, pp. 571–577, 2005.
 18. N. M. Aboel-Fetoh, A. R. Alanazi, A. S. Alanazi, and A. N. Alruwili, “A B O blood groups and risk for obesity in Al-Rar, Northern Saudi Arabia,” *Journal of the Egyptian Public Health Association*, vol. 91, no. 4, pp. 169–173, 2016.
 19. Paraschiv-Ionescu A, Perruchoud C, Buchser E, Aminian K. Barcoding human physical activity to assess chronic pain conditions. *PLoS ONE* 2012;7:e32239.
 20. Paraschiv-Ionescu A, Perruchoud C, Rutschmann B, Buchser E, Aminian K. Quantifying dimensions of physical behavior in chronic pain conditions. *J Neuroeng Rehabil*. 2016;13:85.
 21. Jackson TP, Stabile VS, McQueen KK. The global burden of chronic pain. *ASA Newsl*. 2014;78:24–27.
 22. Institute of Medicine of The National Academies. Relieving pain in America: a blueprint for transforming prevention, care, education, and research [Internet]. 2011

[cited 2017 Jan 23]. <https://www.nationalacademies.org/hmd/~media/Files/Report%20Files/2011/Relieving-Pain-in-America-A-Blueprint-for-Transforming-Prevention-Care-Education-Research/Pain%20Research%202011%20Report%20Brief.pdf>.

23. Green MA, Strong M, Razak F, Subramanian SV, Relton C, Bissell P. Who are the obese? A cluster analysis exploring subgroups of the obese. *J Public Health (Oxf)*. 2016;38(2):258–64.
24. Narouze S, Souzdalnitski D. Obesity and chronic pain: systematic review of prevalence and implications for pain practice. *Regional Anesthesia & Pain Medicine*. 2015;40:91–111.
25. Yoo JJ, Cho NH, Lim SH, Kim HA. Relationships between body mass index, fat mass, muscle mass, and musculoskeletal pain in community residents. *Arthritis Rheumatol*. 2014;66:3511–20.
26. Chai NC, Scher AI, Moghekar A, Bond DS, Peterlin BL. Obesity and headache: part I—a systematic review of the epidemiology of obesity and headache. *Headache*. 2014;54:219–34.
27. Eslick GD. Gastrointestinal symptoms and obesity: a metaanalysis. *Obes Rev*. 2012;13:469–79.
28. Gurian MB, Mitidieri AM, da Silva JB, da Silva AP, Pazin C, Poli-Neto OB, et al. Measurement of pain and anthropometric parameters in women with chronic pelvic pain. *J Eval Clin Pract*. 2015;21(1):21–27.
29. Miscio G, Guastamacchia G, Brunani A, Priano L, Baudo S, Mauro A. Obesity and peripheral neuropathy risk: a dangerous liaison. *J Peripher Nerv Syst*. 2005;10:354–8.
30. Wachholtz A, Binks M, Eisenson H, Kolotkin R, Suzuki A. Does pain predict interference with daily functioning and weight loss in an obese residential treatment-seeking population? *Int J Behav Med*. 2010;17:118–24.

31. Okifuji A, Hare BD. The association between chronic pain and obesity. *J Pain Res.* 2015;8:399–408.
32. McKendall MJ, Haier RJ. Pain sensitivity and obesity. *Psychiatry Res.* 1983;8:119–25.
33. Seaman DR. Body mass index and musculoskeletal pain: is there a connection? *Chiropr Man Therap.* 2013;21:15.
34. Janke EA, Collins A, Kozak AT. Overview of the relationship between pain and obesity: what do we know? where do we go next? *J Rehabil Res Dev.* 2007;44:245–62.
35. Arranz LI, Rafecas M, Alegre C. Effects of obesity on function and quality of life in chronic pain conditions. *Curr Rheumatol Rep.* 2014;16:390.
36. Tukker A, Visscher TL, Picavet HS. Overweight and health problems of the lower extremities: osteoarthritis, pain and disability. *Public Health Nutr.* 2009;12:359–68.
37. Darbor KE, Lench HC, Carter-Sowell AR. Do people eat the pain away? The effects of acute physical pain on subsequent consumption of sweet-tasting food. *PLoS ONE* 2016;11:e0166931.
38. Choi KW, Somers TJ, Babyak MA, Sikkema KJ, Blumenthal JA, Keefe FJ. The relationship between pain and eating among overweight and obese individuals with osteoarthritis: an ecological momentary study. *Pain Res Manag.* 2014;19:e159–163.
39. Caldwell J, Hart-Johnson T, Green CR. Body mass index and quality of life: examining blacks and whites with chronic pain. *J Pain.* 2009;10:60–67.
40. Hitt HC, McMillen RC, Thornton-Neaves T, Koch K, Cosby AG. Comorbidity of obesity and pain in a general population: results from the Southern Pain Prevalence Study. *J Pain.* 2007;8:430–6.

41. Wright LJ, Schur E, Noonan C, Ahumada S, Buchwald D, Afari N. Chronic pain, overweight, and obesity: findings from a community-based twin registry. *J Pain*. 2010;11:628–35.
42. Zahorska-Markiewicz B, Zych P, Kucio C. Pain sensitivity in obesity. *Acta Physiol Pol*. 1988;39:183–7.
43. Pradalier A, Willer JC, Boureau F, Dry J. Relationship between pain and obesity: an electrophysiological study. *Physiol Behav*. 1981;27:961–4.
44. Marks R. Obesity profiles with knee osteoarthritis: correlation with pain, disability, disease progression. *Obesity* (Silver Spring).

