

Original Research Article

Antibacterial activity of endemic *Artocarpus nobilis* Thw found in Sri Lanka

Abstract

Aims: To evaluate antibacterial activity of aqueous, methanol, dichloromethane, and hexane extracts of *Artocarpus nobilis* Thw. leaves and stem bark against *Escherichia coli* and *Staphylococcus aureus* using agar well diffusion method.

Methodology: Matured fully expanded leaves and stem bark parts of *Artocarpus nobilis* were collected, air-dried, and grounded. The extraction was obtained using a decoction extraction method. Antibacterial activity was performed against *Staphylococcus aureus* (ATCC® 25923TM) and *Escherichia coli* (ATCC® 25922TM) using agar well diffusion method and gentamicin was used as a positive control. The whole experiment was done in triplicates and the diameter of the inhibition zone (in mm) was measured and recorded.

Results: Results showed that aqueous bark extract (EC₅₀ 4.286 mg/mL) showed the highest efficacy and potency against *E. coli* while methanol bark extract (EC₅₀ 4.427 mg/mL) showed the highest efficacy and potency against *S. aureus*. R² and P values for aqueous, methanol, dichloromethane, and hexane extracts indicated that there was a strong, statistically significant correlation ($P \leq 0.05$) between concentration and zone of inhibition for all extracts of *A. nobilis* against *E. coli* and *S. aureus*.

Conclusion: This study showed that aqueous and methanol bark extracts of *Artocarpus nobilis* have marked *in vitro* dose-dependent antibacterial activity against *E. coli* and *S. aureus* respectively. Further studies are necessary to ascertain the mechanism and the active constituents responsible for the antibacterial activity of the of plant parts of *Artocarpus nobilis*.

Keywords: *Artocarpus nobilis* Thw., Antibacterial activity, ethanomedicine, Agar well diffusion method, *Escherichia coli*, *Staphylococcus aureus*, Sri Lanka.

1. INTRODUCTION

There are many types of pathogenic microorganisms such as bacteria, viruses, parasites, and fungi that cause infectious diseases. Since these infectious diseases are very common among humans, implementation of preventive measures is a timely requirement [1]. The discovery of anti-microbials is the best preventive measure that was revealed in the last century. Anti-microbial agents inhibit the growth of pathogenic microorganisms by altering their pathogenic mechanisms [2]. Due to poor administration of anti-microbial drugs and irrational use of antibiotics, microorganisms have started to develop resistant mechanisms against anti-microbial agents for their survival. As a consequence, the effectiveness of anti-

microbial agents towards these pathogens got reduced. At the same time this has become a global health issue, scientists have started to investigate and evaluate the effectiveness of many medicinal plants to discover new anti-microbial agents [3 - 8].

Plants are known to be rich sources of anti-microbial agents since ancient time [9]. In most of the studies, it has proven that compounds derived from various parts of the plants have a promising therapeutic effect mainly due to the phytochemistry of the tree [10]. These compounds are very beneficial in the treatment of infectious diseases [11]. Out of many medicinal plants, *Artocarpus nobilis* Thw. was selected for the current study. *Artocarpus nobilis* belongs to the family *Moraceae* with 40 genera and 60 species [12, 41]. It is endemic to Sri Lanka, and it is economically and medicinally significant tree [13, 18]. *Artocarpus nobilis* Thw. is often found in the wet zone, mid-country homesteads, and wet zone of Sri Lanka [14]. It is known in Sinhala as "Wal Del," "Badi Del," in English as "Ceylon wild breadfruit" and in Tamil as "Aresini-pilaka", "Asiri-pillakai" [15]. It is well-known for its anti-helminthic and antibacterial activities. *Artocarpus nobilis* also used in folk and ayurvedic medicine in Sri Lanka. Edible fruits and seeds are high in nutrients. Bark is used for dysentery and muscle strain, bark and latex combinations used for abscesses and blisters. According to the literature *A. nobilis* is a rich source of anti-microbial and anti-inflammatory compounds but most of these properties have not been scientifically explored. Previous studies on the phytochemistry of the genus *Artocarpus* has led to identification of many compounds like flavonoids, geranyl chalcone derivatives, geranylated phenolic constituents, stilbene derivatives, xanthenes, triterpenes and cycloartane-type triterpenoids carbohydrates, protein, fibre, different ions, vitamin C, thiamine, riboflavin, niacin, vitamin A, lutein and β -carotene etc. [16]. According to the folk and Ayurvedic literature, texture of soil, amount of rain, the average temperature of a particular area would alter the quality and other aspects of a tree have been mentioned [17]. Therefore, Sri Lankan varieties of *A. nobilis* may also have variations in therapeutic properties, quality, and quantity of phytoconstituents according to the area where the plant is grown. In addition to phytochemistry, pharmacological properties such as anti-inflammatory, radical scavenging, antioxidant activity, fungicidal ability [18, 19, 20, 7, 8] and acetylcholinesterase inhibitory activity have also been investigated earlier [21].

Artocarpus nobilis has many anti-microbial properties and this study mainly focused on its antibacterial activity against intestinal infections and wound infections. Enteric (intestinal) infection is a very common condition that refers to the infection in Gastrointestinal system caused by microorganisms such as viruses (Norovirus, Rotavirus), bacteria (*E. coli*, *Salmonella*, *Shigella*, *Campylobacter*, *Clostridium*), and parasites (*Giardia*, *Entamoeba*, *Ascaris*) [22]. These diseases most frequently result from consuming contaminated food or water, and some can spread from person to person. Most of the time infections of the intestines result in diarrhoea or dysentery, nausea, vomiting, and abdominal cramping [23]. According to the World Health Organization [24] *Escherichia coli*, is the commonest causative agent of moderate to severe diarrhoea and food poisoning in developing and low-income countries. *Escherichia coli* is a Gram-negative bacterium and they are mammalian intestinal commensals as well as pathogens [25]. Based on the above information, *Escherichia coli* strain was used to investigate the antibacterial effect of *A. nobilis* against gastrointestinal infections. When considering the wound infection, it is one of the most common hospital-acquired infections. The International Wound Infection Institute [26] defines wound infection as 'the invasion of a wound by proliferating microorganisms to a level that causes a local and/or systemic response in the host. In almost all cases wounds are colonized with potential pathogens such as *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and beta-haemolytic *Streptococci* [27]. Among these pathogens *S. aureus* is the most commonly identified pathogen responsible for wound abscesses, skin, and soft tissue infections [28]. Accordingly, to evaluate the antibacterial on wound infections, this study was

selected *Staphylococcus aureus*. *In-vitro* antibacterial activity against these two organisms was evaluated using agar well diffusion method.

2. METHODOLOGY

2.1 Plant material collection, identification, and authentication

Well grown and fully expanded fresh leaves and bark of *Artocarpus nobilis* Thw. (Bedi del / Wal del) about 500g of each were collected at flowering season from an estate in Gampaha district in Western Province, Sri Lanka (Latitude of 7° 23' 59.99" N and Longitude of 79° 08' 59.99"). Plants which were collected were identified and authenticated by a Botanist at National Herbarium, Peradeniya, Sri Lanka. Organoleptic properties such as colour, odour, and morphological characteristics such as appearance, shape, size of the *Artocarpus nobilis* Thw. (Bedi del / Wal del) were studied carefully before collection of above plant material.

2.2 Preparation of aqueous, methanol, dichloromethane, and hexane extracts of *Artocarpus nobilis* Thw. (Bedi del / Wal del) leaves and bark

Using running tap water, selected plant materials were thoroughly washed and air dried until a constant weight was obtained. The dried leaves and bark of *Artocarpus nobilis* Thw. were grounded using a grinder to obtain a fine powder material. For the extraction process, well dried and blended powder samples of each plant material were taken. The extraction was obtained using a decoction extraction method. Using an electronic balance 50 g of the fine powder was weighed and added to 500 ml of solvent measured using a measuring cylinder. This was added to the decoction apparatus then it was boiled slowly for 4 hours. The prepared extract was left for cooling and then was concentrated from the rotary vacuum evaporator into sterile glass vials. The final concentrated solid mass of the extracts was stored at 4°C in the freezer compartment of the refrigerator until used for the experiment.

2.3 Preparation of plant extracts standard concentration and positive control

Serial dilutions yielding concentrations of 1.25, 2.5, 5, 10, 20, 40, 60, 80 and 100 mg/mL were prepared from the concentrated extracts (aqueous, methanol, dichloromethane, and hexane) [29]. Briefly, concentrated extracts from in powder form first dissolved in DMSO and later used Milli-Q water for preparation of serial dilutions from the stock. Gentamicin was used as the positive control of the study. It was prepared by using commercially available IV Gentamicin (80 mg/2 ml) vial. 20 mg/ml of Gentamicin solution was prepared as the final concentration of the positive control.

2.4 Anti-bacterial activity screening using Agar well diffusion method

Antibacterial activity was performed against Gram positive *Staphylococcus aureus* (ATCC® 25923TM) and Gram-negative *Escherichia coli* (ATCC® 25922TM) bacteria by using agar well diffusion method [30, 23]. Bacterial suspension was prepared according to the 0.5 McFarland turbidity [31]. A sterile swab was dipped into the inoculum and each of the test organisms were spread on the separate Mueller Hinton Agar (MHA) plates (100 x 15 mm in size) by using sterile swabs to obtain a confluent growth [2, 30]. After that sterile stainless-steel cylinders were used to make the wells of 6 mm in diameter, 2.5mm deep and 2 cm apart from the wall and each well on the plate [3]. Then Using the sterile inoculation needle, wells were opened and sealed the wells with sterile, melted MHA agar 10µl. Then using micropipette, 100 µL of each diluted extract solution was added into opened wells [32].

Gentamicin was used as positive control (C1) [33]. Sterile DMSO was used as negative control (C2) [34]. Plates were incubated for 24 hours at 37 °C. After overnight incubation, each of the inhibition zone diameters for the extracts against *Staphylococcus aureus* and *Escherichia coli* were measured and recorded in millimeters. The whole experiment was done in triplicates [3].

3. RESULTS AND DISCUSSION

Reports on anti-bacterial activity of *A. nobilis* are limited and many of these studies have conducted on the phytochemistry and few just to find out the zone of inhibition against bacterial pathogens. However, these studies have not investigated the efficacy, potency, and dose response relationship. Current study investigated *in vitro* antibacterial activity and dose response of aqueous, methanol, dichloromethane and hexane extracts of leaves and bark of *A. nobilis* which is only found in Sri Lanka.




Fig. 1.: The antibacterial activity testing results of dichloromethane bark extracts of *Artocarpus nobilis* against *S. aureus* (top panel) and Methanol Leaves extract against *E. coli* (bottom panel) is shown here. C1 is the positive control and C2 negative control.

According to the results obtained (Figure 1), dichloromethane bark extract of *A. nobilis* showed potent antibacterial activity against *E.coli* at concentration of 100 mg/ml with an inhibition zone diameter of 22.33 ± 0.33 mm. On the other hand, methanol bark extract of *A.*

nobilis showed potent antibacterial activity against *S. aureus* at concentration of 100 mg/ml with an inhibition zone diameter of 25.66 ± 0.33 mm. EC_{50} or the 'half maximum effective concentration is the drug concentration required to provide the half of the maximum possible effect. EC_{50} values were obtained using Graphpad Prism 8 (version 8.2.1). As shown in table 1, EC_{50} values of leaves extracts against *E. coli* as follows. Hexane (4.398 mg/mL) > methanol (4.608 mg/mL) > aqueous (8.50 mg/mL) > dichloromethane (29.96 mg/mL). Antibacterial activity of bark extracts against *E. coli* aqueous (4.286 mg/mL) > methanol (4.976 mg/mL) > dichloromethane (5.959 mg/mL) > hexane (8.95 mg/mL). When comparing both leaves and bark extracts, aqueous bark extract (4.286 mg/mL) showed the highest potency. According to table 2, Antibacterial activity of leaves extracts against *S. aureus* methanol (6.101mg/mL) > aqueous (13.26 mg/mL) > dichloromethane (13.5 mg/mL) > hexane (29.36 mg/mL). Antibacterial activity of bark extracts against *S. aureus* methanol (4.427 mg/mL) > dichloromethane (5.373 mg/mL) > hexane (32.92 mg/mL) > aqueous (65.71 mg/mL). When comparing both leaves and bark extracts, methanol bark extract (4.286 mg/mL) showed the highest potency.

According to results in table (1) and dose-response curve (Figure 2), aqueous bark extract has the highest antibacterial efficacy and aqueous leaves extract has the lowest antibacterial efficacy against *E. coli*. On the other hand, results in table (2) and dose-response curve (Figure 3) shows, methanol bark extract has the highest antibacterial efficacy and hexane bark extract has the lowest antibacterial efficacy against *S. aureus*. According to the R^2 and P values shown in table (1, 2) and (1, 2) there was a strong positive statistically significant correlation ($P = 0.05$) between concentration and zone of inhibition by all the extracts of plant *A. nobilis* against *E. coli* and *S. aureus*. P values and R^2 values were obtained using Graphpad Prism 8 (version 8.2.1).

According to the literature of folk and ayurvedic medicine, *A. nobilis* has antibacterial activity against gastrointestinal infections as well as it is used to treat wounds and blisters. To evaluate these properties, Gram-negative *E. coli* (ATCC 25922) and Gram-positive *S. aureus* (ATCC 25923) were selected respectively. Antibacterial activity was evaluated using agar well diffusion method [35, 24]. Generally, for antibacterial investigations of plant extracts methanol and aqueous extracts are considered as the best. Therefore, we have used aqueous and methanol extracts along with dichloromethane and hexane because solubility of active compounds differs with the solvent polarity [36]. The presence of phytoconstituents and secondary metabolites in the extracts, the method used for the extraction (decoction method) and the types of solvents used for the extraction process might contribute immensely for obtaining this positive antibacterial activity against *E. coli* and *S. aureus*. Extracts were rich in phytochemicals which regulate the antibacterial activity including xanthone, xanthoangelol, terpenoids, stilbene, phenols, flavonoids, and flavone [37, 11-15]. Flavonoids act against bacteria utilizing several mechanisms. Inhibition of nucleic acid synthesis, inhibition of cytoplasmic membrane function, inhibition of the porin on the cell membrane and inhibition of the porin on the cell membrane are some well-known examples [38]. Xanthones and xanthone derivatives can serve as antibacterial and antioxidants by disrupting bacterial cell membranes. It induces the release of lipoteichoic acid (LTA) from the cell wall of Gram-positive bacteria, which are covalently bonded to the outside of peptidoglycan that is used in bacterial cell division and osmotic defense [39] while stilbene compounds exert its antibacterial activity by interfering with the membrane stability and permeability of the bacterial cells [40]. At the same time, it is important to mention that there is a new emerging area of research where bacterial chaperones have been used as a novel antimicrobial target and development of novel anti microbials. Hence it is important to investigate the effect of these compounds against chaperone activity as well. Anti-inflammatory activity by plant extracts may be mediated through possible anti-bacterial

Table 1. Dose-response curve details for methanol, dichloromethane, and hexane leaves and bark extract samples of *A. nobilis* Thw plant parts against *E. coli*, Average values are shown, (N = 3)

Table 2. Dose-response curve details for methanol, dichloromethane, and hexane leaves and bark extract samples of *A. nobilis* Thw plant parts against *S. aureus*. Average values are shown, (N = 3)

[illegible]




Fig. 2.: Dose-response curve for different extracts of *A. nobilis* plant parts (leaves and bark) against *E. coli*. (Normalized zone of Inhibition measured in mm) Error bar represent the standard error of mean.




Fig. 3.: Dose-response curve for different extracts of *A. nobilis* plant parts (leaves and bark) against *S. aureus*. (Normalized zone of Inhibition measured in mm). Error bar represent the standard error of mean. Graphs were prepared using Graphpad Prism 8 (version 8.2.1), using non-linear regression model according to the equation: $Span = Top - Bottom$, $Y = Bottom + (Top - Bottom) / (1 + 10^{-(Log\ IC_{50} - X) * HillSlope})$, where X is the log of dose response or concentration. Y= is response

4. CONCLUSION

In conclusion, this study demonstrates, for the first time, potent *in vitro* antibacterial activity aqueous, methanol, dichloromethane, hexane extracts of leaves and bark of *Artocarpus nobilis* against *Escherichia coli* and *Staphylococcus aureus*. The aqueous bark extract showed marked antibacterial activity *E. coli* while methanol bark showed marked antibacterial activity against *S. aureus*. Further studies are necessary to determine the mechanism and active constituents responsible for the antibacterial activity of *A. nobilis*. Moreover, our result indicates a strong possibility of developing safe potent and cheap antibacterial agent from *A. nobilis* bark.

COMPETING INTERESTS DISCLAIMER:

AUTHORS HAVE DECLARED THAT NO COMPETING INTERESTS EXIST. THIS RESEARCH WAS NOT FUNDED BY ANY COMPANY OR FUNDING AGENCY.

REFERENCES

1. Muir's Textbook of Pathology, Fourteenth Edition : David Levison : 9780340740620 <https://www.bookdepository.com/Muir-Textbook-Pathology-Fourteenth-Edition-David-Levison/9780340740620> (accessed 2021 -12 -22).
2. Patel J. Antibiotic Resistance: Understanding and Responding to an Emerging Crisis. *Emerging Infectious Diseases* 2011, 17 (10), 1984. <https://doi.org/10.3201/eid1710.111066>.
3. Mbah-Omeje. *In Vitro* Study on the Anti-microbial Activity of *Curcuma Longa* Rhizome on Some Microorganism. *AJBLs* 2019, 7 (1), 1. <https://doi.org/10.11648/j.ajbls.20190701.11>.
4. Ranaweera CB, Pathirana R, Ambalanduwa KC, Jayakody RA, Ratnasooriya WD. *In vitro* antirheumatoid arthritic activity of aqueous root extract of *Clitoria ternatea*. *International research journal of pharmacy*. 2014;5(12):926-8. DOI: 10.7897/2230-8407.0572188
5. Silva, A.R.N., Ranaweera, C.B., Karunathilaka, R.N., Pathirana, R. and Ratnasooriya, W.D., 2016. Antibacterial activity of water extracts of different parts of *Morinda citrifolia* grown in Sri Lanka. *Int J Sci Res Publ*, 6, pp.124-7. <http://www.ijsrp.org/research-paper-0516/ijsrp-p5322.pdf>
6. Silva AR, Dissanayake DM, Ranaweera CB, Pathirana R, Ratnasooriya WD. Evaluation of *in vitro* antibacterial activity of some Sri Lankan medicinal plants. *Int J Pharmaceutical Res Allied Sci*. 2015; 4:54-7. http://www.journalijar.com/uploads/225_IJAR-12022.pdf
7. Nimantha Karunathilaka RD, Silva AR, Ranaweera CB, Dissanayake DM, Nelumdeniya NR, Pathirana R, Ratnasooriya WD. *In vitro* antibacterial activity of hexane, chloroform and methanolic extracts of different parts of *Acronychia pedunculata* grown in Sri Lanka. *Int. J. of Adv. Res*. 2016;4(8):1574-9. DOI: 10.21474/IJAR01/1364
8. Ranaweera, CB, Chandana AK. *Clitoria ternatea*-Shifting Paradigms: From Laboratory to Industry. *South Asian Journal of Research in Microbiology*11(2): 18-26, 2021. DOI: 10.9734/SAJRM/2021/v11i230247

9. Khatak S, Naagar J, Gupta A, Malik K. Anti-microbial Activity of *Vitex Negundo* against Pathogenic Bacteria. *Journal of Pharmacy Research* 2014, 8 (2), 91–92.
10. Quiroga N, Sampietro R, Vattuone A. Screening Antifungal Activities of Selected Medicinal Plants. *J Ethnopharmacol* 2001, 74 (1), 89–96. [https://doi.org/10.1016/s0378-8741\(00\)00350-0](https://doi.org/10.1016/s0378-8741(00)00350-0).
11. Erdogru Ö T. Antibacterial Activities of Some Plant Extracts Used in Folk Medicine. *Pharmaceutical Biology* 2002, 40 (4), 269–273. <https://doi.org/10.1076/phbi.40.4.269.8474>
12. Jayasinghe U, Puvanendran S, Hara N, Fujimoto Y. Stilbene derivatives with antifungal and radical scavenging properties from the Stem Bark of *Artocarpus nobilis*. *Natural Product Research*. 2007;18(6): 571-574.
13. Jayasinghe L, Rupasinghe G, Hara N, Fujimoto Y. Geranylated phenolic constituents from the fruits of *Artocarpus nobilis*. *Phytochemistry*. 2006;67(13): 1353-1358. DOI: [10.1016/j.phytochem.2006.04.011](https://doi.org/10.1016/j.phytochem.2006.04.011)
14. Madushani I, Fernando T, Wijesundara C, Siriwardane D. First Report of White Root Disease of *Artocarpus Nobilis* in Sri Lanka Caused by Rigidoporus Microporus. *Journal of the National Science Foundation of Sri Lanka* 2014, 42 (2), 197–198. <https://doi.org/10.4038/jnsfsr.v42i2.6998>.
15. Department of Ayurveda. Compendium of medicinal plants: A Sri Lankan study. 4th ed. Department of Ayurveda, Sri Lanka; 2004.
16. Elevitch C, Ragone D, Cole I, Breadfruit production guide: recommended practices for growing, harvesting, and handling; 2014. Available: https://hdoa.hawaii.gov/add/files/2014/05/Breadfruit_Production_Guide_web_edition.pdf
17. Charaka, A. Vimana Sthana https://www.carakasamhitaonline.com/index.php?title=Vimana_Sthana (accessed 2021 -12 -26).
18. Senadeera, S. P. N. N.; Fernando, K. S. K.; Wickramasekara, W. L. L. N.; Fernando, M. Y. S.; Ranaweera, C. B.; Rajapaksha, W.; Silva, A. R. N. *In Vitro* Anti-Inflammatory Activity of Endemic *Artocarpus Nobilis* Thw Found in Sri Lanka. *APRJ* 2021, 116–122. <https://doi.org/10.9734/aprj/2021/v8i430192>.
19. Jayasinghe L, Balasooriya B, Padmini W, Hara N, Fujimoto Y. Geranyl chalcone derivatives with antifungal and radical scavenging properties from the leaves of *Artocarpus nobilis*. *Phytochemistry*. 2004;65(9): 1287-1290. DOI: [10.1016/j.phytochem.2004.03.033](https://doi.org/10.1016/j.phytochem.2004.03.033)
20. Jayasinghe U, Samarakoon T, Kumarihamy B, Hara N, Fujimoto Y. Four new prenylated flavonoids and xanthenes from the root bark of *Artocarpus nobilis*. *Fitoterapia*. 2008;79(1): 37-41. DOI: [10.1016/j.fitote.2007.07.014](https://doi.org/10.1016/j.fitote.2007.07.014)
21. Zahid S, Ata A, Samarasekera R. New Cycloartane-Type Triterpenoids from *Artocarpus Nobilis*. *Zeitschrift für Naturforschung B* 2007, 62 (2), 280–284. <https://doi.org/10.1515/znbn-2007-0222>
22. Enteric Diseases and Food-Borne Diseases, Public Health Ontario <https://www.publichealthontario.ca/en/diseases-and-conditions/infectious-diseases/enteric-foodborne-diseases> (accessed 2021 -12 -26).
23. Gastrointestinal Infections <https://www.atsu.edu/faculty/chamberlain/website/lectures/lecture/gi4.htm> (accessed 2021 -12 -26).
24. Diarrhoeal disease <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease> (accessed 2021 -12 -26).
25. Riley W. Pandemic Lineages of Extraintestinal Pathogenic *Escherichia Coli*. *Clinical Microbiology and Infections* 2014, 20 (5), 380–390. <https://doi.org/10.1111/1469-0691.12646>.
26. Calne S. Wound infection in clinical practice, An international consensus. <https://healthservice.hse.ie/filelibrary/onmsd/hse-national-wound-management-guidelines-2018.pdf>

27. Bowler G, Duerden I, Armstrong G. Wound Microbiology and Associated Approaches to Wound Management. *Clin Microbiol Rev* 2001, 14 (2), 244–269.
<https://doi.org/10.1128/CMR.14.2.244-269.2001>.
28. McCaig F, McDonald C, Mandal S, Jernigan B. *Staphylococcus Aureus*–Associated Skin and Soft Tissue Infections in Ambulatory Care. *Emerging Infectious Diseases* 2006, 12 (11), 1715–1723. <https://doi.org/10.3201/eid1211.060190>.
29. Valgas C, Souza M, Smânia A, Smânia Jr A. Screening Methods to Determine Antibacterial Activity of Natural Products. *Braz. J. Microbiol.* 2007, 38 (2), 369–380.
<https://doi.org/10.1590/S1517-83822007000200034>.
30. Abeyasinghe D, Pathirana N, Wanigatunge P. Evaluation of Antibacterial Activity of Different Mangrove Plant Extracts. *Ruhuna Journal of Science* 2012, 1 (1).
31. CLSI. Performance standards for Anti-microbial Susceptibility Testing; Twenty-Fifth Informational Supplement. Clinical and Laboratory Standards Institute. 2015, 35(3), 1–231
32. Ali Khan M, Yaqoob S, Ahmad S. Anti-microbial Activity of *Azadirachta Indica*, against Target Pathogens and Its Utility as a Disinfectant and Floor Cleaner. *jemds* 2021, 10 (25), 1899–1903. <https://doi.org/10.14260/jemds/2021/392>.
33. Rajapaksha W, Silva N, Bulathwelage A, Wanigasekara D. Evaluation of *in Vitro* Antibacterial Activity of Extract of Semi-Aquatic Plants Growing in the Polluted Water of Sri Lanka. 2019.
<https://doi.org/10.13140/RG.2.2.19704.39689>.
34. Muddukrishnaiah K, Singh S. Anti-microbial, Synergistic Activity and Antioxidant Studies on Multidrug Resistance Human Pathogen Using Crude Extract of *Azadirachta Indica* Leaf and *Withania Somnifera* Rhizome. *Journal of Plant Pathology & Microbiology* 2015, 0 (0), 1–3.
<https://doi.org/10.4172/2157-7471.1000S3-009>.
35. Mohammadzadeh T, Sadjadi S, Habibi P, Sarkari B. Comparison of Agar Dilution, Broth Dilution, Cylinder Plate and Disk Diffusion Methods for Evaluation of Anti-Leishmanial Drugs on *Leishmania* Promastigotes. *Iran J Parasitol* 2012, 7 (3), 43–47.
36. Nawaz H, Shad A, Rehman N, Andaleeb H, Ullah N. Effect of Solvent Polarity on Extraction Yield and Antioxidant Properties of Phytochemicals from Bean (*Phaseolus Vulgaris*) Seeds. *Braz. J. Pharm. Sci.* 2020, 56. <https://doi.org/10.1590/s2175-97902019000417129>.
37. Somashekhar M, Nayeem N, Sonnad B. A Review on Family Moraceae (Mulberry) With a Focus on *Artocarpus* Species. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2013, 2, 2614–2621.
38. Xie Y, Yang W, Tang F, Chen X, Ren L. Antibacterial Activities of Flavonoids: Structure-Activity Relationship and Mechanism. *Curr Med Chem* 2015, 22 (1), 132–149.
<https://doi.org/10.2174/0929867321666140916113443>
39. Miladiyah, I.; Rachmawaty, F. J. Potency of Xanthone Derivatives as Antibacterial Agent against Methicillin-Resistant *Staphylococcus Aureus* (MRSA). *JKKI : Jurnal Kedokteran dan Kesehatan Indonesia* 2017, 8 (2), 124–135. <https://doi.org/10.20885/JKKI.Vol8.Iss2.art8>.
40. Mattio M, Catinella G, Dallavalle S, Pinto A. Stilbenoids: A Natural Arsenal against Bacterial Pathogens. *Antibiotics* 2020, 9 (6), 336. <https://doi.org/10.3390/antibiotics9060336>.
41. Park J, Lee J, Jung E, Park Y, Kim K, Park B, Jung K, Park E, Kim J, Park D. *In Vitro* Antibacterial and Anti-Inflammatory Effects of Honokiol and Magnolol against *Propionibacterium* Sp. *Eur J Pharmacol* 2004, 496 (1–3), 189–195.
<https://doi.org/10.1016/j.ejphar.2004.05.047>.
42. Hashim, N. Mohd.; Rahmani, M.; Ee, G. C. L.; Sukari, M. A.; Yahayu, M.; Amin, M. A. M.; Ali, A. M.; Go, R. Antioxidant, Anti-microbial and Tyrosinase Inhibitory Activities of Xanthones Isolated from *Artocarpus Obtusus* F.M. Jarrett. *Molecules* 2012, 17 (5), 6071–6082.
<https://doi.org/10.3390/molecules17056071>.
43. Ranaweera, Chathuranga B., Przemyslaw Glaza, and Michal Zolkiewski. "Interaction of substrate-mimicking peptides with the AAA+ ATPase ClpB from *Escherichia coli*." (2018).
<https://doi.org/10.1016/j.abb.2018.08.002>

44. Glaza, Przemyslaw, Chathurange B. Ranaweera, Sunitha Shiva, Anuradha Roy, Brian V. Geisbrecht, Frank J. Schoenen, and Michal Zolkiewski. "Repurposing p97 inhibitors for chemical modulation of the bacterial ClpB–DnaK chaperone system." *Journal of Biological Chemistry* 296 (2021). DOI: <https://doi.org/10.1074/jbc.RA120.015413>

UNDER PEER REVIEW